

Survey on the Disaster Prevention System in Rural Areas and Isolated Islands: Preparing for the Nankai Trough Earthquake

Dr. Takamitsu Kodama, Mr. Kazuhiro Ozawa,
Dr. Takashi Nakagawa

Aichi Medical University, Nagakute, Japan

Introduction: An expert government committee in Japan decided to revise the chance of a Nankai Trough earthquake in the next 30 years from 70% to between 70 and 80% in 2017. If a mega-earthquake occurs, medical institutions in disaster regions are required to perform self-contained activities during the super-acute phase. Human damage varies depending on whether medical functions can be sustained, particularly in rural areas and isolated islands. Here we examined actual situations.

Aim: To identify the issues of a disaster medical system in rural areas and isolated islands that need solving.

Methods: Regarding disaster preparedness planning, we conducted a survey on 10 hospitals undesignated as disaster key hospitals of remote area medical care bases (survey group), and 69 hospitals designated as disaster key hospitals (control group) in three prefectures in the Tokai region (Aichi, Mie, and Shizuoka).

Results: We received responses from four hospitals in the survey group and 52 hospitals in the control group. The hospitals in the survey group responded that they could accept 74 severe casualties and 85 moderate casualties. We identified problems such as insufficient stockpiling of fuel, water, and oxygen, and lack of a prioritized lifeline supply contract.

Discussion: It was predicted that human damage would be relatively minor given smaller populations in rural areas and isolated islands in the Tokai region. However, the number of patients would exceed their acceptance capacity. Moreover, the system for sustaining infrastructure is not adequate for providing medical services. Thus, it was indicated that these regions would be isolated in terms of disaster measures. It is imperative to establish a disaster medical system in rural areas and isolated islands that lack adequate disaster medical systems to manage Nankai Trough earthquakes.

Prehosp Disaster Med 2019;34(Suppl. 1):s170

doi:10.1017/S1049023X19003881

Susceptibility of Host Immunity to Infectious Diseases in the Spacecraft Environment

Miss Vienna Tran

School of Medicine, The University of Adelaide, Adelaide, Australia

Introduction: The principles of Disaster and Emergency Medicine are applicable beyond the confines of planet Earth. With the accelerating rate of climate change, natural disasters, and overpopulation, as well as the innate human appetite for knowledge and technological advancement, there has recently been an increased interest in the prospect of long-duration spaceflight with a view to colonize extra-terrestrial bodies, such as Mars. However, there is a need to understand the risk of adverse medical events in the hostile environment of space.

For example, previous incidences of infectious disease and immune dysregulation during a short-term mission have threatened to jeopardize the crew dynamic and the mission objectives. The risk of infectious diseases to the astronaut is one of the many knowledge gaps that must be addressed before long-duration flight is considered.

Aim: To review how spaceflight impacts an astronaut's in-flight susceptibility to infectious diseases.

Methods: Research was guided by the Microbiology section of the NASA Human Research Roadmap Program. Search terms in the University of Adelaide Library Search database collection included: "infectious diseases + spaceflight," "astronaut + immunity," "analog," and "inflammatory marker."

Results: Studies that have been conducted in-flight and on Earth demonstrate that both the astronaut and the microbe are affected by spaceflight. Stress, microgravity, and the isolated nature of the spacecraft have been found to compromise the immunity of the astronaut, as shown by reduced T cell counts and increased viral shedding of dormant viruses. Microbes have demonstrated rapid adaptation mechanisms, including genetic mutation and increased virulence.

Discussion: This paper identifies a significant need for further research into host immunity during spaceflight to mitigate infectious disease risk. It is recommended that in-situ studies and terrestrial space analogs are most effective and that current knowledge on the principles of wilderness and expedition medicine be applied where possible.

Prehosp Disaster Med 2019;34(Suppl. 1):s170

doi:10.1017/S1049023X19003893

The System of Disaster Dispatch Team Corresponding to the Nuclear Disaster of Nagasaki University

Dr. Yoshihiro Nozaki

Nagasaki University Hospital, Nagasaki, Japan

Introduction: After accidents of Fukushima Daiichi power plant, the Japanese Government distinguished some medical institutions corresponding to the nuclear disaster by roles and functions. Nuclear Regulation Authority is managing these medical institutions. The Nagasaki University was designated as two centers for "the advanced radiation emergency medical support" and "nuclear emergency medical support". We established "Headquarters for Nuclear Disaster Response and Preparedness in Nagasaki University" (NDRP) and prepared for emergency ordinarily. The staff of headquarters are mainly concentrating their power on the network construction and joint training with each facility. We are improving the dispatch system of nuclear emergency medical assistance team, but some problems were found through the experiences of some trainings.

Aim: To stimulate discussion and listen to opinions from several facilities.

Methods: The dispatch system of nuclear emergency medical assistance team imitated the system of Japan Disaster Medical Assistance Team (DMAT). Specifically, activity days of all teams are limited, and all teams should come under the command of the head of support acceptance medical institution of a disaster area. Particularly the main duties of the dispatch