

Changes in neuropsychological functioning following treatment for late-life generalised anxiety disorder

Meryl A. Butters, Rishi K. Bhalla, Carmen Andreescu, Julie Loebach Wetherell, Rose Mantella, Amy E. Begley and Eric J. Lenze

Background

Generalised anxiety disorder (GAD) in older adults is associated with neuropsychological impairment.

Aims

We examined neuropsychological functioning in older adults with GAD in comparison with psychiatrically healthy older adults at baseline, and we examined changes following a 12-week placebo-controlled trial of escitalopram.

Method

A total of 160 participants without dementia aged ≥ 60 with current GAD and 37 individuals in a comparison group without psychiatric history underwent neuropsychological assessment. Of these, 129 participants with GAD were reassessed post-treatment (trial registration: NCT00105586).

Results

The participants with GAD performed worse than the comparison group in information processing speed, working memory, inhibition, problem-solving (including concept formation and mental flexibility) and immediate and delayed

memory. Neuropsychological functioning was correlated with everyday functioning. After treatment, those with low cognitive scores experienced working memory, delayed memory and visuospatial ability improvement and those who reported clinical improvement in anxiety exhibited improvement in the ability to engage inhibition and episodic recall. These improvements were modest and of similar magnitude in both treatment conditions.

Conclusions

Generalised anxiety disorder in older adults is associated with neuropsychological impairments, which are associated with functional impairment. Those with GAD who either have a low cognitive performance or report clinical improvement in anxiety post-treatment, show improvement in multiple cognitive domains. These findings underscore the importance of treatments that aid cognition as well as anxiety symptoms.

Declaration of interest

R.M. is employed by Abbott Laboratories.

Generalised anxiety disorder (GAD) is a serious and chronic illness, defined by excessive and difficult-to-control worry. Somatic and mental symptoms include poor concentration, restlessness, sleep disturbance, fatigue and muscle tension. The prevalence of GAD is as high as 7.3% in community-dwelling older adults.^{1–3} Older adults with GAD worry about medical problems, family and finances; typical concerns in this age group but greatly amplified in duration, severity and distress.⁴ The disorder tends to be chronic in the absence of effective treatment, with an average duration of 20 years or more^{5–8} and strongly associated with functional disability.⁹

To our knowledge, no single study has provided a comprehensive examination of neuropsychological functioning in late-life GAD. Moreover, there is very little available data on whether neuropsychological function changes with treatment. Thus, we carried out a neuropsychological evaluation pre- and post-treatment, in a large group of older adults with GAD in a randomised placebo-controlled trial of the selective serotonin reuptake inhibitor (SSRI) escitalopram. We also compared participants with GAD (the GAD group) at baseline with a group of psychiatrically healthy older adults (comparison group), equated for gender, ethnicity, years of education and medical burden. The goal of this study was twofold: to characterise neuropsychological function among a large group of older adults with a principal diagnosis of GAD; and to identify any cognitive changes related to treatment of anxiety. Based on the published literature (including our own work) with the rationale that worry takes up cognitive capacity and leaves less attentional resources for the tasks at hand, we hypothesised that those in the GAD group

would perform worse than the comparison group on measures of attention (digit span), information processing speed (coding), and executive functions including working memory (letter–number sequencing) and problem-solving, conceptual ability and mental flexibility (sorting test), as well as multiple measures of immediate and delayed recall. In addition, we examined measures of visuospatial function and language and we did not expect to find differences on these measures. We also hypothesised that impairments would improve with successful treatment of anxiety and that there would be a significant relationship between disability and cognitive performance, particularly in the executive domain.

Method

Study design

This was a National Institute of Mental Health sponsored 12-week double-blind randomised controlled trial of escitalopram *v.* placebo in older adults with a principal diagnosis of GAD, conducted in primary care practices and a specialty academic mental health centre in Pittsburgh, Pennsylvania, from 2005 to 2008 (trial registration NCT00105586).

Participants

This clinical trial recruited 177 individuals aged 60 or older meeting DSM-IV criteria for GAD.¹⁰ We excluded a total of 17 participants, due to central nervous system disease ($n = 13$), physical impairment that precluded neuropsychological assessment ($n = 3$) or refusal to

undergo neuropsychological assessment ($n=1$). Thus in the present analysis there were 160 participants in the GAD group whose mean age at onset was 39.6 years (s.d.=26.89) with a mean duration of 32.01 years (s.d.=26.6). Participants were assessed at baseline (pre-treatment) and after the 12-week trial. For comparison, we also assessed 37 older adults without dementia and without a psychiatric history equated with the GAD participants on gender and ethnicity as well as years of education and medical (including vascular disease) burden with the same neuropsychological battery, at a single time point.

Details on participant recruitment, retention and evaluation have been described elsewhere.^{11,12} During screening, all potential participants were evaluated by a board-certified geriatric psychiatrist (E.J.L.). Psychiatric diagnosis was established with the Structured Clinical Interview for Axis I DSM-IV Disorders (SCID-IV)¹³ administered by formally trained master's and doctoral degree level clinicians and a consensus diagnostic conference attended by the raters and at least two geriatric psychiatrists. Potential participants with an established diagnosis of dementia were excluded from the study; anyone with suspected dementia based on the screening evaluation was excluded. Moreover, the Mini-Mental State Examination (MMSE)¹⁴ was administered in this study and all data on anyone with a score <26 were closely reviewed by a geriatric psychiatrist (E.J.L.).

Measures

Participants underwent a broad-based pre-treatment assessment that included clinical, psychosocial and biological measures (described previously¹¹) as well as neuropsychological assessment (supervised by M.A.B.). We used a brief but comprehensive battery of neuropsychological measures that is widely used with older adults. The neuropsychological measures included the Repeatable Battery for the Assessment of Neuropsychological Status (RBANS).¹⁵ Forms A and B were administered in a counter-balanced manner to minimise practice effects. The RBANS provides a total index score and subscale index scores measuring language (confrontation naming and category fluency), visuo-spatial construction skills (figure copy and line orientation tasks), attention (forward digit span and coding tasks), immediate memory (list and story recall) and delayed memory (list, story and figure recall). Because the RBANS does not assess executive functioning and other higher-level abilities such as working memory, we also administered the Delis-Kaplan Executive Function System (D-KEFS)¹⁶ sorting test and used the confirmed correct sorts that measure problem-solving, conceptual ability and mental flexibility. We also used the colour-word interference score of the Stroop Neuropsychological Screening Test¹⁷ to measure the ability to inhibit automatic responses, and the Wechsler Adult Intelligence Scale (WAIS-III)¹⁸ letter-number sequencing subtest as a measure of working memory.

Among individuals with anxiety disorders both depression and anxiety symptoms are confounded making it extremely difficult to tease the two apart. Moreover, many of the most widely used mood scales (e.g. the Hamilton Rating Scale for Depression (HRSD)¹⁹) contain items that measure symptoms common to both depression and anxiety. In order to help tease apart the effects of depression *v.* anxiety, we examined depressive symptoms with the least confounded measure, the HRSD core depression items ($q1 + q2 + q3 + q7$).²⁰

We assessed disability with the Function and Disability Instrument (FDI) limitations and frequency subscales.²¹ These subscales measure a person's self-reported performance of socially defined life tasks that are expected of an individual within a typical environment, along two domains: how much difficulty

he or she has performing activities (activity limitations subscale) and how often he or she performs activities (frequency subscale). The limitation and frequency subscales of the FDI correspond to the disability domains of activity limitation and participation restriction, consistent with the International Classification of Function, Disability and Health.²² Both of these domains of function are impaired in late-life GAD.⁹

Statistical analysis

Prior to statistical analysis, we examined the data for normality and used transformations where necessary. If the distribution could not be normalised a non-parametric test was used. We generated descriptive statistics to characterise the GAD and comparison groups on key demographic and clinical characteristics.

Neuropsychological functioning of GAD and comparison groups at baseline

To characterise neuropsychological function among older adults with a principal diagnosis of GAD, we compared neuropsychological functioning of the GAD and comparison groups at baseline using analysis of variance. Seven main neuropsychological measures were chosen *a priori* to represent the domains of attention (digit span forward), information processing speed (coding), and the specific executive functions of inhibition (Stroop), working memory (letter-number sequencing), and problem-solving, conceptual ability and mental flexibility (sorting test), as well as immediate memory (RBANS immediate memory index) and delayed memory (RBANS delayed memory index). Age and education were used as covariates in these models except for those involving the RBANS memory scores, because these scores are age-adjusted based on test norms. We also examined baseline differences in index scores of the remaining two RBANS domains, visuospatial construction and language. In these models only education was used as a covariate since the derived scaled scores are adjusted for age. We report the results both with and without correction for multiple comparisons. Generalised anxiety disorder is frequently accompanied by depressive symptoms so to examine the effect of depression on neuropsychological functioning without obscuring the effect of GAD, we re-ran the analyses, once excluding the participants who also met criteria for major depressive disorder (MDD, $n=15$) and once covarying for MDD diagnosis.

Functional correlates of neuropsychological performance at baseline

Because group and depression scores are highly multicollinear and could not be entered into the same model, to further examine the potential influence of depression, we examined the correlations of HRSD total score and the core HRSD depression items score with the neuropsychological data. The correlations were not significant and therefore we did not control for either in the analyses comparing cognitive performance between the groups.

To examine the relationship between cognitive function and everyday functioning in late-life GAD, we calculated Spearman correlation coefficients between the nine neuropsychological variables and the FDI frequency and limitations subscales.

Comparing neuropsychological functioning pre- and post-treatment in the GAD group

As cognitive function in cognitively normal individuals may not improve with efficacious treatment for anxiety, and to identify

any cognitive changes related to treatment of anxiety, we compared neuropsychological functioning pre- and post-treatment in the low-scoring GAD treatment completers group using a repeated measures mixed-effect model with random intercept and slope. We examined treatment (drug *v.* placebo), time, and treatment \times time interactions. A participant was considered to be a low cognitive scorer at baseline if he or she performed in the lower half of a median split of the GAD group's RBANS total index scores.

We also examined neuropsychological change scores among the GAD group in participants whose anxiety symptoms significantly improved, to see which, if any, neuropsychological measures also improved, using the Wilcoxon Signed Rank Exact test. A participant was considered 'improved' if he or she reported having 'much improved' or 'very much improved' on the Clinical Global Impressions (CGI)²³ scale (CGI score ≤ 2) post-treatment.

Results

Descriptive analyses

Individuals in the GAD group were younger than those in the comparison group (71.6 (s.d.=7.7) *v.* 74.9 (s.d.=6.2) years, $P=0.012$; Table 1). Otherwise, the GAD and comparison groups did not differ in terms of relevant demographic and clinical characteristics, including medical and vascular disease burden, as measured with the Cumulative Illness Rating Scale – Geriatrics.²⁵

Participants who dropped out before week 12 ($n=31$) were not significantly different from completers ($n=129$) in age, gender, ethnicity, years of education, age at onset of GAD or rates of comorbid MDD. However, participants who dropped out before week 12 were significantly different from completers in terms of baseline severity on the Hamilton Rating Scale for Anxiety²⁶ (HRSA; mean 25.1 (95% CI 22.9–27.2) *v.* 22.6 (95% CI 21.9–23.2, $P=0.003$) and HRSD (mean 14.0 (95% CI 12.3–15.7) *v.* 11.6 (95% CI 11.0–12.2), $P=0.02$), co-prescription of benzodiazepine (drop-out rate for benzodiazepine users 33.3%

($n=9/27$) and for non-users 16.0% ($n=24/150$), $P=0.04$) and ethnicity (drop-out rate for participants who were White 15.9% ($n=23/145$) and for those who were Black 31.2% ($n=10/32$), $P=0.04$).

Neuropsychological functioning of GAD and comparison groups at baseline

Among the main outcome measures (Table 2 and Fig. 1), after controlling for age and education, those in the GAD group performed significantly worse than those in the comparison group on RBANS coding ($F(1,192)=6.44$, $P=0.012$), the Stroop test ($F(1,189)=3.92$, $P=0.049$), letter–number sequencing ($F(1,192)=8.37$, $P=0.004$), the D–KEFS sorting test ($F(1,192)=10.28$, $P=0.002$) and both RBANS immediate memory ($F(1,193)=13.38$, $P=0.0003$) and delayed memory ($F(1,193)=9.03$, $P=0.003$) indexes. There was no difference between the groups on the RBANS digit span ($F(1,192)=0.96$, $P=0.33$) or on the other neuropsychological domains assessed, language ($F(1,193)=0.10$, $P=0.75$) and visuospatial construction ($F(1,193)=2.67$, $P=0.10$). When the comparisons were corrected using the stepwise Bonferroni method, all of the findings remained except that the two groups performed similarly on the Stroop and RBANS coding (Table 2). Size and strength of the results did not change when excluding participants who met criteria for MDD or using MDD as a covariate. We also re-ran the analyses using lorazepam co-prescription as a covariate and the results did not significantly change from the previous results (results not reported).

Functional correlates of neuropsychological performance at baseline

Table 3 displays correlations between the two FDI subscales and nine neuropsychological variables. Of the 18 correlations, 9 were statistically significant at $P<0.05$ suggesting that in late-life GAD, neuropsychological impairments are correlated with functional disability and lending credence to the importance of

Table 1 Participants: descriptive information

	Generalised anxiety disorder group ($n=160$)	Comparison group ($n=37$)	Test statistic		
			t (d.f.)	χ^2 (d.f.)	P
Age, years: mean (s.d.) ^a	71.6 (7.7)	74.9 (6.2)	-2.53 (195)		0.012
Female, % (n)	68.8 (110)	67.6 (25)		0.02 (1)	0.89
White, % (n)	82.5 (132)	91.9 (34)		2.00 (1)	0.16
Education, years: mean (s.d.) ^a	13.9 (2.8)	14.7 (3.0)	-1.48 (195)		0.14
Cumulative Illness Rating Scale – Geriatrics, mean (s.d.)					
Total	8.5 (3.7)	7.9 (3.1)	0.85 (195)		0.40
Heart and vascular scales	2.3 (1.7)	2.1 (1.3)	0.70 (195)		0.48
Hamilton Rating Scale for Anxiety, ^a mean (s.d.), n	22.8 (4.5)	4.8 (3.7) 36	19.89 (40.4)		<0.0001 ^b
Penn State Worry Questionnaire, ²⁴ mean (s.d.), n	56.3 (12.6) 158	28.1 (6.8) 30	17.71 (73.3)		<0.0001 ^b
Generalised anxiety disorder, mean (s.d.), n					
Onset, years	39.6 (26.9) 159				
Duration, years	32.01 (26.00)				
Comorbid anxiety disorder, % (n)	34.0 (53/156)				
Comorbid major depressive disorder, % (n)	15.4 (24/156)				
Hamilton Rating Scale for Depression, mean (s.d.)					
Total	11.93 (3.89)	1.78 (2.20)			<0.0001
Core depression items	3.01 (1.85)	0.11 (0.31)			<0.0001
Prescribed lorazepam, mean (s.d.)	22.5	2.7		772 (1)	0.006
Mini-Mental State Exam, mean (s.d.), n	28.14 (1.72) 159	29.03 (1.16)			0.0015

a. Square root (X) transformation used in the analyses. Means and standard deviations reported in the original units.
b. Satterthwaite test used because of unequal variances.

Table 2 Neuropsychological measures

	GAD group (n = 160)	Comparison group (n = 37)	Test statistics															
			Age				Education				Group				Partial Eta ²			
			F (d.f.)	P	F (d.f.)	P	F (d.f.)	P	F (d.f.)	P	Age	Education	Group	Stepwise Bonferroni corrected, group: P				
RBANS, mean (s.d.), n	10.4 (2.7) 159	10.9 (2.3)	1.05 (1,192)	0.31	6.21 (1,192)	0.014	0.96 (1,192)	0.33	0.005	0.031	0.005	0.005	0.66					
Digit span	39.0 (10.7) 159	42.9 (9.2)	17.22 (1,192)	<0.0001	10.15 (1,192)	0.002	6.44 (1,192)	0.012	0.082	0.050	0.032	0.06						
Coding	3.6 (3.1) 156	3.0 (1.7)	14.48 (1,189)	0.0002	1.57 (1,189)	0.21	3.92 (1,189)	0.049	0.071	0.008	0.020	0.20						
Stroop ^a , mean (s.d.), n	8.7 (2.7) 159	9.9 (2.1)	15.04 (1,192)	0.0001	15.58 (1,192)	0.0001	8.37 (1,192)	0.004	0.073	0.075	0.042	0.026						
Letter–number sequencing, mean (s.d.), n	7.2 (3.4) 159	9.0 (2.8)	8.67 (1,192)	0.004	45.13 (1,192)	<0.0001	10.28 (1,192)	0.002	0.043	0.190	0.051	0.013						
D–KEFS, confirmed correct sorts: mean (s.d.), n	95.2 (17.4) 159	107.3 (13.5)	13.10 (1,193)	0.0004	13.38 (1,193)	0.0004	13.38 (1,193)	0.0003	0.064	0.064	0.065	0.003						
RBANS, mean (s.d.), n	92.5 (17.4) 159	102.2 (10.8)	7.16 (1,193)	0.008	7.16 (1,193)	0.008	9.03 (1,193)	0.003	0.036	0.036	0.045	0.021						
Immediate memory index score	89.4 (18.5) 159	95.7 (14.7)	11.07 (1,193)	0.001	11.07 (1,193)	0.001	2.67 (1,193)	0.10	0.05	0.05	0.014	0.31						
Delayed memory index score	98.7 (14.7) 159	100.6 (10.5)	20.71 (1,193)	<0.0001	20.71 (1,193)	<0.0001	0.10 (1,193)	0.75	0.10	0.10	0.001	0.75						
Visuospatial construction index score																		
Language index score																		

GAD, generalised anxiety disorder; RBANS, Repeatable Battery for the Assessment of Neuropsychological Status; D–KEFS, Delis–Kaplan Executive Function System.
a. Natural Log (X) transformation used in the analyses. Means and standard deviations reported in the original units.

neuropsychological functioning in late-life GAD. Significant correlations included: the FDI frequency subscale with RBANS coding, letter–number sequencing, D–KEFS sorting, and RBANS immediate and delayed memory index scores as well as the language index score; and the FDI limitations subscale with RBANS coding and letter–number sequencing and the RBANS language index score.

Comparing neuropsychological functioning pre- and post-treatment in GAD participants

Low cognitive scorers

Sixty-six participants scored below the group median score (total index score ≤ 94) on RBANS (30 in the active and 36 in the placebo treatment conditions) and thus were considered low cognitive scorers for the purpose of these analyses. The scores of the 66 participants who performed below the median RBANS total score ranged from the low end of the average range (RBANS digit span and the RBANS language index score) to mildly impaired (approximately 1 standard deviation below the age-corrected mean: letter–number sequencing, RBANS immediate and delayed memory index scores) to moderately impaired (1 to 2 standard deviations below the age-corrected mean: D–KEFS sorting, RBANS coding task, RBANS visuospatial construction index), to moderately to severely impaired (2 to 3 standard deviations below the age-correlated mean: the Stroop test). There was no difference between the lower and upper halves in age at onset of GAD (39 years for both groups; $P=0.99$) or per cent who had comorbid MDD (20 v. 9.7, $P=0.10$), but the lower half did have slightly higher anxiety as measured by the HRSA (23.42 v. 21.37, $P=0.05$).

The repeated measures ANOVA revealed no significant main effects for treatment group, time, or treatment \times time interactions for the Stroop test, RBANS digit span, coding or immediate memory or language index scores. There were significant main effects for time but not for treatment group or treatment \times time interactions on letter–number sequencing ($F(1,64)=4.82$, $P=0.032$) and RBANS delayed memory ($F(1,64)=6.65$, $P=0.01$) and visuospatial construction index scores, suggesting that working memory, delayed memory and visuospatial ability improved in both treatment conditions over the course of the study. There was a significant treatment \times time interaction on the D–KEFS sorting task ($F(1,62)=4.06$, $P=0.048$) indicating that the group receiving escitalopram improved in problem-solving, conceptual ability and mental flexibility more than did the group receiving placebo.

Reported improvement in anxiety

Forty-two participants reported ‘improvement’ in anxiety (CGI score ≤ 2) over the course of the study. These participants experienced significant improvement on the Stroop (change score 0.60 (s.d.=5.38), $P=0.0063$) and both RBANS immediate (change score 3.74 (s.d.=10.08), $P=0.0114$) and delayed memory (change score 5.41 (s.d.=10.08), $P=0.0019$), indicating an association between improved anxiety and improved ability to engage in inhibition and episodic recall in both treatment conditions over the course of the study.

Discussion

Main findings

To our knowledge, this is the first large-scale study to comprehensively evaluate neuropsychological function in late-life GAD and its response to treatment in a systematic manner. We found broad based, but not global impairments. Older participants with




Fig. 1 Boxplots depicting performance of generalised anxiety disorder (GAD) and comparison groups on the neuropsychological variables.

(a) Repeatable Battery for the Assessment of Neuropsychological Status (RBANS) digital span; (b) RBANS coding; (c) Stroop; (d) letter-number sequencing; (e) Delis-Kaplan Executive Function System (D-KEFS) confirmed correct sorts; (f) RBANS immediate memory index score; (g) RBANS delayed memory index score.

GAD performed worse than those in the comparison group on measures of information processing speed, working memory, inhibition and problem-solving (including concept formation and mental flexibility), as well as immediate and delayed memory. These findings are consistent with published literature describing decrements in memory and executive functions in young and middle-aged adults²⁷ and memory and working memory in older adults with GAD.^{28,29} The impairments of highest magnitude were

in memory, as immediate and delayed memory in the GAD group were about one standard deviation below that of the comparison group. This finding fits well with the literature review of Beaudreau & O'Hara,³⁰ who noted that memory impairments tended to be the most consistent cognitive deficits in late-life anxiety. Our findings also suggest that worry may compete for cognitive resources thereby interfering with execution of some especially vulnerable cognitive functions, including

Table 3 Functional correlates of neuropsychological performance at baseline ($n = 155$)

	Rho	
	FDI, frequency	FDI, limitation
RBANS		
Digit span	0.08	0.04
Coding	0.29**	0.23**
Stroop ^a	-0.11	-0.09
Letter-number sequencing	0.29**	0.17*
D-KEFS: confirmed correct sorts	0.20*	0.07
RBANS		
Immediate memory index score	0.26**	0.14
Delayed memory index score	0.17*	0.07
Visuospatial construction index score	0.08	0.03
Language index score	0.26**	0.17*

FDI, Function and Disability Instrument; RBANS, Repeatable Battery for the Assessment of Neuropsychological Status; D-KEFS, Delis-Kaplan Executive Function System.
a. Natural Log (X) transformation used in the analyses.
* $P < 0.05$, ** $P < 0.01$.

processing speed, some aspects of executive function and episodic memory.

We also found that neuropsychological functioning was significantly, but relatively weakly correlated with everyday functioning, consistent with the hypothesis that impaired cognitive function plays a significant role in the functional disability associated with late-life GAD.³¹

Finally, we found that among participants with GAD with poor cognitive performance at baseline, multiple domains showed improvement during the course of the clinical trial, suggesting that either clinical improvement in anxiety led to improvements in poor cognitive performance or clinical improvement in poor cognitive performance led to improvements in anxiety. Further, those who received escitalopram had greater improvements compared with placebo on the D-KEFS sorting task, which measures problem-solving, concept formation and mental flexibility. In addition, those participants with GAD who reported improved anxiety over the course of the trial experienced significant improvement in inhibition and both immediate and delayed recall.

Findings from other studies

Most studies of cognitive function and anxiety in older adults examine community-based populations with symptomatic measures of anxiety and report that higher levels of anxiety symptoms are associated with poorer fluid intelligence,³² complex visuospatial skills,³³ learning and memory,³³⁻³⁶ information processing speed^{30,34,37} and executive functions^{34,36} including inhibition.³⁰ By contrast, some community-based studies report no association between anxiety symptoms and cognitive function in late life.^{38,39} Three clinic-based studies have examined cognitive functioning in older adults who met specific clinical criteria for GAD. We previously reported that relative to a comparison group, individuals with late-life GAD exhibited impairment on measures of immediate and delayed recall as well as mental set shifting.²⁸ Caudle and colleagues²⁹ found that greater GAD symptom severity was associated with poorer working memory and Price & Mohlman⁴⁰ found an association with better inhibitory control (as measured with the Stroop).

The causal relationship between generalised anxiety and cognitive impairment is unclear. One prevalent model posits that

worry is a cognitive process designed to avoid the anxiogenic images that induce somatic activation through increased noradrenergic discharge.⁴¹ However, it is not clear whether the affective (increased anxiety) or cognitive (avoidance) component of GAD plays a more prominent role in the interplay between GAD and cognitive impairment. Affective interference (e.g. selective processing of anxious information at the expense of other cognitive tasks⁴²) seems to be related to a particular aspect of anxiety, namely ruminative worry. Rumination might mitigate the ability to shift resources between emotional and cognitive tasks, thus reducing task performance.⁴³ In a recent report, our group has shown that older adults with depression and high comorbid anxiety have elevated activation of several brain areas involved in cognitive performance, including dorsal prefrontal cortex and dorsal anterior cingulate cortex.⁴⁴

Generalised anxiety disorder may be particularly detrimental to cognitive function in older adults,³⁶ who have less cognitive reserve against central nervous system insults than do younger adults. Cognitive reserve refers to the degree to which an individual is able to maintain cognitive function in the face of mounting neuropathology. Moreover, worry may compete for the cognitive resources necessary for working memory thereby interfering with execution of some cognitive functions.³⁰ Additionally, ageing increases vulnerability to cognitive impairment because homeostatic mechanisms that prevent an excessive biological stress response are diminished.^{31,37} Consequently, some deleterious effects of excessive stress response – such as neurotoxic hypercortisolemia – worsen with age. Despite these threats to cognition, few studies have examined neuropsychological functioning in late-life anxiety disorders, and there are very few reports on late-life GAD in particular.

Whereas observational studies provide some support for a cross-sectional association between anxiety and cognitive impairment in older adults, better-designed cross-sectional and longitudinal studies may be the more appropriate approach. Experimental research designs that involve manipulating anxiety levels, rather than simply assessing them repeatedly, could also provide informative data about the possible causal impact of anxiety on cognitive change.

Limitations

Our study has some limitations. Although the RBANS has parallel forms to reduce practice effects, improved performance over time may be attributable to repetition rather than true treatment effects. Also, the clinical trial was only 12 weeks long; longer protocolised evaluations may be more robust for finding differences in the course of cognitive decline and its amelioration with effective treatment. Comparing individuals with late-life GAD with non-psychiatric controls leaves open the possibility that the impairments that were identified could be the result of having a psychiatric disorder in general and may not be specific to late-life GAD. Although future studies should employ this approach, in the case of GAD, the most appropriate psychiatric control group is unclear. It is possible that IQ differences could explain our findings. However, all of our participants were recruited from the same sources and the average years of education was similar in both the GAD and comparison groups, minimising the likelihood that there were substantial IQ differences. We did not measure sleep quality or quantity, and it is possible that sleep disturbance may account for some or all of the effects on memory. Finally, our understanding of the neurobiology of late-life GAD is lacking, and better understanding of the structural or functional changes leading to cognitive impairment in this disorder may yield more informative neuropsychological hypotheses and, eventually, better

treatments. The greatest strength of this study is that it examines the relationship between anxiety and cognition in the context of an experimental manipulation (i.e. treatment). In the context of a 12-week clinical trial, it is more likely that reductions in anxiety were driving improvements in cognition rather than the reverse.

Implications

In conclusion, GAD in older adults is associated with widespread neuropsychological impairments, including information processing speed, working memory, inhibition, problem-solving and both immediate and delayed memory. Although there was specific improvement in problem-solving, concept formation and mental flexibility attributable to SSRI treatment, many of these cognitive domains showed improvement during the course of treatment regardless of treatment assignment. Although medication-specific benefits are minimal in this analysis, we found that improvement in anxiety symptoms was associated with neuropsychological improvements. Such findings are not a treatment effect *per se* but could be interpreted as showing that the clinical fluctuation of anxiety symptoms also affects cognitive performance. Future treatment research might usefully focus on providing long-term remission stability, both clinically and cognitively and should examine whether such improvements translate to meaningful gains in function and quality of life. These findings underscore the importance of cognitive functioning as a potential treatment target to reduce the impairment associated with anxiety disorders in older adults.

Meryl A. Butters, PhD, Department of Psychiatry, University of Pittsburgh, Pittsburgh, Pennsylvania, USA; **Rishi K. Bhalla**, PhD, Department of Psychiatry, University of British Columbia, Vancouver, Canada; **Carmen Andreescu**, MD, Department of Psychiatry, University of Pittsburgh, Pittsburgh, Pennsylvania, USA; **Julie Loebach Wetherell**, PhD, Department of Psychiatry, University of California at San Diego and Psychology Service, VA San Diego Healthcare System, San Diego, California, USA; **Rose Mantella**, PhD, Abbott Laboratories, Illinois, USA; **Amy E. Begley**, MA, Department of Psychiatry, University of Pittsburgh, Pittsburgh, Pennsylvania, USA; **Eric J. Lenze**, MD, Department of Psychiatry, Washington University in St. Louis, Missouri, USA

Correspondence: Meryl A. Butters, PhD, Western Psychiatric Institute and Clinic, 3811 O'Hara St, Pittsburgh, PA 15213, USA. Email: buttersma@upmc.edu

First received 30 Nov 2010, final revision 15 Mar 2011, accepted 6 Apr 2011

Funding

This research was supported by United States Public Health Service grants MH070547, MH072947 and MH52247, as well as a 2009 NARSAD Young Investigator Award to C.A.

Acknowledgements

We would like to thank Michelle Zmuda, Program Coordinator of the Geriatric Neuropsychology Research Program for overseeing the neuropsychological assessments.

References

- 1 Beekman AT, Bremner MA, Deeg DJ, van Balkom AJ, Smit JH, de Beurs E, et al. Anxiety disorders in later life: a report from the Longitudinal Aging Study Amsterdam. *Int J Geriatr Psychiatry* 1998; **13**: 717–26.
- 2 Bryant C, Jackson H, Ames D. The prevalence of anxiety in older adults: methodological issues and a review of the literature. *J Affect Disord* 2008; **109**: 233–50.
- 3 Gum AM, King-Kallimanis B, Kohn R. Prevalence of mood, anxiety, and substance-abuse disorders for older Americans in the national comorbidity survey-replication. *Am J Geriatr Psychiatry* 2009; **17**: 769–81.
- 4 Diefenbach GJ, Stanley MA, Beck JG. Worry content reported by older adults with and without generalized anxiety disorder. *Aging Ment Health* 2001; **5**: 269–74.
- 5 Chou KL. Age at onset of generalized anxiety disorder in older adults. *Am J Geriatr Psychiatry* 2009; **17**: 455–64.
- 6 Le Roux H, Gatz M, Wetherell JL. Age at onset of generalized anxiety disorder in older adults. *Am J Geriatr Psychiatry* 2005; **13**: 23–30.
- 7 Lenze EJ, Rollman BL, Shear MK, Dew MA, Pollock BG, Ciliberti C, et al. Escitalopram for older adults with generalized anxiety disorder: a randomized controlled trial. *JAMA* 2009; **301**: 295–303.
- 8 Stanley MA, Beck JG, Novy DM, Averill PM, Swann AC, Diefenbach GJ, et al. Cognitive-behavioral treatment of late-life generalized anxiety disorder. *J Consult Clin Psychol* 2003; **71**: 309–19.
- 9 Porensky EK, Dew MA, Karp JF, Skidmore E, Rollman BL, Shear MK, et al. The burden of late-life generalized anxiety disorder: effects on disability, health-related quality of life, and healthcare utilization. *Am J Geriatr Psychiatry* 2009; **17**: 473–82.
- 10 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th edn) (DSM-IV)*. APA, 1994.
- 11 Lenze EJ, Wetherell JL. Bringing the bedside to the bench, and then to the community: a prospectus for intervention research in late-life anxiety disorders. *Int J Geriatr Psychiatry* 2009; **24**: 1–14.
- 12 Mantella RC, Butters MA, Amico JA, Mazumdar S, Rollman BL, Begley AE, et al. Salivary cortisol is associated with diagnosis and severity of late-life generalized anxiety disorder. *Psychoneuroendocrinology* 2008; **33**: 773–81.
- 13 First MB, Spitzer RL, Gibbon M, Williams JBW. *Structured Clinical Interview for DSM-IV Axis I Disorders – Nonpatient Edition*. Biometrics Research Department, New York State Psychiatric Institute, 1996.
- 14 Folstein M, Folstein S, McHugh P. 'Mini-Mental State', a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; **12**: 189–98.
- 15 Randolph C. *The Repeatable Battery for the Assessment of Neuropsychological Status*. Psychological Corporation, 1998.
- 16 Delis D, Kaplan E, Kramer J. *Delis-Kaplan Executive Function Scale*. Psychological Corporation, 2001.
- 17 Trennery M, Crosson B, DeBoe J, Leber W. *Stroop Neuropsychological Screening Test Manual*. Psychological Assessment Resources, 1989.
- 18 Wechsler D. *WAIS-III Administration and Scoring Manual*. Psychological Corporation, 1997.
- 19 Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960; **23**: 56–62.
- 20 Dombrovski AY, Blakesley-Ball RE, Mulsant BH, Mazumdar S, Houck PR, Szanto K, et al. Speed of improvement in sleep disturbance and anxiety compared with core mood symptoms during acute treatment of depression in old age. *Am J Geriatr Psychiatry* 2006; **14**: 550–4.
- 21 Jette AM, Haley SM, Coster WJ, Kooyomjian JT, Levenson S, Heeren T, et al. Late life function and disability instrument: I. Development and evaluation of the disability component. *J Gerontol A Biol Sci Med Sci* 2002; **57**: M209–16.
- 22 World Health Organization. *The World Health Report: Reducing Risks, Promoting Healthy Life*. WHO, 2002.
- 23 Guy W. *ECDEU Assessment Manual for Psychopharmacology*. US Department of Health, Education, and Welfare Public Health Service Alcohol, Drug Abuse, and Mental Health Administration, 1976.
- 24 Meyer TJ, Miller ML, Metzger RL, Borkovec TD. Development and validation of the Penn State Worry Questionnaire. *Behav Res Ther* 1990; **28**: 487–95.
- 25 Miller MD, Paradis CF, Houck PR, Mazumdar S, Stack JA, Rifai AH, et al. Rating chronic medical illness burden in geropsychiatric practice and research: application of the Cumulative Illness Rating Scale. *Psychiatry Res* 1992; **41**: 237–48.
- 26 Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol* 1959; **32**: 10.
- 27 Airaksinen E, Larsson M, Forsell Y. Neuropsychological functions in anxiety disorders in population-based samples: evidence of episodic memory dysfunction. *J Psychiatr Res* 2005; **39**: 207–14.
- 28 Mantella RC, Butters MA, Dew MA, Mulsant BH, Begley AE, Tracey B, et al. Cognitive impairment in late-life generalized anxiety disorder. *Am J Geriatr Psychiatry* 2007; **15**: 673–9.
- 29 Caudle DD, Senior AC, Wetherell JL, Rhoades HM, Beck JG, Kunik ME, et al. Cognitive errors, symptom severity, and response to cognitive behavior therapy in older adults with generalized anxiety disorder. *Am J Geriatr Psychiatry* 2007; **15**: 680–9.
- 30 Beaudreau SA, O'Hara R. The association of anxiety and depressive symptoms with cognitive performance in community-dwelling older adults. *Psychol Aging* 2009; **24**: 507–12.
- 31 Wetherell JL, Le Roux H, Gatz M. DSM-IV criteria for generalized anxiety disorder in older adults: distinguishing the worried from the well. *Psychol Aging* 2003; **18**: 622–7.

- 32 Cohen D, Eisdorfer C, Vitaliano PP, Bloom V. The relationship of age, anxiety, and serum immunoglobulins with crystallized and fluid intelligence. *Biol Psychiatry* 1980; **15**: 699–709.
- 33 Wetherell JL, Reynolds CA, Gatz M, Pedersen NL. Anxiety, cognitive performance, and cognitive decline in normal aging. *J Gerontol B Psychol Sci Soc Sci* 2002; **57**: P246–55.
- 34 Bierman EJ, Comijs HC, Jonker C, Beekman AT. Effects of anxiety versus depression on cognition in later life. *Am J Geriatr Psychiatry* 2005; **13**: 686–93.
- 35 Deptula D, Singh R, Pomara N. Aging, emotional states, and memory. *Am J Psychiatry* 1993; **150**: 429–34.
- 36 Paterniti S, Dufouil C, Bisserbe JC, Alperovitch A. Anxiety, depression, psychotropic drug use and cognitive impairment. *Psychol Med* 1999; **29**: 421–8.
- 37 Hogan MJ. Divided attention in older but not younger adults is impaired by anxiety. *Exp Aging Res* 2003; **29**: 111–36.
- 38 Biringer E, Mykletun A, Dahl AA, Smith AD, Engedal K, Nygaard HA, et al. The association between depression, anxiety, and cognitive function in the elderly general population—the Hordaland Health Study. *Int J Geriatr Psychiatry* 2005; **20**: 989–97.
- 39 Lindsay J, Briggs K, Murphy E. The Guy's/Age Concern survey. Prevalence rates of cognitive impairment, depression and anxiety in an urban elderly community. *Br J Psychiatry* 1989; **155**: 317–29.
- 40 Price RB, Mohlman J. Inhibitory control and symptom severity in late life generalized anxiety disorder. *Behav Res Ther* 2007; **45**: 2628–39.
- 41 Borkovec TD, Inz J. The nature of worry in generalized anxiety disorder: a predominance of thought activity. *Behav Res Ther* 1990; **28**: 153–8.
- 42 Ingram RE, Kendall PC, Smith TW, Donnell C, Ronan K. Cognitive specificity in emotional distress. *J Pers Soc Psychol* 1987; **53**: 734–42.
- 43 Whalen PJ, Bush G, McNally RJ, Wilhelm S, McInerney SC, Jenike MA, et al. The emotional counting Stroop paradigm: a functional magnetic resonance imaging probe of the anterior cingulate affective division. *Biol Psychiatry* 1998; **44**: 1219–28.
- 44 Andreescu C, Butters M, Lenze EJ, Venkatraman VK, Nable M, Reynolds 3rd CF, et al. fMRI activation in late-life anxious depression: a potential biomarker. *Int J Geriatr Psychiatry* 2009; **24**: 820–8.

100
years ago

Thomas Drapes, Medical Superintendent of the Enniscorthy

Dermot Walsh

In these days of some perturbation concerning the overinclusiveness of proposed psychiatric classificatory systems such as DSM-5, it is salutary to reflect that this concern is nothing new. Dr Thomas Drapes, Medical Superintendent of the Enniscorthy, Co. Wexford Asylum, voiced his worry in 1911, when Eugen Bleuler was introducing the word 'schizophrenia' to the psychiatric dictionary. In an article 'The personal equation in alienism' in the *Journal of Mental Science* he wrote (May 1911): 'In classification we have another example of the effects of the personal equation. That there is need for some kind of classification is agreed on by all; what should be the basis of classification is agreed by none. And so we have scheme after scheme ushered into existence by one writer after another, framed according to the mental proclivities of each individual authority, all different, and each constructed, not on a single basis, but on sundry and diverse bases, pathological, aetiological, symptomatological, and so on, forming a veritable medley or hotch-potch, most embarrassing to the student of psychiatry. . . A new variety of insanity is announced. It is ushered into existence by some illustrious psychiatrist. It is accepted with acclaim by one class of mind, regarded as a new discovery by which diagnosis and prognosis will be in future rendered comparatively easy, the crooked shall be made straight, and the rough places plain. (607)'

His obituarist, fellow countryman, psychiatrist and Inspector of Lunacy Sir W. R. Dawson, on the first page of the *Journal* in April 1920, reflected that Drapes 'could never bring himself to adopt the doctrines of Kraepelin' and thought that 'Freudism is, in much of its theory, scientifically unsound'. Drapes was probably the most enquiring psychiatrist in Ireland of his generation and contributed frequently to the *Journal* on topics as diverse as 'On the alleged increase of insanity in Ireland' and 'Acute hallucinatory insanity of traumatic origin'. Literate in French and German, he reviewed many publications in these languages for the *Journal*. In 1912, he became co-editor of the *Journal* and in 1915 editor-in-chief. In 1911/1912, he was elected president of the Royal Medico-Psychological Association, but declined the honour on health grounds.

Earlier in 2011, the Enniscorthy Asylum, over which Drapes had presided from 1883 to 1919, closed. One wonders what he would have thought of the 'community psychiatry' which led to its demise.

The British Journal of Psychiatry (2011)
199, 218. doi: 10.1192/bjp.bp.111.098541