

# Clustering of adherence to personalised dietary recommendations and changes in healthy eating index within the Food4Me study

Katherine M Livingstone<sup>1</sup>, Carlos Celis-Morales<sup>1</sup>, Jose Lara<sup>1</sup>, Clara Woolhead<sup>2</sup>, Clare B O'Donovan<sup>2</sup>, Hannah Forster<sup>2</sup>, Cyril FM Marsaux<sup>3</sup>, Anna L Macready<sup>4</sup>, Rosalind Fallaize<sup>4</sup>, Santiago Navas-Carretero<sup>5,6</sup>, Rodrigo San-Cristobal<sup>6</sup>, Silvia Kolossa<sup>7</sup>, Lydia Tsigoti<sup>8</sup>, Christina P Lambrinou<sup>8</sup>, George Moschonis<sup>8</sup>, Agnieszka Surwitto<sup>9</sup>, Christian A Drevon<sup>10</sup>, Yannis Manios<sup>8</sup>, Iwona Traczyk<sup>9</sup>, Eileen R Gibney<sup>2</sup>, Lorraine Brennan<sup>2</sup>, Marianne C Walsh<sup>2</sup>, Julie A Lovegrove<sup>4</sup>, J Alfredo Martinez<sup>6</sup>, Wim HM Saris<sup>3</sup>, Hannelore Daniel<sup>7</sup>, Mike Gibney<sup>2</sup> and John C Mathers<sup>1,\*</sup> on behalf of the Food4Me Study

<sup>1</sup>Human Nutrition Research Centre, Institute of Cellular Medicine, Newcastle University, Biomedical Research Building, Campus for Ageing and Vitality, Newcastle upon Tyne NE4 5PL, UK; <sup>2</sup>UCD Institute of Food and Health, University College Dublin, Dublin, Republic of Ireland; <sup>3</sup>Department of Human Biology, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre, Maastricht, The Netherlands; <sup>4</sup>Hugh Sinclair Unit of Human Nutrition and Institute for Cardiovascular and Metabolic Research, University of Reading, Reading, UK; <sup>5</sup>Center for Nutrition Research, University of Navarra, Pamplona, Spain; <sup>6</sup>CIBER Fisiopatología Obesidad y Nutrición (CIBERObn), Instituto de Salud Carlos III, Madrid, Spain; <sup>7</sup>ZIEL Research Center of Nutrition and Food Sciences, Biochemistry Unit, Technical University of Munich, Munich, Germany; <sup>8</sup>Department of Nutrition and Dietetics, Harokopio University, Athens, Greece; <sup>9</sup>National Food & Nutrition Institute (IZZ), Warsaw, Poland; <sup>10</sup>Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo, Norway

Submitted 20 January 2016: Final revision received 18 May 2016: Accepted 20 June 2016: First published online 8 August 2016

## Abstract

**Objective:** To characterise clusters of individuals based on adherence to dietary recommendations and to determine whether changes in Healthy Eating Index (HEI) scores in response to a personalised nutrition (PN) intervention varied between clusters.

**Design:** Food4Me study participants were clustered according to whether their baseline dietary intakes met European dietary recommendations. Changes in HEI scores between baseline and month 6 were compared between clusters and stratified by whether individuals received generalised or PN advice.

**Setting:** Pan-European, Internet-based, 6-month randomised controlled trial.

**Subjects:** Adults aged 18–79 years ( $n$  1480).

**Results:** Individuals in cluster 1 (C1) met all recommended intakes except for red meat, those in cluster 2 (C2) met two recommendations, and those in cluster 3 (C3) and cluster 4 (C4) met one recommendation each. C1 had higher intakes of white fish, beans and lentils and low-fat dairy products and lower percentage energy intake from SFA ( $P < 0.05$ ). C2 consumed less chips and pizza and fried foods than C3 and C4 ( $P < 0.05$ ). C1 were lighter, had lower BMI and waist circumference than C3 and were more physically active than C4 ( $P < 0.05$ ). More individuals in C4 were smokers and wanted to lose weight than in C1 ( $P < 0.05$ ). Individuals who received PN advice in C4 reported greater improvements in HEI compared with C3 and C1 ( $P < 0.05$ ).

**Conclusions:** The cluster where the fewest recommendations were met (C4) reported greater improvements in HEI following a 6-month trial of PN whereas there was no difference between clusters for those randomised to the Control, non-personalised dietary intervention.

**Keywords**  
Clustering  
Personalised nutrition  
Dietary recommendations  
Healthy Eating Index

Global obesity prevalence has reached epidemic proportions, with 37% of men and 38% of women now either overweight or obese<sup>(1)</sup>. Poor dietary choices and

inadequate physical activity are the primary causes of obesity<sup>(2)</sup>. Current strategies for improving diet and other lifestyle behaviours, such as consuming five portions of

\*Corresponding author: Email john.mathers@newcastle.ac.uk

fruit and vegetables per day<sup>(3)</sup>, are based on 'one size fits all' generalised dietary guidelines. Given that the burden of obesity is increasing<sup>(1)</sup>, alternative strategies for improving dietary behaviours are being developed, including predictive, personalised, preventive and participatory interventions<sup>(4)</sup>. Recent evidence suggests that genetic-based personalised nutrition (PN) improves dietary intakes more than non-personalised advice<sup>(5)</sup>. However, since dietary intakes tend to cluster<sup>(6,7)</sup>, it may be possible to enhance the efficacy of interventions by further characterisation of participants according to their dietary and lifestyle behaviours and, subsequently, use this information to strengthen the basis for personalisation of the intervention. For example, lower intakes of fruit, vegetables and whole grains are often associated with higher intakes of red or processed meat<sup>(8)</sup>. In addition, less healthy dietary clusters are associated with increased disease risk<sup>(9)</sup> and unhealthy dietary and lifestyle behaviours are associated with higher levels of sedentary behaviour<sup>(7)</sup> and mortality<sup>(10,11)</sup>. Clustering individuals based on whether they meet dietary recommendations may be a useful predictive tool for estimating response to an intervention<sup>(12–14)</sup> and may help to stratify or personalise interventions.

The Food4Me Proof-of-Principle study was the first Internet-based study to demonstrate that PN advice was more effective in improving dietary intakes, including lowering intakes of red meat, compared with conventional 'one size fits all' population-based advice<sup>(15,16)</sup>. However, the characteristics of individuals clustered on the basis of adherence to current recommended dietary intakes of fruit and vegetables, whole grains, oily fish, dairy products and red and processed meat are unknown. Thus, the aims of the current analysis were to: (i) characterise European adults participating in the Food4Me study<sup>(17)</sup> according to clustering based on European recommendations for healthy eating; and (ii) determine whether cluster membership predicted dietary changes following a PN intervention.

## Methods

### *Study design and population*

The Food4Me study was a 6-month, four-arm, Internet-based, randomised controlled trial (RCT) in 1607 individuals conducted across seven European countries<sup>(17,18)</sup>. Participants were recruited via the Food4Me website<sup>(16)</sup> to emulate a web-based PN service. This was aided by local and national advertising of the study via the Internet, radio, newspapers, posters, e-flyers, social media and word of mouth. Recruitment took place between August 2012 and August 2013 in the following sites: University College Dublin (Ireland), Maastricht University (The Netherlands), University of Navarra (Spain), Harokopio University (Greece), University of Reading (UK), National Food and Nutrition Institute (Poland) and Technical University of Munich (Germany). The Research Ethics Committees at

each University or Research Centre delivering the intervention granted ethical approval for the study. The Food4Me trial was registered as an RCT (NCT01530139) at Clinicaltrials.gov. All participants expressing an interest in the study were asked to sign online consent forms at two stages in the screening process.

### *Intervention arms*

Participants were randomised to receive non-personalised, generalised dietary advice (Control) or one of three levels of PN (Level 1, Level 2 or Level 3). Briefly, non-personalised dietary advice was based on national dietary recommendations in each of the seven European countries. These 'standardised' recommendations included advice on energy intake and on the consumption of fruits and vegetables, whole grains, fish, dairy products, meat, type of fat and salt. Participants randomised to Level 1 received personalised dietary advice on how their intakes of these food groups at baseline compared with guideline amounts. Participants randomised to Level 2 received advice based on their dietary intake (as for Level 1) and also on their baseline phenotypic data. The phenotypic feedback was based on anthropometric measurements and nutrient- and metabolic-related biomarkers. Participants randomised to Level 3 received advice based on their dietary intake, phenotypic and genotypic data collected at baseline. The genotypic feedback was based on specific variants in five nutrient-responsive genes selected specifically for the study. Further details are provided elsewhere<sup>(17)</sup>.

### *Screening questionnaires and dietary intakes*

Participants eligible for inclusion in the RCT completed an online questionnaire to collect detailed information on sociodemographic, health and anthropometric characteristics and dietary habits. Following completion of this questionnaire, participants were asked to complete an online FFQ to estimate usual dietary intake. This FFQ, which was developed and validated for the study<sup>(19,20)</sup>, included 157 food items consumed frequently in each of the seven recruitment countries. Intakes of foods and nutrients were computed in real time using a food composition database based on *McCance and Widdowson's The Composition of Foods*<sup>(21)</sup>. Intakes of nutrients were assessed based on standardised recommendations (see online supplementary material, Supplemental Table 1) for dietary intakes of foods and food groups<sup>(17)</sup>, which were integrated and harmonised across eight European countries (UK, Ireland, Germany, The Netherlands, Spain, Greece, Poland and Norway)<sup>(22–25)</sup>. The following four food group recommendations were used in the present analysis: (i) eat at least five portions of fruit and vegetables every day (operationalised as  $\geq 400$  g); (ii) eat at least three portions of wholegrain products daily ( $\geq 50$  g); (iii) eat at least one portion of oily fish per week ( $\geq 150$  g); and (iv) eat less than three portions of red or processed meat

per week ( $\leq 450$  g)<sup>(17)</sup>. The Healthy Eating Index 2010 (HEI) was derived based on intakes of the following components: ratio of MUFA + PUFA to SFA, protein, salt, 'empty calories', refined grains, seafood and plant protein, fruit, whole fruit, vegetables, greens and beans, whole grains and dairy products<sup>(26)</sup>.

### **Personalised feedback report**

Participants randomised to PN received personalised reports via email at baseline, month 3 and month 6 of the intervention based on diet, anthropometric measurements and physical activity. Using information on the individual's intakes of nutrients, algorithms were used to rank information on need for dietary change and to provide participants with three specific dietary, food-based goals. For participants randomised to Level 2 and Level 3, the dietary advice was also based on phenotypic data (Level 2) and phenotypic plus genotypic data (Level 3). Reported intakes were compared with recommended intakes and determined to be adequate, high or low. If intakes were too high or too low, contributing foods were identified and specific messages developed to advise change in intake of those foods. Dietary intakes relative to recommendations were illustrated using a three-colour sliding scale: green representing 'Good, no change recommended', amber representing 'Improvement recommended' and red representing 'Improvement strongly recommended'. For the genotype-based information, risk was indicated using 'Yes' or 'No' according to whether the participant did or did not carry the higher-risk variant for each of the five nutrient-related genes included in the study. Additionally, each report contained a personalised message from the dietitian/nutritionist to the participant. Further details of the protocol are provided elsewhere<sup>(17)</sup>.

### **Anthropometric, sociodemographic and physical activity measures**

Detailed standardised online instructions were given for participants to self-measure and self-report their body weight, height and waist circumference via the Food4Me website<sup>(16)</sup>. BMI was estimated from body weight and height. Self-reported measurements were validated in a sub-sample of the participants ( $n$  140) and showed a high degree of reliability<sup>(27)</sup>. Physical activity level and time spent in sedentary behaviours were estimated from triaxial accelerometers (TracmorD, Philips Consumer Lifestyle, The Netherlands). Participants self-reported smoking habits and occupation. Occupations were grouped according to the European classifications of occupations and their salaries (the European-wide average salary for each occupation was compared with the mean overall salary; if the SD of the salary was  $>0.5$ , they were placed in Group 1, between 0.5 to  $-0.5$  were placed into Group 2, and  $<-0.5$  were placed into Group 3) as follows: Group 1, Professional and managerial (professionals; managers); Group 2, Intermediate (armed forces occupations;

technicians and associate professionals; clerical support workers); Group 3, Routine and manual (craft and related trades workers; plant and machine operators and assemblers; service and sales workers; elementary occupations; skilled agricultural, forestry and fishery workers)<sup>(28,29)</sup>. Categories for 'Students' and 'Retired and unemployed' were added.

### **Statistical analysis**

Data were analysed using the statistical software packages Stata version 13 and IBM SPSS Statistics Version 22.0. Clusters of dietary recommendations were generated based on whether participants met the following four food group recommendations at baseline and were coded as 0 or 1 accordingly: (i) eat at least five portions of fruit and vegetables every day (operationalised as  $\geq 400$  g); (ii) eat at least three portions of wholegrain products daily ( $\geq 50$  g); (iii) eat at least one portion of oily fish per week ( $\geq 150$  g); and (iv) eat less than three portions of red or processed meat per week ( $\leq 450$  g). Clusters were derived using the SPSS Two Step cluster analysis procedure<sup>(30)</sup>. Small pre-clusters were generated based on the log-likelihood distance criterion (Step 1) and were merged into distinct groups using agglomerative hierarchical clustering (Step 2). Automatic selection and the Bayesian information criterion were used to determine the optimal number of clusters. Robustness and stability of the final clusters were re-evaluated by random ordering of cases (four times). This clustering methodology identified the percentage of participants within each cluster who met recommended intakes of each of the four food groups of public health importance. Logistic regression was used to test for significant differences across categorical variables and ANOVA was used for continuous variables. Tukey pairwise comparisons were used to test for significant differences between clusters. Analyses were adjusted for age, sex, country, BMI, physical activity level and smoking, except when those (or related) variables were being assessed (i.e. analyses were not adjusted for BMI when assessing BMI, body weight or waist circumference). Results were deemed significant at  $P < 0.05$ . To exclude extreme intakes of the food groups used for clustering, the top and bottom 3 SD of these intakes were excluded prior to clustering.

### **Results**

Of the 5562 individuals who registered on the Food4Me website, 1607 were randomised into the study and a total of 1480 provided baseline data on dietary intakes<sup>(17)</sup>.

### **Dietary adequacies across Food4Me cohort**

Recommended intakes for nutrients are summarised in the online supplementary material, Supplemental Table 1. On average, 50% of individuals met the recommendations for

total fat (Supplemental Table 2). The percentage of individuals who met the recommendations for SFA, MUFA and PUFA intake was 54, 24 and 36%, respectively. Only 56% of individuals met the recommendation for carbohydrate intake, whereas 91% of individuals had adequate protein intakes. Only 7 and 46% of individuals met the recommendations for salt and dietary fibre intakes, respectively. Meeting recommended micronutrient intakes ranged from 61% (folate) to 99% (vitamin B<sub>12</sub>).

As summarised in the online supplementary material, Supplemental Table 2, approximately half (52%) of participants reported consuming at least five portions of fruit and vegetables per day and 32% consumed at least one portion of oily fish per week. Nearly three-quarters (74%) of participants consumed more than three servings of whole grains per day and approximately half of participants (51%) consumed less than three servings of red meat per week. Fourteen per cent of individuals met the recommendation for dairy product intake ( $\geq 600$  g/d).

### **Cluster characterisation**

Clustering of individuals according to whether they met the intake recommendations for dairy products, fruits and vegetables, oily fish, red meat and whole grains at baseline did not create clear clustering due to the low percentage of individuals who met the recommendation for dairy products (two clusters). Exclusion of dairy products as a clustering variable provided improved clustering, as estimated by the silhouette measure of cohesion and separation (average silhouette: 0.3 *v.* 0.5; four clusters; see online supplementary material, Supplemental Table 3). Cluster 1 (C1) was the largest (*n* 475) and was particularly characterised by individuals meeting the recommended intake for oily fish (100% of individuals); 74 and 69% of C1 members met the recommendations for whole grains and fruit and vegetables, respectively, whereas only 46% met the recommendation for red meat. Cluster 2 (C2; *n* 398) was the second largest and was particularly characterised by all members meeting recommendations for whole grains (100%) and red meat (100%), only 50% meeting the recommendation for fruit and vegetables and no one meeting the recommendation for oily fish. All individuals in cluster 3 (C3; *n* 348) met the recommendation for whole grains, but no one met the recommendation for oily fish or red meat, whereas only 48% met the recommended intake for fruit and vegetables. None of the participants in cluster 4 (C4; *n* 259) met the recommended intake for either oily fish or whole grains; only 50 and 29% of C4 members achieved the recommended intakes for red meat and fruit and vegetables, respectively (Supplemental Table 3).

### **Dietary intakes by cluster**

Intakes of oily fish and fruit and vegetables were higher in C1 than in C2, C3 and C4 ( $P < 0.05$ ), and wholegrain

intakes were higher in C1, C2 and C3 than in C4 (Table 1;  $P < 0.05$ ). Red meat intake was lower in C1, C2 and C3 than in C4 ( $P < 0.05$ ). Intakes of fruit juice, eggs, chicken, white fish, fish products, beans and lentils and low-fat dairy products were higher in C1 than C4, whereas intakes of non-wholegrain products were lower ( $P < 0.05$ ). Participants in C2 had lower intakes of chips and pizza and fried foods than C3 and C4 ( $P < 0.05$ ; Table 1). Total energy intake and ratio of energy intake to BMR were higher in C1 than in C2 and C4 and higher in C3 than in C2 ( $P < 0.05$ ; Table 1). Individuals in C1 derived higher percentages of energy intake from PUFA and protein than those in C2 and C4 ( $P < 0.05$ ) and individuals in C2 derived a higher percentage energy from carbohydrates than participants in C3 and C4 ( $P < 0.05$ ). In contrast, individuals in C1 had lower percentage energy intakes from total fat and SFA than those in C4 ( $P < 0.05$ ) and higher percentage energy intake from MUFA than participants in C2 and C3 ( $P < 0.05$ ). Individuals in C1 had lower percentage energy intake from sugar than those in C2 ( $P < 0.05$ ). Participants in C1 consumed more dietary fibre and salt than those in C2 and C4 ( $P < 0.05$ ).

More individuals in C1 met the recommendations for total fat intake (51%), SFA (62%), PUFA (42%) and dietary fibre (56%) than C4 cluster members (see online supplementary material, Supplemental Table 4). Fewer individuals in C1 met the recommendation for protein intake (86%) than those in C2 (97%) and C3 (93%). Furthermore, fewer individuals in C1 met the recommendation for salt intake (5%) than C2 (11%) and C4 (17%; Supplemental Table 4).

### **Sociodemographic, anthropometric and health characteristics by cluster**

Individuals in C1 were on average 4.5 years older than those in C4 ( $P < 0.05$ ; Table 2). Body weight was significantly lower in C1 than in C3, and in C2 compared with C3 and C4 ( $P < 0.05$ ). Individuals in C1 had 1.4 kg/m<sup>2</sup> lower BMI and 5.0 cm lower waist circumference than participants in C3 ( $P < 0.05$ ) and physical activity level was higher in C1 than C2 and C4 ( $P < 0.05$ ). Eleven per cent more individuals in C4 wanted to lose weight than in C1 ( $P < 0.05$ ; Table 2) and C4 was characterised by more current smokers than C1 ( $P < 0.05$ ). Twelve per cent more individuals in C1 than in C4 had a professional or managerial occupation, and similarly 7% more individuals had a manual occupation in C4 compared with C1 ( $P < 0.05$ ; Table 2). No other significant differences were observed (Table 2).

### **Changes in Healthy Eating Index by cluster after 6-month intervention**

Baseline and follow-up HEI scores and their components are presented in Table 3. There were no significant differences in changes in HEI between clusters for those randomised to non-personalised dietary advice. In contrast, for individuals who received PN advice (based

**Table 1** Food and nutrient and intakes by cluster of adherence to dietary recommendations at baseline among adults aged 18–79 years (*n* 1480), Food4Me study

	Cluster								<i>P</i> *
	C1 ( <i>n</i> 475)		C2 ( <i>n</i> 398)		C3 ( <i>n</i> 348)		C4 ( <i>n</i> 259)		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
<b>Dietary recommendations (g/d)</b>									
Oily fish	<b>48<sup>2,3,4</sup></b>	32	8	7	10	7	8	7	<b>&lt;0.001</b>
Whole grains	<b>183<sup>2,4</sup></b>	182	<b>216<sup>3,4</sup></b>	184	<b>205<sup>4</sup></b>	165	22	16	<b>&lt;0.001</b>
Red meat	<b>85<sup>2,3,4</sup></b>	81	<b>30<sup>3,4</sup></b>	20	<b>119<sup>4</sup></b>	53	84	96	<b>&lt;0.001</b>
Fruit and vegetables	<b>610<sup>2,3,4</sup></b>	371	<b>470<sup>3,4</sup></b>	303	456	288	339	218	<b>&lt;0.001</b>
<b>Other food intakes (g/d)</b>									
Fruit juice	<b>117<sup>3,4</sup></b>	181	114	165	94	144	76	108	<b>0.008</b>
Non-wholemeal	<b>116<sup>2,4</sup></b>	140	<b>78<sup>4</sup></b>	76	<b>114<sup>4</sup></b>	103	149	189	<b>&lt;0.001</b>
Eggs	<b>41<sup>2,3</sup></b>	41	22	24	31	47	30	51	<b>&lt;0.001</b>
Chicken, grilled or roast	<b>36<sup>2,3,4</sup></b>	37	<b>17<sup>3</sup></b>	21	28	25	25	27	<b>&lt;0.001</b>
White fish	<b>26<sup>2,3,4</sup></b>	26	10	14	13	14	11	14	<b>&lt;0.001</b>
Fish products	<b>19<sup>2,4</sup></b>	30	10 <sup>3</sup>	11	14	16	13	15	<b>&lt;0.001</b>
Beans and lentils	<b>30<sup>2,3</sup></b>	40	15	24	16	27	22	28	<b>&lt;0.001</b>
Butter	<b>4<sup>3</sup></b>	9	<b>6<sup>3</sup></b>	11	<b>9<sup>4</sup></b>	18	5	12	<b>0.005</b>
Low-fat dairy	<b>293<sup>2,3,4</sup></b>	296	217	203	221	212	173	219	<b>&lt;0.001</b>
High-fat dairy	64	120	60	119	83	113	83	204	0.44
Sugar-sweetened beverages	36	176	18	55	40	139	41	84	0.39
Low-calorie soft drinks	66	194	46	154	80	239	72	190	0.53
Added sugar	4	9	4	11	5	13	7	13	0.11
Chocolate and sweets	21	37	19	23	26	61	17	26	0.10
Cakes	22	31	18	25	20	25	22	39	0.08
Biscuits	30	55	21	37	35	88	27	55	0.38
Ice cream	7	19	6	11	7	12	7	13	0.62
Pastries	8	34	4	6	8	10	10	39	0.49
Crisps	4	10	<b>3<sup>3</sup></b>	5	5	10	4	8	0.06
Chips and pizza	30	41	<b>24<sup>3,4</sup></b>	22	35	30	34	35	<b>0.001</b>
Fried foods	<b>33<sup>2</sup></b>	52	<b>21<sup>3,4</sup></b>	28	34	35	33	30	<b>0.047</b>
<b>Nutrient intakes</b>									
Total energy (MJ/d)	<b>12.02<sup>2,4</sup></b>	5.10	<b>9.29<sup>3</sup></b>	3.12	<b>11.95<sup>4</sup></b>	4.46	8.82	4.10	<b>&lt;0.001</b>
Total energy (kcal/d)	<b>2870<sup>2,4</sup></b>	1219	<b>2218<sup>3</sup></b>	745	<b>2855<sup>4</sup></b>	1065	2106	978	<b>&lt;0.001</b>
EI:BMR	<b>1.9<sup>2,4</sup></b>	0.7	<b>1.5<sup>3</sup></b>	0.5	<b>1.8<sup>4</sup></b>	0.6	1.4	0.6	<b>&lt;0.001</b>
Total fat (% energy)	<b>36.0<sup>2,4</sup></b>	5.7	<b>34.1<sup>3,4</sup></b>	5.6	<b>36.4</b>	5.5	37.9	6.6	<b>&lt;0.001</b>
SFA (% energy)	<b>13.4<sup>3,4</sup></b>	2.8	<b>13.6<sup>3,4</sup></b>	3.3	<b>14.9</b>	3.0	15.3	3.3	<b>&lt;0.001</b>
MUFA (% energy)	<b>14.2<sup>2,3</sup></b>	3.2	<b>12.6<sup>3,4</sup></b>	2.8	<b>13.6<sup>4</sup></b>	2.6	14.8	3.5	<b>&lt;0.001</b>
PUFA (% energy)	<b>6.0<sup>2,4</sup></b>	1.4	5.7	1.4	5.6	1.3	5.5	1.7	<b>0.003</b>
Protein (% energy)	<b>18.3<sup>2,3,4</sup></b>	4.1	<b>15.5<sup>3,4</sup></b>	3.2	17.0	2.9	17.3	3.7	<b>&lt;0.001</b>
Carbohydrate (% energy)	<b>44.5<sup>2,3</sup></b>	7.5	<b>49.6<sup>3,4</sup></b>	7.0	45.6	6.4	43.7	8.3	<b>&lt;0.001</b>
Sugars (% energy)	<b>21.0<sup>2</sup></b>	5.9	<b>22.5<sup>3,4</sup></b>	6.1	19.8	5.6	20.8	5.9	<b>&lt;0.001</b>
Dietary fibre (g/d)†	<b>34.0<sup>2,4</sup></b>	15.8	<b>30.2<sup>4</sup></b>	14.4	<b>31.7<sup>4</sup></b>	12.8	18.7	8.2	<b>&lt;0.001</b>
Salt (g/d)†	<b>8.3<sup>2,4</sup></b>	4.0	<b>6.1<sup>3</sup></b>	2.7	<b>8.7<sup>4</sup></b>	3.6	5.9	3.6	<b>&lt;0.001</b>

EI:BMR, ratio of energy intake to BMR.

\*ANOVA was used to test for significant differences across clusters. Models were adjusted for age, sex, BMI, physical activity level, smoking habits and country; significant *P* values are highlighted in bold. *Post hoc* Tukey tests were performed to test for significant differences between clusters; superscript numbers denote where the differences lie across the clusters. For example, 2 indicates that the mean value is significantly different from that in cluster 2 (*P*<0.05).†*P* values are also adjusted for total energy intake.

on information of current diet alone or combined with information on phenotype and genotype), changes in HEI differed between clusters (*P*<0.001). There were bigger improvements in HEI for participants in C4 compared with C1 and C2 (*P*<0.05) and in C2 compared with C4 (*P*<0.05; Fig. 1). There were no significant differences in changes in HEI between clusters when PN was stratified by Level 1, Level 2 or Level 3 (data not shown).

### Sensitivity analyses

Exclusion of participants with reported intakes more than 3 SD above or below the mean dietary intakes of whole grains,

oily fish, red meat and fruit and vegetables revealed similar clusters (see online supplementary material, Supplemental Table 5). The pattern of the main results remained the same, with individuals in C3 and C4 making greater changes in HEI at month 6 than those in C1, as did participants in C4 compared with those in C2 (*P*<0.05).

## Discussion

### Main findings

Based on our secondary analysis in the Food4Me Proof-of-Principle study, we identified four distinct clusters of

**Table 2** Sociodemographic characteristics by cluster of adherence to dietary recommendations at baseline among adults aged 18–79 years (*n* 1480), Food4Me study

	Cluster								<i>P</i> *
	C1 ( <i>n</i> 475)		C2 ( <i>n</i> 398)		C3 ( <i>n</i> 348)		C4 ( <i>n</i> 259)		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Age (years)	41.2 <sup>2,3</sup>	12.7	39.2 <sup>4</sup>	14.2	41.2 <sup>4</sup>	12.7	36.7	11.5	<0.001
Female (%)	56.0		67.3		47.4		64.1		0.79
Ethnicity (%)									
Caucasian	95.6		96.5		97.7		98.1		0.16
Occupation (%)									
Professional and managerial	44.2 <sup>4</sup>		37.8		39.4		32.2		<b>0.014</b>
Intermediate occupations	25.9		22.4		28.5		28.7		0.16
Routine and manual	7.4 <sup>4</sup>		6.8		12.9		14.3		<b>0.006</b>
Student	13.5		21.7		9.5		14.7		0.18
Not currently working	9.1		11.3		9.8		10.1		0.38
Anthropometrics									
Body weight (kg)	74.6 <sup>3</sup>	15.1	70.5 <sup>3,4</sup>	15.0	80.3 <sup>4</sup>	16.0	74.1	16.3	<0.001
BMI (kg/m <sup>2</sup> )	25.4 <sup>2,3</sup>	4.4	24.1 <sup>3,4</sup>	4.4	26.8	4.9	26.0	5.7	<0.001
Waist circumference (cm)	85.4 <sup>3</sup>	13.0	81.8	13.2	90.4	14.1	85.9	14.1	<0.001
Physical activity									
Physical activity level	1.8 <sup>2,4</sup>	0.2	1.7 <sup>3</sup>	0.2	1.8 <sup>4</sup>	0.2	1.7	0.2	<0.001
Sedentary behaviour (min/d)	746	73	742	77	750	76	744	7	0.96
Dietary conditions (%)									
Want to lose weight	46.1 <sup>4</sup>		41.2		48.6		57.5		<b>0.013</b>
Restricted diet	6.1		11.6		3.7		5.8		0.47
Medication use (%)									
Prescribed medication	26.1		35.7		29.9		27.0		0.79
Non-prescribed medication	8.6		10.6		9.2		11.2		0.18
Health and disease									
Current smoker (%)	9.8 <sup>4</sup>		9.0		10.3		22.0		<b>0.005</b>
Total cholesterol (mmol/l)	4.6	0.9	4.5	1.0	4.7	1.0	4.6	0.9	0.09
High blood pressure (%)	8.2		7.0		9.8		5.8		0.89
Heart disease (%)	2.1		1.8		0.6		1.2		0.17

Values are presented means and standard deviations or percentages.

\*ANOVA and logistic regression were used to test for significant differences across clusters in continuous and categorical variables, respectively. Analyses were adjusted for age, sex, BMI, physical activity level, smoking habits and country; significant *P* values are highlighted in bold. *Post hoc* Tukey tests (continuous data) and logistic regression (categorical) were used to test for significant differences between clusters; superscript numbers denote where the differences lie across the clusters. For example, 2 indicates that the mean value or percentage is significantly different from that in cluster 2 (*P* < 0.05).

individuals according to their adherence to current European dietary recommendations. Individuals in C1 and C2 met more dietary recommendations than those in C3 and C4. Moreover, on average, individuals in C1 and C2 had a healthier diet, lower BMI and waist circumference, and smoked less, compared with those in C3 and C4. When randomised to a 6-month PN intervention, participants in C4 made the greatest improvements in their diets (as estimated by HED), compared with participants receiving non-personalised 'one size fits all' generalised advice. The present study is the first to investigate clusters of adherence to European dietary recommendations and to determine the responsiveness of cluster members to PN advice.

### Comparison with other studies

Previous studies have used cluster analysis to categorise individuals<sup>(31)</sup>. We used cluster analysis to categorise individuals based on their adherence to current European food-based dietary guidelines at baseline for participants in the Food4Me intervention study. This approach identified groups of individuals who differed in the number, and

groupings, of dietary recommendations they met. Clusters where more individuals met the recommendations were characterised by being slightly older and in more highly educated occupations, which is a well-established characteristic of healthy dietary clusters<sup>(32)</sup>.


Clustering of dietary intakes and adequacies has been investigated in relation to several health outcomes<sup>(7,8,33)</sup> and can be strong predictors of these outcomes<sup>(34)</sup>. A recent review of dietary clusters and health outcomes by the US Department of Agriculture<sup>(35)</sup> concluded that the strongest evidence for an association between unhealthy dietary patterns and increased disease risk is for CVD, followed by obesity and then type 2 diabetes. The review concluded that there was a lack of studies assessing dietary intakes at follow-up and using a universal and quantitative indicator of dietary intake. Our study is in line with these recommendations as we utilised the HEI, which is a validated estimate of dietary adequacy, and we assessed dietary change using the same instrument at both baseline and follow-up. Although more limited, some prospective and RCT studies have investigated the effect of clustering on changes in health outcomes<sup>(12,36,37)</sup> and

**Table 3** Healthy Eating Index 2010 (HEI) score and its constituents at baseline and month 6 by cluster of adherence to dietary recommendations among adults aged 18–79 years (*n* 1480), Food4Me study

	Cluster								<i>P</i> *
	C1 ( <i>n</i> 475)		C2 ( <i>n</i> 398)		C3 ( <i>n</i> 348)		C4 ( <i>n</i> 259)		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
<b>Baseline score</b>									
Total HEI	53.3 <sup>2,3,4</sup>	8.9	50.5 <sup>3,4</sup>	8.9	47.5 <sup>4</sup>	8.9	41.8	10.1	<0.001
Fatty acid ratio†	3.2 <sup>2,3,4</sup>	2.4	2.2 <sup>3,4</sup>	2.4	1.7	1.7	2.0	2.0	<0.001
Protein	3.7 <sup>2,3,4</sup>	0.7	3.2 <sup>3,4</sup>	0.6	3.5	0.6	3.5	0.7	<0.001
Salt	0.1	0.5	0.1	0.7	0.1	0.6	0.1	0.6	0.002
Empty calories	8.8 <sup>2</sup>	4.0	7.7	4.3	8.5	3.8	7.5	4.1	0.012
Refined grains	6.1 <sup>2,3,4</sup>	3.7	4.8	3.7	4.4	3.7	4.7	4.0	<0.001
Seafood and plant protein	5.0 <sup>2,3,4</sup>	0.2	4.5	1.0	4.3	1.1	4.4	1.1	<0.001
Fruit	3.8 <sup>3</sup>	1.3	3.8 <sup>3</sup>	1.4	3.3	1.5	3.3	1.5	<0.001
Whole fruit	4.2 <sup>3,4</sup>	1.3	4.1 <sup>3,4</sup>	1.3	3.6	1.5	3.6	1.6	<0.001
Vegetables	2.5 <sup>3,4</sup>	1.1	2.3 <sup>3,4</sup>	1.1	2.0	0.9	2.1	1.1	<0.001
Greens and beans	4.2 <sup>2,3,4</sup>	1.1	3.8 <sup>3</sup>	1.3	3.5	1.3	3.7	1.4	<0.001
Whole grains	7.3	3.5	9.5	1.2	8.8	1.9	2.9	2.2	<0.001
Dairy products	4.7 <sup>2,3,4</sup>	2.6	4.7 <sup>4</sup>	2.7	4.3 <sup>4</sup>	2.2	4.4	2.7	0.27
<b>Follow-up score</b>									
Total HEI	55.7 <sup>2,3,4</sup>	9.1	53.3 <sup>4</sup>	9.6	51.4	8.7	48.0	10.3	<0.001
Fatty acid ratio†	3.8 <sup>2,3,4</sup>	2.6	3.1 <sup>3</sup>	2.7	2.5	2.1	2.6	2.2	<0.001
Protein	3.8 <sup>2,3,4</sup>	0.7	3.3 <sup>3,4</sup>	0.6	3.6	0.6	3.6	0.6	<0.001
Salt	0.1	0.6	0.2 <sup>3</sup>	0.9	0.1	0.6	0.1	0.6	0.002
Empty calories	8.7 <sup>2</sup>	4.0	7.4	4.1	8.8	4.0	8.1	4.1	0.002
Refined grains	6.2 <sup>4</sup>	3.8	5.4	3.8	5.1	3.8	4.9	3.8	0.004
Seafood and plant protein	5.0 <sup>2,3</sup>	0.2	4.7	0.8	4.6	1.0	4.7	0.9	<0.001
Fruit	4.1	1.3	4.2 <sup>3</sup>	1.2	3.7	1.4	3.7	1.5	0.009
Whole fruit	4.4	1.2	4.4	1.1	4.1	1.4	4.0	1.5	0.023
Vegetables	2.8 <sup>3,4</sup>	1.2	2.7 <sup>3,4</sup>	1.3	2.3	1.0	2.4	1.0	<0.001
Greens and beans	4.3 <sup>2,3</sup>	1.0	4.0	1.2	3.9	1.2	4.1	1.2	0.001
Whole grains	7.9 <sup>2,3,4</sup>	3.1	9.2 <sup>4</sup>	1.9	8.5 <sup>4</sup>	2.7	5.5	3.7	<0.001
Dairy products	4.8	2.7	4.7	2.8	4.4	2.3	4.5	2.6	0.52

\*Fatty acid ratio is the ratio of unsaturated fatty acids (mono- and polyunsaturated fatty acids) to saturated fatty acids.

\*ANOVA was used to test for significant differences across clusters. Models were adjusted for age, sex, BMI, physical activity level, smoking habits and country; significant *P* values are highlighted in bold. *Post hoc* Tukey tests were used to test for significant differences between clusters; superscript numbers denote where the differences lie across the clusters. For example, 2 indicates that the mean value is significantly different from that in cluster 2 (*P* < 0.05).



**Fig. 1** Changes from baseline to month 6 in Healthy Eating Index 2010 (HEI) score by cluster of adherence to dietary recommendations at baseline among adults aged 18–79 years (*n* 1480), Food4Me study. Values are predicted means with their standard errors represented by vertical bars. Models were adjusted for age, sex, BMI, physical activity level, smoking habits and country ( $P_{\text{trend}} < 0.001$ ); *post hoc* Tukey tests were used to test for significant differences between clusters (C); C4 > C1 (*P* < 0.001), C3 > C1 (*P* = 0.005)

some studies have used adherence to dietary recommendations to derive clusters<sup>(12–14,38,39)</sup>. Dietary recommendations used in studies included in the systematic review by the US Department of Agriculture<sup>(35)</sup> varied according to the study, but all included a measure of fruit and vegetable, wholegrain and meat intake.

To our knowledge, no previous research has evaluated the impact of clustering of dietary recommendations on the response to a PN intervention. We observed that individuals in the cluster where the fewest recommendations were met (C4) reported the biggest improvement in HEI following PN intervention but there were no differences between clusters in response to conventional, non-personalised dietary advice. Given that adverse lifestyle behaviours and the prevalence and risk of death from obesity-related diseases are strongly socio-economically patterned<sup>(40)</sup>, it is important that appropriate interventions are targeted to those most in need of improved lifestyle. While research on the development and implementation of PN interventions and their effects on changing diets is in its infancy<sup>(41)</sup>, the findings from the present study provide

encouragement that PN interventions can be more effective than current 'one size fits all' interventions and that they may be particularly effective among individuals with the poorest diets. There have been concerns that PN may be taken up only by the 'worried well'<sup>(42)</sup>, who already have adequate dietary intakes. However, our findings suggest that PN is most effective in people who have the least adequate diets, and therefore the greatest need for improvement in dietary intakes, with the potential for significant reductions in disease risk.

### **Strengths and limitations**

The present study had a number of strengths. Our findings are derived from a relatively large number of participants who were broadly representative of European adults from seven different European countries<sup>(18)</sup>. The Food4Me RCT collected extensive information on anthropometrics, physical activity and sociodemographic and health-related data, which contributed to detailed characterisation of participants in the clusters. Our study design allowed us to estimate changes in dietary intakes using the same validated instrument at baseline and at month 6. Furthermore, we quantified responses using the HEI, which has been shown to be an effective indicator of overall diet quality<sup>(26)</sup> and, therefore, a better measure of overall dietary change than outcomes based on single foods or nutrients.

A limitation of the study is that our data were self-reported via the Internet, which may have introduced measurement error. However, the validity of Internet-based, self-reported anthropometric data is high<sup>(43)</sup> and has been confirmed in the present study<sup>(27)</sup>. We were not able to include dairy products as a dietary recommendation in the present analyses due to so few individuals meeting the recommendation. However, dairy products do not have a recommended intake in the UK and so habitual diets would not necessarily be expected to comply with this recommendation, even if they were very health conscious. Dietary intakes were estimated by an FFQ, which is known to be subject to misreporting error<sup>(44)</sup>, but this was minimised by validating our FFQ against a 4 d weighed food record<sup>(20)</sup>. Moreover, our estimation of dietary change was based on the HEI, which is a validated indicator of overall diet<sup>(26)</sup> and which may be less susceptible to reporting errors than approaches measuring change in specific nutrients or individual foods. Our study participants were almost solely Caucasian; thus, further research in wider ethnicity groups is required to generalise our findings to other populations. One of the primary aims of the Food4Me Proof-of-Principle study was to evaluate change in intakes of food groups across four treatment arms. Thus, although the present study is a secondary analysis of these data, clustering was based on how individuals adhered to food group recommendations and included four clusters. As a result, our analyses are likely to be powered to detect differences between clusters.

### **Implications of findings**

Our findings suggest that the efficacy of PN in modifying dietary intakes depends on the clustering of adherence to dietary recommendations, with those with the poorest diets benefiting most from the PN intervention. As a result, the implementation of PN-based interventions in individuals with the least healthy diets may help to address health inequalities. Understanding the characteristics of individuals within coherent clusters which are linked with their responsiveness to interventions may help in the design and implementation of more effective health promotion actions. Future PN interventions may benefit from tailoring PN advice based on clustering of overall dietary behaviours rather than on single nutrients or foods.

### **Conclusions**

We identified four distinct clusters of individuals based on adherence to current food-based dietary recommendations. Participants in the cluster where the fewest recommendations were met (C4) reported significantly greater improvements in their diets (as estimated by the HEI) following a 6-month trial of PN, whereas there was no difference between clusters for those randomised to the Control, non-personalised dietary intervention.

### **Acknowledgements**

*Acknowledgements:* The authors would like to thank Antoneta Granic for her valuable input into the design of the cluster analysis. J.A.M. and S.N.-C. are grateful to CIBERobn Fisiopatología de la Obesidad y Nutrición (Instituto Carlos III, Madrid, Spain) for general support in research. *Financial support:* This work was supported by the European Commission under the Food, Agriculture, Fisheries and Biotechnology Theme of the 7th Framework Programme for Research and Technological Development (265494). The European Commission had no role in the design, analysis or writing of this article. *Conflict of interest:* None of the authors had a personal or financial conflict of interest. *Authorship:* K.M.L. and C.C.-M. are joint first authors. Author responsibilities were as follows: Y.M., I.T., C.A.D., E.R.G., L.B., J.A.L., J.A.M., W.H.M.S., H.D., M.G. and J.C.M. contributed to the research design. J.C.M. was the Food4Me Proof-of-Principle study leader. C.C.-M., C.F.M.M., H.F., C.B.O., C.W., A.L.M., R.F., S.N.-C., R.S.-C., S.K., L.T., C.P.L., A.S., M.C.W., E.R.G., L.B. and J.C.M. contributed to the developing the standardised operating procedures for the study. C.C.-M., S.N.-C., R.S.-C., C.W., C.B.O., H.F., C.F.M.M., A.L.M., R.F., S.K., L.T., C.P.L., A.S., M.C.W. and J.C.M. conducted the intervention. C.C.-M., C.F.M.M. and W.H.M.S. contributed to physical activity measurements. K.M.L. and C.C.-M. wrote the paper, performed the statistical analysis for the manuscript.

All authors contributed to a critical review of the manuscript during the writing process. All authors approved the final version to be published. *Ethics of human subject participation*: This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects/patients were approved by the Research Ethics Committees at each University or Research Centre delivering the intervention. The Food4Me trial was registered as an RCT (NCT01530139) at Clinicaltrials.gov. All participants expressing an interest in the study were asked to sign online consent forms at two stages in the screening process. These consent forms were automatically directed to the local study investigators to be counter-signed and archived.

### Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1368980016001932>

### References

- Ng M, Fleming T, Robinson M *et al.* (2014) Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* **384**, 766–781.
- Hill JO, Wyatt HR & Peters JC (2012) Energy balance and obesity. *Circulation* **126**, 126–132.
- NHS Choices (2015) Live Well – Healthy living for everyone. <http://www.nhs.uk/Livewell/Pages/Livewellhub.aspx> (accessed March 2015).
- Hood L & Friend SH (2011) Predictive, personalized, preventive, participatory (P4) cancer medicine. *Nat Rev Clin Oncol* **8**, 184–187.
- Nielsen DE & El-Sohehy A (2014) Disclosure of genetic information and change in dietary intake: a randomized controlled trial. *PLoS ONE* **9**, e112665.
- Berrigan D, Dodd K, Troiano RP *et al.* (2003) Patterns of health behavior in US adults. *Prev Med* **36**, 615–623.
- Leech R, McNaughton S & Timperio A (2014) The clustering of diet, physical activity and sedentary behavior in children and adolescents: a review. *Int J Behav Nutr Phys Act* **11**, 4.
- Newby PK, Muller D, Hallfrisch J *et al.* (2003) Dietary patterns and changes in body mass index and waist circumference in adults. *Am J Clin Nutr* **77**, 1417–1425.
- Moore L, Singer M, Bradlee ML *et al.* (2015) Adolescent dietary intakes predict cardiometabolic risk clustering. *Eur J Nutr* **55**, 461–468.
- Khaw K-T, Wareham N, Bingham S *et al.* (2008) Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk Prospective Population Study. *PLoS Med* **5**, e12.
- van Dam RM, Li T, Spiegelman D *et al.* (2008) Combined impact of lifestyle factors on mortality: prospective cohort study in US women. *BMJ* **337**, a1440.
- Kesse-Guyot E, Castetbon K, Estaquio C *et al.* (2009) Association between the French Nutritional Guideline-based score and 6-year anthropometric changes in a French middle-aged adult cohort. *Am J Epidemiol* **170**, 757–765.
- Chiuve SE, Fung TT, Rexrode KM *et al.* (2011) Adherence to a low-risk, healthy lifestyle and risk of sudden cardiac death among women. *JAMA* **306**, 62–69.
- Zamora D, Gordon-Larsen P, He K *et al.* (2011) Are the 2005 dietary guidelines for Americans associated with reduced risk of type 2 diabetes and cardiometabolic risk factors? Twenty-year findings from the CARDIA study. *Diabetes Care* **34**, 1183–1185.
- Livingstone KM, Celis-Morales C, Navas-Carretero S *et al.* (2016) Effect of an Internet-based, personalized nutrition randomized trial on dietary changes associated with the Mediterranean diet: the Food4Me Study. *Am J Clin Nutr* (Epublication ahead of print version).
- Food4Me (2016) An integrated analysis of opportunities and challenges for personalised nutrition. <http://www.food4me.org/> (accessed February 2016).
- Celis-Morales C, Livingstone KM, Marsaux CFM *et al.* (2015) Design and baseline characteristics of the Food4Me study: a web-based randomised controlled trial of personalised nutrition in seven European countries. *Genes Nutr* **10**, 450.
- Livingstone KM, Celis-Morales C, Navas-Carretero S *et al.* (2016) Profile of European adults interested in internet-based personalised nutrition: the Food4Me study. *Eur J Nutr* **55**, 759–769.
- Forster H, Fallaize R, Gallagher C *et al.* (2014) Online dietary intake estimation: the Food4Me food frequency questionnaire. *J Med Internet Res* **16**, e150.
- Fallaize R, Forster H, Mcready AL *et al.* (2014) Online dietary intake estimation: reproducibility and validity of the Food4Me food frequency questionnaire against a 4-day weighed food record. *J Med Internet Res* **16**, e190.
- Food Standards Agency (2002) *McCance and Widdowson's The Composition of Foods*, sixth summary edition ed. Cambridge: Royal Society of Chemistry.
- Institute of Medicine (2005) Dietary Reference Intakes for Energy, Carbohydrate, Fibre, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. <http://www.nap.edu/openbook.php?isbn=0309085373> (accessed March 2015).
- Institute of Medicine (2011) Dietary Reference Intakes Tables and Applications. <http://www.iom.edu/Activities/Nutrition/SummaryDRIs/DRI-Tables.aspx> (accessed March 2015).
- World Health Organization (2007) *Protein and Amino Acid Requirements in Human Nutrition. Report of a Joint WHO/FAO/UNU Expert Consultation*. WHO Technical Report Series no. 935. Geneva: WHO.
- Food and Agriculture Organization of the United Nations (2010) Fats and fatty acids in human nutrition. Report of an expert consultation. [http://www.who.int/nutrition/publications/nutrientrequirements/fatsandfattyacids\\_human\\_nutrition/en/](http://www.who.int/nutrition/publications/nutrientrequirements/fatsandfattyacids_human_nutrition/en/) (accessed March 2016).
- Guenther PM, Casavale KO, Reedy J *et al.* (2013) Update of the Healthy Eating Index: HEI-2010. *J Acad Nutr Diet* **113**, 569–580.
- Celis-Morales C, Livingstone KM, Woolhead C *et al.* (2015) How reliable is internet-based self-reported identity, socio-demographic and obesity measures in European adults? *Genes Nutr* **73**, 476.
- European Commission (2015) European skills, competences, qualifications and occupations. <https://ec.europa.eu/esco/web/guest/hierarchybrowser/-/browser/occupation> (accessed April 2015).
- European Commission (2015) Mean annual earnings by sex, age and occupation. [http://ec.europa.eu/eurostat/web/products-datasets/-/earn\\_ses\\_agt28](http://ec.europa.eu/eurostat/web/products-datasets/-/earn_ses_agt28) (accessed March 2015).
- Chiu T, Fang D, Chen J *et al.* (2001) A robust and scalable clustering algorithm for mixed type attributes in large database environment. In *Proceedings of the Seventh ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pp. 263–268. San Francisco, CA: ACM.

31. Ocké MC (2013) Evaluation of methodologies for assessing the overall diet: dietary quality scores and dietary pattern analysis. *Proc Nutr Soc* **72**, 191–199.
32. Kant AK (2004) Dietary patterns and health outcomes. *J Am Diet Assoc* **104**, 615–635.
33. Ma Y, Bertone ER, Stanek EJ *et al.* (2003) Association between eating patterns and obesity in a free-living US adult population. *Am J Epidemiol* **158**, 85–92.
34. Wirt A & Collins CE (2009) Diet quality – what is it and does it matter? *Public Health Nutr* **12**, 2473–2492.
35. US Department of Agriculture (2014) A series of systematic reviews on the relationship between dietary patterns and health outcomes. <http://www.nel.gov/vault/2440/web/files/DietaryPatterns/DPRptFullFinal.pdf> (accessed July 2016).
36. Gao SK, Beresford SAA, Frank LL *et al.* (2008) Modifications to the Healthy Eating Index and its ability to predict obesity: the Multi-Ethnic Study of Atherosclerosis. *Am J Clin Nutr* **88**, 64–69.
37. Jacobs DR, Sluik D, Rokling-Andersen MH *et al.* (2009) Association of 1-y changes in diet pattern with cardiovascular disease risk factors and adipokines: results from the 1-y randomized Oslo Diet and Exercise Study. *Am J Clin Nutr* **89**, 509–517.
38. Cheng G, Gerlach S, Libuda L *et al.* (2010) Diet quality in childhood is prospectively associated with the timing of puberty but not with body composition at puberty onset. *J Nutr* **140**, 95–102.
39. Park S-Y, Murphy SP, Wilkens LR *et al.* (2005) Dietary patterns using the food guide pyramid groups are associated with sociodemographic and lifestyle factors: the Multiethnic Cohort Study. *J Nutr* **135**, 843–849.
40. Di Cesare M, Khang Y-H, Asaria P *et al.* (2013) Inequalities in non-communicable diseases and effective responses. *Lancet* **381**, 585–597.
41. Celis-Morales C, Lara J & Mathers JC (2014) Personalising nutritional guidance for more effective behaviour change. *Proc Nutr Soc* **74**, 130–138.
42. Ferguson L (editor) (2013) *Nutrigenomics and Nutrigenetics in Functional Foods and Personalized Nutrition*. Boca Raton, FL: CRC Press.
43. Pursey K, Burrows LT, Stanwell P *et al.* (2014) How accurate is web-based self-reported height, weight, and body mass index in young adults? *J Med Internet Res* **16**, e4.
44. Macdiarmid J & Blundell J (1998) Assessing dietary intake: who, what and why of under-reporting. *Nutr Res Rev* **11**, 231–253.