

49 adolescents meeting ICD 10 criteria for anorexia nervosa or atypical anorexia nervosa. The same diagnostic interview was administered twice, once via face-to-face and once as an online version, within a week. Method order was counterbalanced among participants and temporal stability was controlled. The Eating Disorder Inventory-2 (EDI-2) was used as a control variable.

Results: Both the equivalence test and the null hypothesis test were significant for the sum score of the EDE. Measures of psychopathology in eating disorders demonstrated equivalence across face-to-face and online format of the EDE.

Conclusions: The aim was to examine the equivalence of face-to-face and online methodologies, controlling for temporal change in the variable under investigation over one week and order of administration. Results demonstrate equivalence across face-to-face and online format of the EDE. These findings suggest that data obtained using EDE online can be interpreted in comparison with normative data obtained in the face-to-face Interview and that corrections through transformations are not necessary.

Disclosure of Interest: None Declared

EPV0902

Psychiatric role in physician-assisted death requests – a study protocol for a literature review

R. Barranha^{1,2*}, A. R. Ferreira^{2,3}, E. Gonçalves^{4,5} and L. Fernandes^{1,2,3}

¹Psychiatry Service, Centro Hospitalar Universitário de São João;

²Department of Clinical Neurosciences and Mental Health;

³CINTESIS@RISE, Faculty of Medicine, University of Porto;

⁴Palliative Care Service, Centro Hospitalar Universitário de São João

and ⁵Department of Community Medicine, Information and Health Decision Sciences (MEDCIDS), Faculty of Medicine, University of Porto, Porto, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2201

Introduction: The prospect of a medium-term approval of physician-assisted death in Portugal raises relevant ethical and deontological issues that need to be addressed, namely the framework of psychiatric assessments in this process. Such assessments are undermined by the lack of scientific precision in the methods used to determine decision-making capacity, making it possible for the final decision to be affected by psychiatrists' personal beliefs. As such, outlining scientific evidence and legislation pieces defining the psychiatrists' role and scope is of utmost importance to frame this debate.

Objectives: To synthesize the accumulated evidence worldwide regarding the psychiatrists' involvement in the global process of physician-assisted death requests by reviewing scientific literature, published protocols, official reports and international promulgated or amended legislation related to hasten death practices.

Methods: *PubMed*, *Scopus*, *Web of Science*, *PsycInfo* and *Google Scholar* electronic bibliographic databases will be searched for eligible articles, as well as grey literature, using the following search terms: Psychiatry AND (Euthanasia OR (Suicide AND Assisted)). Official governments' and countries health authorities' websites will also be searched for relevant reports and legislation documents,

as well as right-to-die organizations and akin associations. No language, date of publication, or geographical restrictions will be applied. The full text of potentially relevant results will be retrieved from the different sources for review after screening titles and abstracts. This two-stage process will be conducted independently by two researchers. Outcomes of interest will be the descriptions of psychiatric role in the process of physician-assisted death requests, assessment methods, and measurement techniques used.

Results: Given the fact that physician-assisted death is legalized only in a few jurisdictions, we believe the number of eligible results will be limited. Data will be extracted and a descriptive summary of the evidence will be provided. We anticipate finding a significant variability, but also to identify points of consensus. The findings will be published in a peer-review indexed journal and presented at national and international conferences.

Conclusions: To our knowledge, this is the first review of both, scientific published literature, and international legislation on the role of psychiatrists in physician-assisted death requests. We hope to provide an international overview to frame the public debate by pinpointing the most consensual assessment methodology, allowing to design an optimized assessment protocol before the implementation of the law in Portugal.

Disclosure of Interest: None Declared

Schizophrenia and other psychotic disorders

EPV0903

Folie a deux / induced delusional disorder – case report and literature review

A. Lourenço*, A. L. Falcão, G. Soares, J. Petta, C. Rodrigues, M. Nascimento and C. Oliveira

Clínica 3, Centro Hospitalar Psiquiátrico de Lisboa, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2202

Introduction: *Folie a deux*, also known as shared psychotic disorder or induced delusional disorder, is a rare mental disorder that was first described in France in the late 19th century and was referred to delusions shared between two individuals in close relationship. The concept has evolved and according to ICD-10 the following criteria for the diagnosis is phenomenology-based-only.

Objectives: To describe a clinical case and review the existing evidence on *folie a deux*.

Methods: Clinical case and non-systematic review of the literature, from the last 15 years, on *folie a deux*. For this research, the keywords “*folie a deux*”, “shared psychotic disorder” and “induced delusional disorder” were used in the MEDLINE/PubMed database.

Results: The clinical case presented refers to a 56-year-old female patient with no known psychiatric history. The patient stated that 5 years ago when his mother died, neighbors began to persecute her and her sister. She was medicated with a second-generation anti-psychotic without total remission of symptoms. Generally, in *folie a deux* there is a close and prolonged relationship between the