

Short Communication

Increased dietary protein consumed at breakfast leads to an initial and sustained feeling of fullness during energy restriction compared to other meal times

Heather J. Leidy^{1,2*}, Mandi J. Bossingham¹, Richard D. Mattes¹ and Wayne W. Campbell¹

¹Department of Foods & Nutrition, Ingestive Behavior Research Center, Purdue University, West Lafayette, IN 47907, USA

²Department of Dietetics & Nutrition, The University of Kansas Medical Center, 3901 Rainbow Blvd, Kansas City, KS 66160, USA

(Received 28 March 2008 – Revised 2 June 2008 – Accepted 23 June 2008 – First published online 2 September 2008)

The objective of the study was to assess whether the timing of increased dietary protein throughout the day influences the feelings of fullness during energy balance (EB) and restriction (ER). Nine men (age 48 (SEM 6) years; BMI 32.7 (SEM 0.7) kg/m²) randomly completed five controlled feeding trials, each consisting of 3 d of EB, followed by 3 d of ER of a 3138 kJ/d (750 kcal/d) reduction). The diet was composed of a normal amount of protein (NP) (0.8 g protein/kg per d), or an additional amount of protein (HP) (+0.6 g protein/kg per d) given at breakfast (HP-B), lunch (HP-L), dinner (HP-D) or equally divided among all meals (HP-E). Meal-related (3 h postprandial) and overall (15 h composite) feelings of fullness were assessed from thirteen-point, numbered, linear category scale questionnaires (reported as arbitrary units (au)). When comparing HP treatments, the data are presented as difference from NP. No differences in meal-related or overall fullness were observed among HP treatments during EB. During ER, the HP-B led to greater meal-related fullness (+137 (SEM 44) au × 180 min) compared to HP-D (−1 (SEM 37) au × 180 min; *P*=0.003), but not for HP-L (+62 (SEM 53) au × 180 min; *P*=0.188) or HP-E-B (+92 (SEM 85) au × 180 min; *P*=0.587). HP-B also led to greater overall (15 h) fullness (+404 (SEM 162) au × 900 min) v. HP-L (+33 (SEM 162) au × 900 min; *P*=0.009) and HP-D (−60 (SEM 132) au × 900 min; *P*=0.05), but not HP-E (+274 (SEM 165) au × 900 min; *P*=0.188). The initial and sustained feelings of fullness following protein consumption at breakfast suggests that the timing of protein intake differentially influences satiety during ER.

Satiety: Breakfast: Dietary protein

The long-standing parental advice to ‘eat your breakfast; it’s the most important meal of the day’ has recently acquired scientific support as breakfast skipping is associated with an array of unhealthful outcomes⁽¹⁾ including body weight gain, overweight and obesity^(2–5). Several recent studies have shown that men and women consume more total energy throughout the day, especially in the evening, when breakfast is skipped^(2,6). The types of foods consumed at breakfast may also impact total energy intake. Dietary protein is the most satiating of the macronutrients during energy restriction (ER) and energy balance (EB) conditions⁽⁷⁾. We have previously reported that acute consumption of a higher dietary protein (28 g/meal) breakfast during ER leads to reduced hunger and desire to eat and increased feelings of fullness compared to consumption of a normal amount of dietary protein (17 g/meal)⁽⁸⁾. Data are limited concerning whether the consumption of additional dietary protein at breakfast leads to a greater differential appetitive response compared

to other meal times or when protein is given at each meal. The purpose of the present study was to examine whether increased dietary protein intake at breakfast, lunch, dinner, or dividing it equally across meals influences short-term self-reported fullness during EB and ER.

Materials and methods

Participants

Participants were recruited through newspaper advertisements. Eligibility was based on the following criteria: (1) men ≥21 years; (2) BMI between 25 and 39.9 kg/m²; (3) not dieting and no weight loss or gain (>4.5 kg) within the last 6 months; (4) non-smoking; and (5) non-diabetic. Ten men began the study; nine men completed all study procedures (48 (SEM 6) years; BMI 32.7 (SEM 0.7) kg/m²). Each subject was informed of the purpose, procedures and potential risks of participation in the

Abbreviations: au, arbitrary units; EB, energy balance; ER, energy restriction; HP, higher protein; HP-B, HP breakfast; HP-D, HP dinner; HP-E, HP equally divided among all meals; HP-L, HP lunch; NP, normal protein.

* **Corresponding author:** Dr Heather J. Leidy, The University of Kansas Medical Center, fax +1 913 588 8946, email hleidy@kumc.edu

study before signing an informed consent form approved by the University Biomedical Institutional Review Board.

Experimental design

A randomized cross-over design was utilized in which each subject completed the following five controlled feeding trials: normal protein (NP), higher protein (HP) breakfast (HP-B), lunch (HP-L), dinner (HP-D) and HP equally divided among all meals (HP-E). Each trial consisted of 6 d. During the first 3 d (days 1–3) of each trial, each subject was provided with an EB diet. During the second 3 d (days 4–6), each subject was provided with an individualized ER diet. During day 7, the subjects were asked to follow their habitual energy intake. In each trial, a three meal per day pattern was provided. Appetite was assessed throughout the 15 h period during days 3 and 6.

Diet

The subjects' energy needs were estimated to be 1.5 times their resting energy expenditure calculated using the Harris–Benedict equation for men⁽⁹⁾. The ER diet was then determined as 3138 kJ/d (750 kcal/d) less than their estimated energy need. The NP-EB diet consisted of 11% of total energy from protein, 64% of total energy from carbohydrate and 25% of total energy from fat, whereas the NP-ER diet consisted of 14% of total energy from protein, 61% from carbohydrate and 25% from fat. Both NP diets contained 0.8 g protein/kg per d equally divided among breakfast, lunch and dinner and was void of eggs and all striated tissue, including pork. The HP-EB diet was composed of 18% of total energy from protein, 57% from carbohydrate and 25% from fat, whereas the HP-ER diet was 25% of total energy from protein, 50% from carbohydrate and 25% from fat. Both HP diets contained 1.4 g protein/kg per d, with 15 and 25% of total protein provided from egg and pork products, respectively. The additional 0.6 g protein/kg per d was provided as animal protein from eggs and pork and was consumed at breakfast, lunch, dinner or equally divided among all meals (0.2 g protein/meal). For the breakfast, lunch and dinner trials, the meals not containing additional protein were identical to the NP meals. Additional meal nutrient content information is provided in Table 1 and a sample menu is provided in Appendix 1.

Participants reported to the metabolic kitchen for all meals on 6 d consecutively for 5 weeks. Breakfast, lunch and dinner were served between 07.00 and 09.00, 11.00 and 13.00, and 16.00 and 18.00 hours, respectively. Participants ate under supervision of research staff and consumed all foods provided to them. During each testing day, the subjects were instructed to eat only the foods provided during the supervised breakfast, lunch and dinner meals at the metabolic kitchen. While the subjects were allowed to leave the kitchen after each meal was consumed, they were not permitted to eat or drink anything besides water during the time between each meal. If a subject consumed anything during this time, the quantity and type of food or drink was recorded. None of the volunteers reported eating or drinking anything extra.

Appetite

Fullness, hunger and desire to eat were assessed over 15 h on days 3 and 6 by responses to questions such as 'How strong is your feeling of fullness?' during the following times: pre-meal (–30 min), +30, +60, +90, +120, +180 min post-meal for breakfast, lunch and dinner. Responses were recorded on a thirteen-point, numbered, linear category scale. The end anchors were 'not at all' to 'extremely'. Participants were instructed to circle the vertical dash along the horizontal line corresponding to their feelings at that moment, and the results are reported using arbitrary units (au).

Data and statistical analyses

To assess the acute appetite responses, the meal-related 3 h postprandial fullness, hunger and desire to eat area under the curves were calculated for the HP meal in each HP treatment, each NP meal in the NP treatment, and each HP meal in the HP-E treatment. To assess the sustained appetite response, the 15 h (overall) area under the curves were determined for each treatment. All area under the curve measurements were calculated using the trapezoidal rule⁽¹⁰⁾. The NP treatment was incorporated into the study design to provide the reference (i.e. 'normal') responses for the meal-related and overall feelings of fullness, hunger and desire to eat. All data throughout each of the HP treatments are therefore expressed as the difference (Δ) from NP.

Table 1. Dietary characteristics of the normal protein (NP) v. higher protein breakfast (HP-B), lunch (HP-L), dinner (HP-D) and equally distributed (HP-E) energy balance (EB) and energy restriction (ER) diets in nine volunteers*

(Mean values with their standard errors)

Dietary characteristics	NP meals				HP-B, HP-L, HP-D†				HP-E‡			
	EB		ER		EB		ER		EB		ER	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
Meal (kJ/meal)	4280 ^a	109	3314 ^c	109	4431 ^b	117	3452 ^c	117	4301 ^a	113	3494 ^c	126
Protein (g/meal)	28.8 ^a	0.8	28.0 ^a	0.8	92.2 ^b	2.8	92.0 ^b	2.7	49.8 ^c	1.3	50.0 ^c	1.5
Carbohydrate (g/meal)	168.2 ^a	4.4	124.5 ^d	4.1	105.3 ^b	4.0	61.4 ^e	3.5	147.2 ^c	3.9	110.8 ^f	4.9
Fat (g/meal)	29.1 ^a	0.7	22.4 ^b	0.8	29.8 ^a	0.8	23.2 ^b	1.0	29.4 ^a	0.8	23.1 ^b	0.8

^{a–f} Mean values within a row with unlike superscript letters were significantly different (repeated measures ANOVA within and between treatments; $P < 0.05$).

* For details of subjects and procedures, see Materials and methods.

† Dietary characteristics only correspond to the HP meal. During each of these trials, the non-HP meals were the same as the NP meals.

‡ Dietary characteristics were consumed at each of the three meals of a given day of the trial.

To examine the main effect of protein timing, a repeated measures ANOVA was performed on the meal-related and overall appetite responses during EB and ER. When main effects were detected, *post hoc* analyses were performed using Least Significant Difference procedures. This statistical approach led to greater than 80% power (0.87) to detect differences in meal-related and overall appetite among treatments. All measurements are expressed as means and their standard errors. An α level of $P < 0.05$, two-tailed, was considered statistically significant. Statistical analyses were performed using the Statistical Package for the Social Sciences version 15.0 (SPSS Inc., Chicago, IL, USA).

Results

Meal-related (3 h postprandial) fullness

While no difference in meal-related fullness was observed among the HP treatments during EB (data not shown), the HP-B during negative EB led to greater meal-related fullness (+137 (SEM 44) au \times 180 min) *v.* HP-D (-1 (SEM 37) au \times 180 min; $P = 0.003$). The response to the HP-B did not differ significantly from the HP-L (+62 (SEM 53) au \times 180 min; $P = 0.188$) or HP-E-B (+92 (SEM 85) au \times 180 min; $P = 0.587$).

Overall (15 h composite) fullness

No difference in overall fullness was observed among the HP treatments during EB (data not shown). During ER, the HP-B elicited greater fullness throughout the 15 h (+404 (SEM 162) au \times 900 min) compared to HP-L (+33 (SEM 162) au \times 900 min; $P = 0.009$) and HP-D (-60 (SEM 132) au \times 900 min; $P = 0.05$) (Fig. 1). The response to the HP-B did not differ significantly from the HP-E (+274 (SEM 165) au \times 900 min; $P = 0.188$) (Fig. 1).

Meal-related and overall hunger and desire to eat

No differences in the meal-related and overall hunger and desire to eat responses were observed during EB. During ER, the meal-related and overall hunger and desire to eat responses tended to be comparable to the fullness responses but were not as statistically strong (data not shown).

Discussion

While the satiating property of increased dietary protein has been independently observed during breakfast^(7,8) and lunch^(7,11), the relative efficacy of consuming protein in different patterns is not known. We found that during ER, increased dietary protein at breakfast led to greater initial and/or sustained increases in fullness compared to lunch and dinner.

Hunger and desire to eat were measured along with the feelings of fullness. We chose to specifically report the fullness (satiety) response due to the well-established satiating properties of protein⁽⁷⁾ and the strong association with food intake which is not consistently observed with the hunger and/or desire to eat responses^(12,13).

The differences in fullness observed between breakfast and the other meals may be a reflection of the greater change in macronutrients from the subject's habitual breakfast consumption. Since most Americans typically consume a relatively small amount of dietary protein at breakfast (15% of total daily protein) in comparison to other meal times⁽¹⁴⁾, the exposure to a disproportionately higher protein test-meal breakfast could potentially lead to a greater feeling of fullness than at other meal times. This is supported by Long *et al.*⁽¹⁵⁾ who examined whether the appetite responses to a higher protein test meal were influenced by habitual protein intake. The subjects habitually consuming a lower protein diet experienced greater




Fig. 1. Fullness responses across the day during the higher protein (HP)-breakfast (B), lunch (L), dinner (D) or equally divided (E) treatments during energy restriction. (A), Fullness responses at each time-point (○, HP-B; ●, HP-L; △, HP-D; ▲, HP-E; ■, normal protein (NP)). Values are means. au, arbitrary units. (B), Fullness area under the curve over the entire 15 h period. Values are means with their standard errors depicted by vertical bars. ^{a,b} Mean values with unlike superscript letters were significantly different (repeated measures ANOVA between treatments; $P < 0.05$).

postprandial satiety following the higher protein test meal compared to those who habitually consumed a higher protein diet. This would suggest that our differential satiety response at breakfast may have been simply due to the change in diet and not the timing of protein consumption. We previously evaluated whether protein habituation during weight loss influenced the acute, meal-related appetitive responses⁽⁸⁾. Unlike that of Long *et al.*⁽¹⁵⁾, the appetite responses following the higher protein meal *v.* the normal protein meal in the present study were unaffected by chronic protein intake⁽⁸⁾.

Along these lines, our experimental design incorporated three meals of equivalent energy intake. Since breakfast is typically much smaller than the lunch or dinner meals⁽⁶⁾, the larger breakfast in the current study may have influenced the raw fullness responses. We chose to compare each treatment according to the difference from the NP treatment, which also incorporated a larger breakfast, to counter any differences. We believe the satiating effects of protein at breakfast in the current study are not a result of the change in habitual diet.

The effects of increased protein when equally divided among all meals were also examined. We anticipated greater overall fullness during this treatment as a small amount of protein at every eating occasion should theoretically sustain feelings of fullness throughout the day. No difference in fullness was observed between this treatment and any of the other higher protein treatments. Further, this treatment led to similar fullness compared to the normal protein treatment. This may be explained by the fact that most increased protein studies incorporate protein amounts on the order of approximately 75 g protein/meal or greater⁽⁷⁾. Our HP-E treatment contained approximately 50 g/meal, suggesting a threshold or graded effect may exist for the amount of protein consumed at one time to yield differences in appetite. Additional studies are needed to characterize the dose–response relationship.

The present study also indicates that the satiating properties of protein are affected by energy status as no difference in fullness was observed between the normal protein and increased dietary protein during EB. In a comprehensive review by Halton & Hu⁽⁷⁾, eleven of fourteen studies found greater satiety with increased dietary protein. This finding is also supported by two recent studies^(16,17). Several main differences exist when comparing these studies to our current study: (1) the dietary protein in these studies ranged from 30 to 100% of total energy intake^(7,16,17), whereas the current study was only 18% in the EB segment; (2) many of these studies incorporated healthy men and/or women, while the current study targeted overweight and/or obese men; (3) lastly, none of these studies compared the increased dietary protein among all three meals^(7,16,17). The present data imply that the satiating effect of increased dietary protein is blunted when consuming 18% of total energy intake as protein in an EB state while other data support that consuming larger amounts of dietary protein will lead to greater satiety during EB.

We have suggested an increased sensitivity to dietary protein during breakfast as compared to other meal times. Due to the fact that the satiating properties of dietary protein consumed at dinner were completely abolished (as evidenced by the similar postprandial fullness response to that of the normal protein treatment), it may, in actuality, be the reduced sensitivity to dietary protein during lunch and dinner. This would support the concept that individuals appear to be less

satiated towards evening leading to increased meal size and shorter inter-meal intervals throughout the day⁽²⁾.

The present data lend further support for the need to incorporate increased dietary protein at breakfast when designing effective energy-restricted diets. Several study limitations exist. First, the findings are derived from acute subjective measures of fullness (satiety) from questionnaires. While the data indicate differential fullness responses among the treatments, the present study design does not allow for the examination of whether these fullness responses would lead to differences in food intake. According to Parker *et al.*⁽¹²⁾ and Flint *et al.*⁽¹⁸⁾, subjective sensations assessing fullness from questionnaires are highly predictive and/or correlated with food intake. Thus, we are confident that the present data provide important and unique findings concerning the timing of protein consumption in relation to appetite and food intake.

The hormones involved with the regulation of food intake and body weight were not examined. Further research is necessary to confirm the present findings, document changes in food intake, and to explore the underlying mechanisms and long-term implications for EB and body weight control.

Lastly, the present study only examined the effect of increased animal protein (provided from the pork and eggs). While research suggests comparable fullness (satiety) between meals containing animal-based (egg albumin) protein *v.* plant-based protein (i.e. soya)⁽¹⁹⁾, we did not test this in the present study. Thus, caution should be made concerning the generalizability of the present data.

Conclusions

The differential appetitive responses following increased protein intake at breakfast compared to other meal times suggest that the satiating property of dietary protein is influenced by the timing of protein consumption.

Acknowledgements

The authors would like to thank Nadine S. Carnell, MS, who provided nutritional support concerning the test meals. This study was funded through the National Pork Board and the American Egg Board – Egg Nutrition Center. Support for H. L. was also provided, in part, by an Ingestive Behavior Research Center post-doctoral fellowship from Purdue University. None of the authors had any personal or financial conflicts of interest.

References

1. Keski-Rahkonen A, Kaprio J, Rissanen A, Virkkunen M & Rose RJ (2003) Breakfast skipping and health-compromising behaviors in adolescents and adults. *Eur J Clin Nutr* **57**, 842–853.
2. de Castro JM (2004) The time of day of food intake influences overall intake in humans. *J Nutr* **134**, 104–111.
3. van der Heijden AAWA, Hu FB, Rim EB & van Dam RM (2007) A prospective study of breakfast consumption and weight gain among U.S. men. *Obesity* **15**, 2463–2469.
4. Purslow LR, Sandhu MS, Forouhi N, Young EH, Luben RN, Welch AA, Khaw K-T, Bingham SA & Wareham NJ (2008) Energy intake at breakfast and weight change: prospective

- study of 6764 middle-aged men and women. *Am J Epidemiol* **167**, 188–192.
5. Song WO, Chun OK, Obayashi S, Cho S & Chung CE (2005) Is consumption of breakfast associated with body mass index in US adults? *J Am Diet Assoc* **105**, 1373–1382.
 6. de Castro JM (2008) The time of day and the proportions of macronutrients eaten are related to total daily food intake. *Br J Nutr* **98**, 1077–1083.
 7. Halton TL & Hu FB (2004) The effects of high protein diets on thermogenesis, satiety and weight loss: a critical review. *J Am Coll Nutr* **23**, 373–385.
 8. Leidy HJ, Mattes RD & Campbell WW (2007) Effects of acute and chronic protein intake on metabolism, appetite, and ghrelin during weight loss. *Obesity* **15**, 1215–1225.
 9. Harris JA & Benedict FG (1919) *A Biometric Study of Basal Metabolism in Man*. Washington, DC: Carnegie Institute of Washington.
 10. Wolever TM, Jenkins DJ, Jenkins AL & Josse RG (1991) The glycemic index: methodology and clinical implications. *Am J Clin Nutr* **54**, 846–854.
 11. Latner JD & Schwartz M (1999) The effects of a high-carbohydrate, high-protein or balanced lunch upon later food intake and hunger ratings. *Appetite* **33**, 119–128.
 12. Parker BA, Sturm K, MacIntosh CG, Feinle C, Horowitz M & Chapman IM (2004) Relation between food intake and visual analogue scale ratings of appetite and other sensations in healthy older and young subjects. *Eur J Clin Nutr* **58**, 212–218.
 13. Raben A, Tagliabue A & Astrup A (1995) The reproducibility of subjective appetite scores. *Br J Nutr* **73**, 517–530.
 14. Moshfegh A, Goldman J & Cleveland L (2005) *What We Eat in America, NHANES 2001–2002: Usual Nutrient Intakes from Food Compared to Dietary Reference Intakes*. Washington, DC: US Department of Agriculture, Agricultural Research Service.
 15. Long SJ, Jeffcoat AR & Millward DJ (2000) Effect of habitual dietary-protein intake on appetite and satiety. *Appetite* **35**, 79–88.
 16. Moran LJ, Luscombe-Marsh ND, Noakes M, Wittert GA, Keogh JB & Clifton PM (2005) The satiating effect of dietary protein is unrelated to postprandial ghrelin secretion. *J Clin Endocrinol Metab* **90**, 5205–5211.
 17. Lejeune MPGM, Westerterp KR, Adam TCM, Luscombe-Marsh ND & Westerterp-Plantenga MS (2006) Ghrelin and glucagon-like peptide 1 concentrations, 24-h satiety, and energy and substrate metabolism during a high-protein diet and measured in a respiration chamber. *Am J Clin Nutr* **83**, 89–94.
 18. Flint A, Raben A, Blundell JE & Astrup A (2000) Reproducibility, power and validity of visual analogue scales in assessment of appetite sensations in single test meal studies. *Int J Obes Relat Metab Disord* **24**, 38–48.
 19. Lang V, Bellisle F, Oppert JM, Craplet C, Bornet FR, Slama G & Guy-Grand B (1998) Satiating effect of proteins in healthy subjects: a comparison of egg albumin, casein, gelatin, soy protein, pea protein, and wheat gluten. *Am J Clin Nutr* **67**, 1197–1204.

Appendix 1. Sample subject energy restriction menus

Normal protein (NP)	Higher protein breakfast (HP-B)	Higher protein lunch (HP-L)	Higher protein dinner (HP-D)	Higher protein equally divided (HP-E)
Breakfast	Breakfast	Breakfast	Breakfast	Breakfast
Entre: 5 Eggo waffles 85 g pancake syrup 14 g margarine 5 slices meatless bacon Sides: 367 g FF milk	Sandwich: 2 slices whole wheat bread 9 slices Canadian bacon 126 g egg substitute Sides: 1 large hard-boiled egg 126 g cranberry juice 245 g FF milk 59 g banana 226 g low-carbohydrate yogurt	Entre: 5 Eggo waffles 57 g pancake syrup 14 g margarine 5 slices meatless bacon Sides: 124 g orange juice 306 g FF milk	Entre: 5 Eggo waffles 85 g pancake syrup 14 g margarine 5 slices meatless bacon Sides: 367 g FF milk	Sandwich: 2 English muffins 3 slices Canadian bacon 63 g egg substitute 2 meatless sausage patties 1 slice FF American cheese Sides: 173 g melon 186 g apple juice 15 grapes
Lunch	Lunch	Lunch	Lunch	Lunch
Entre: 42 g meatless beef 110 g lettuce 28 g blue cheese 28 g Italian dressing Sides: 2 slices sourdough bread 92 g broccoli 124 g pears with 38 g low-fat whipped topping 214 g fruit cocktail 248 g soda	Sandwich: 2 slices whole wheat bread 3 slices of meatless bacon 20 g lettuce 8 g FF mayonaise 2 slices FF American cheese 2 slices tomatoes 1 dill pickle Sides: 25 potato chips 132 g pie filling 26 g low-fat whipped topping 1 frozen ice cream sandwich	Sandwich: 2 slices whole wheat bread 198 g extra lean ham 168 g egg substitute 20 g lettuce 8 g FF mayonaise 4 slices FF American cheese 60 g tomatoes Sides: 10 potato chips 245 g FF milk	Sandwich: 2 slices whole wheat bread 4 slices meatless bacon 20 g lettuce 5 g FF mayonaise 1 slice FF American cheese 60 g tomatoes 1 dill pickle Sides: 25 potato chips 132 g pie filling 10 g low fat whipped topping 1 frozen ice cream sandwich 184 g FF milk	Salad: 71 g extra lean ham 63 g egg substitute 56 g meatless beef 35 g mushrooms 68 g green beans 3 dried prunes 30 g Italian dressing 110 g lettuce 82 g corn Sides: 1 dinner roll 5 g margarine 2 chocolate éclairs
Dinner	Dinner	Dinner	Dinner	Dinner
Entre: 4 wheat tortillas 25 g peppers 74 g carrots 105 g onions 5 g olive oil 28 g cheddar cheese 32 g sour cream 62 g salsa 120 g red beans Sides: 96 g sherbet	Entre: 4 meatless meatballs 158 g mashed potatoes 105 g green beans 112 g carrots 55 g lettuce 40 g tomatoes 3 g olive oil 3 g vinegar Sides: 2 dinner rolls 143 g fruit cocktail 19 g low-fat whipped topping 2 low-fat devil's food cookies	Entre: 210 g spaghetti noodles 21 g meat crumbles 125 g spaghetti sauce 5 g olive oil Sides: 55 g lettuce 40 g tomatoes 28 g FF mozzarella cheese 28 g Italian dressing 1 slice sourdough bread 14 g margarine 62 g peaches 9 g low-fat whipped topping 1 piece of angel food cake	Entre: 2 wheat tortillas 170 g pork loin 168 g egg substitute 3.5 slices American cheese 32 g sour cream Sides: 245 g FF milk	Entre: 57 g pork loin 63 g egg substitute 42 g stuffing Sides: 1 dinner roll 14 g margarine 1 snack-size chocolate pudding 1 low-fat devil's food cookie 490 g FF milk

FF, fat-free.

Protein at breakfast increases fullness