

The beneficial effect of the sap of *Acer mono* in an animal with low-calcium diet-induced osteoporosis-like symptoms

Geun-Shik Lee¹, Hyuk-Soo Byun¹, Man-Hee Kim¹, Bo-Mi Lee¹, Sang-Hwan Ko¹, Eui-Man Jung¹, Ki-Seob Gwak², In-Gyu Choi², Ha-Young Kang³, Hyun-Jin Jo³, Hak-Ju Lee³ and Eui-Bae Jeung^{1*}

¹Laboratory of Veterinary Biochemistry and Molecular Biology, College of Veterinary Medicine, Chungbuk National University, Cheongju, Chungbuk 361-763, Republic of Korea

²Department of Forest Sciences, College of Agriculture & Life Sciences, Seoul National University, Seoul 151-921, Republic of Korea

³Division of Wood Chemistry and Microbiology, Department of Forest Products, Korea Forest Research Institute, Seoul 130-712, Republic of Korea

(Received 18 September 2007 – Revised 18 January 2008 – Accepted 12 February 2008 – First published online 1 April 2008)

The sap of *Acer mono* has been called ‘bone-benefit-water’ in Korea because of its mineral and sugar content. In particular, the calcium concentration of the sap of *A. mono* is 37.5 times higher than commercial spring water. In the current study, we examined whether *A. mono* sap could improve or prevent osteoporosis-like symptoms in a mouse model. Male mice (3 weeks old) were fed a low-calcium diet supplemented with 25, 50 or 100% *A. mono* sap, commercial spring water or a high calcium-containing solution as a beverage for 7 weeks. There were no differences in weekly weight gain and food intake among all the groups. Mice that were given a low-calcium diet supplemented with commercial spring water developed osteoporosis-like symptoms. To assess the effect of sap on osteoporosis-like symptoms, we examined serum calcium concentration, and femur density and length, and carried out a histological examination. Serum calcium levels were significantly lower in mice that received a low-calcium diet supplemented with commercial spring water (the negative control group), and in the 25% sap group compared to mice fed a normal diet, but were normal in the 50 and 100% sap and high-calcium solution groups. Femur density and length were significantly reduced in the negative control and 25% sap groups. These results indicate that a 50% sap solution can mitigate osteoporosis-like symptoms induced by a low-calcium diet. We also examined the regulation of expression of calcium-processing genes in the duodenum and kidney. Duodenal *TRPV6* and renal *calbindin-D9k* were up-regulated dose-dependently by sap, and the levels of these factors were higher than those attained in the spring water-treated control. The results demonstrate that the sap of *A. mono* ameliorates the low bone density induced by a low-calcium diet, most likely by increasing calcium ion absorption.

Acer mono: Calcium-processing genes: Osteoporosis: Sap

Sap is the fluid transported in the xylem cells or phloem sieve tube elements of a plant. Sap consists primarily of water, but also contains mineral elements, sugars, hormones and other nutrients. Other liquid components found in plants or that are exuded by plants, such as latex, resins and mucilage, are sometimes incorrectly referred to as sap⁽¹⁾. *Acer mono* is a rounded, deciduous tree with five to seven bright green, almost heart-shaped, lobular leaves that turn yellow in autumn. Yellow flowers appear on erect umbels with these leaves, but may also appear without them. The sap of *A. mono* contains sugar, and can be ingested directly as a drink, or concentrated into a syrup by boiling⁽²⁾ and then used as a sweetener. In Korea, the sap of *A. mono* has been called ‘bone-benefit-water’ because of its abundant calcium and magnesium ion content. The medicinal uses of *A. mono* include the leaves as an irritant and the bark as an

astringent⁽³⁾, however, the pharmacological effect of the sap on bone health has not been examined.

The most common abnormality of bone is osteoporosis, which is a decrease in the density of the bone mass that causes thinning and weakening of the bone. There are two main categories of osteoporosis: type I and type II. Type I osteoporosis occurs only in post-menopausal women, and is due to oestrogen deficiency. Type II osteoporosis occurs in both men and women (about two times more frequently in women), and is due to ageing and calcium deficiency. In animal models, a low-calcium diet leads to the development of radiologic rickets and histologic features of osteomalacia⁽⁴⁾. These symptoms were defined as osteoporosis-like symptoms in the present study.

To clarify the potential therapeutic nature of ‘bone-benefit-water’, an animal model of low-calcium diet-induced

Abbreviations: CaBP-9k, calbindin-D9k; 1,25-(OH)₂D₃, 1,25-dihydroxyvitamin D₃; PTH, parathyroid hormone; TRPV5, transient receptor potential vanilloid 5; TRPV6, transient receptor potential vanilloid 6.

* **Corresponding author:** Dr Eui-Bae Jeung, fax +82 43 267 3150, email ebyeung@chungbuk.ac.kr

osteoporosis-like symptoms has been generated. Using this model, the effect of sap on the active calcium transport system has also been examined. Calcium ions are actively absorbed from the duodenum and reabsorbed in kidney. In the calcium active transport system, several proteins mediate calcium influx, transfer through the cytosol and extrusion into the bloodstream. In the active calcium-transporting organs, these steps are carried out by calcium entry channel proteins of the outer cell membrane, cytosolic buffering or transfer proteins, and excretory pump proteins^(5–7). Two highly selective calcium channels located in the apical cell membrane are the main portals of calcium ion entry, transient receptor potential vanilloid 6 (TRPV6) and 5 (TRPV5). Calbindin-D9k (CaBP-9k) is an intracellular calcium ion-binding protein that is thought to participate in shuttling calcium ions from the apical to the basolateral membrane, where calcium ions are extruded. However, the proteins involved in intracellular calcium transport vary somewhat depending on the tissue. In the epithelial cells of the duodenum, the transport pathway involves calcium entry primarily through specialized TRPV6 channels, and is highly dependent on intracellular free-calcium concentration. Maintenance of the 'open' state of TRPV6 requires the constant buffering action of intracellular calcium-binding proteins. In mammalian intestinal cells, CaBP-9k is the major intracellular calcium-binding protein. The final step of extrusion of calcium into the blood requires ATP. In the kidney, trans-cellular calcium ion transport is carried out by cells lining the distal part of the nephron. Calcium ion entry is facilitated by TRPV5 in the apical membrane. Calcium then binds CaBP-9k and diffuses through the cytosol to the basolateral membrane, where it is extruded, directly or via CaBP-9k, by excretory pump proteins⁽⁶⁾.

In the current study, we examined the effect of the sap of *A. mono* on osteoporosis-like symptoms. Mice were fed a low-calcium diet to induce osteoporosis-like symptoms during their growth stage (from 3 to 10 weeks of age). Solutions of increasing percentages of sap, commercial spring water or a high-calcium-content solution were supplied to the mice, and after 7 weeks, bone mineral density and length, and the osteoid thickness of the femur and tibia were analysed. We also examined the expression of duodenal and renal calcium-processing genes. Serum calcium and parathyroid hormone (PTH) levels were examined as markers of the development of osteoporosis-like symptoms in the mice.

Materials and methods

Mineral concentration of the sap of *Acer mono*

The mineral contents of *A. mono* sap and commercial spring water (Samdasu, Jeju Special Self-governing Province Development Corp., Jeju-si, Jeju-do, Korea) were analysed using an inductively coupled plasma emission spectrometer (ICPS-1000 IV; Shimadzu, Tokyo, Japan). All samples (10 ml) were digested with the addition of 20 ml nitric acid to remove all organic matter, reduced in volume with heating, filtered through no. 42 filter paper (Whatman, Maidstone, UK), and then diluted to a final volume of 50 ml before being processed by an inductively coupled plasma emission spectrometer.

Animals and treatments

Male ICR mice (3 weeks old) were obtained from KOATECH (Pyeongtaek-si, Gyeonggi-do, Korea). All animals were housed in polycarbonate cages and acclimatized to an environmentally controlled room ($23 \pm 2^\circ\text{C}$, relative humidity $50 \pm 10\%$, frequent ventilation and 12 h light/dark cycle) prior to use. Thirty mice were divided into six groups (five in each group). Osteoporosis-like symptoms were induced in five groups by feeding with a low-calcium diet (AIN-76A purified rodent diet with 1% phosphorus and 0.02% calcium, DYET 113294; Dyets Inc., Bethlehem, PA, USA) for 7 weeks. One group was fed a normal diet (AIN-76A purified rodent diet with 0.8% phosphorus and 1.1% calcium, DYET #113295; Dyet Inc.; normal group) for the same period of time.

To assess the preventative effect of *A. mono* sap on osteoporosis-like symptoms, a daily total of 30 ml commercial spring water (negative control group), or a solution of 25, 50 or 100% sap, or a high-calcium-content rescue solution (7.5 mM-CaCl₂ and 3.0 mM-NaH₂PO₄ based on the Krebs–Ringer buffer, rescue group) were restrictively supplied to the mice from 3 to 10 weeks of age. Body weights were measured weekly for the duration of the experiment. The Ethics Committee of the Chungbuk National University approved all experimental procedures and use of animals.

Bone mineral density and histomorphometry

Mice were killed at 10 weeks of age by CO₂ inhalation. The hind legs were removed and fixed in a 10% neutralized formalin solution for 3 d. The legs were then washed with water, and the skin and muscle were eliminated. The femur was dried overnight and bone mineral density was analysed using a Direct View CR 500 (Eastman Kodak Co., Rochester, NY, USA). The tibia was decalcified using Plank-Rycholo's solution, and the bone was embedded in paraffin, sectioned and deparaffinized in xylene. Sections were hydrated in a series of descending ethanol concentrations, then stained with haematoxylin and subjected to histomorphometric analyses using light microscopy. Osteoid thickness was measured at the centre point of the tibia.

Real-time PCR using TaqMan™ Probe

Total RNA was prepared from the duodenum and kidney using the TRIzol reagent (Invitrogen, Carlsbad, CA, USA). First stand complementary DNA (cDNA) was prepared by subjecting total RNA (1 µg) to reverse transcription using mMLV RT (Invitrogen Life Technologies Inc.) and random primers (9 mer; TaKaRa Bio. Inc., Otsu, Shiga, Japan)^(8,9). Real-time PCR was performed in a 20 µl reaction volume containing 10 µl TaqMan Universal PCR Master Mix (Applied Biosystems, Foster, CA, USA), 1 µl 20 × Assays-on-Demand™ Gene Expression Assay Mix (Applied Biosystems; *TRPV6*, Mm00499069_m1, NM_022413-2; *TRPV5*, Mm01166029_m1, NM_001007572-2; *CaBP-9k*, Mm00486654_m1, NM_009789-2; *HPRT1*, Mm00446968_m1, NM_013556-2) and 3 µl cDNA. Amplification was carried out using a 7300 Real-Time PCR System (Applied Biosystems) and the following cycle parameters: initial denaturation at 50°C for 2 min; 90°C for 10 min; forty cycles of denaturation at 95°C for 15 s; annealing and extension at 60°C

for 1 min. Expression levels were determined using RQ software (Applied Biosystems). The expression of *TRPV6*, *TRPV5* and *CaBP-9k* was normalized to that of *HPRT1*.

Biochemical and hormonal analysis

Blood was collected from each mouse into serum separator tubes and subjected to centrifugation at 3000 rpm for 10 min. Serum calcium levels were determined using ASAN Ca-Lq Reagents (Asanpharm, Kyeongdo, Korea). Serum PTH levels were measured using a Mouse Intact PTH ELISA Kit (ALPCO Diagnostics, Salem, NH, USA), according to manufacturer's instructions.

Data analysis

Data were analysed by non-parametric one-way ANOVA using the Kruskal–Wallis test, followed by Dunnett's test for multiple comparisons to the negative control group. Data were ranked according to these tests. Statistical analysis was performed using SPSS for Windows (SPSS Inc., Chicago, IL, USA). $P < 0.05$ was considered statistically significant.

Results

The mineral content of *Acer mono* sap

The concentration of sixteen mineral ions was measured in the sap of *A. mono* and commercial spring water. The potassium, calcium and magnesium levels of sap were 16, 37 and 3.9 times higher than commercial spring water, respectively, while the level of sodium in sap was 0.68 times lower (Table 1). A small amount of silicon was detected in both solutions. Manganese was detected only in the sap, and iron was detected only in commercial spring water. No other minerals were detected in either liquid.

Table 1. Mineral content (mg/l) of the sap of *Acer mono* and commercial spring water*

(Mean values with their standard errors)

	Commercial spring water		Sap of <i>A. mono</i>		Rescue solution
	Mean	SEM	Mean	SEM	
K	7.34	0.05	117.96	2.86	
Ca	4.15	0.27	155.32	3.94	300.6
Na	27.86	0.97	18.90	0.49	68.97
Mg	3.68	0.12	14.39	0.44	
P	ND		ND		92.92
Si	0.08	0.00	0.07	0.01	
Al	ND		ND		
Mn	ND		0.35	0.09	
Fe	0.31	0.08	0.10		
Cu	ND		ND		
Zn	ND		ND		
As	ND		ND		
Mo	ND		ND		
Cd	ND		ND		
Pb	ND		ND		
Cr	ND		ND		

ND, not detected.

* For details of procedures, see Materials and methods.

The effect of *Acer mono* sap on osteoporosis-like symptoms

To elucidate the beneficial effect of sap on osteoporosis-like symptoms, mice from the weaning (3 weeks old) to mature (10 weeks old) stage were fed low-calcium diets supplemented with increasing percentage solutions (0, 25, 50 and 100 %) of sap, or a rescue solution containing a high concentration of calcium ions, and femur mineral densities were analysed using computer radiography (Fig. 1). The femurs of the normal group were bright and clear in radiographic images, while the corresponding images of the negative control group, which received commercial spring water, were dark and indistinct, which indicated that the mice had developed osteoporosis-like symptoms (Fig. 1 (A)). The rescue group (which received high calcium-containing water) exhibited improved femur brightness in radiographs. To determine the effect of sap on osteoporosis-like symptoms more precisely, the density of the distal (D) and proximal (P) femur heads was analysed using Direct View CR 500 software (Fig. 1 (B)). The densities of the distal and proximal femur heads of the normal group were significantly higher than all of the low-calcium diet groups. However, the bone densities of the 50 and 100 % sap solution groups and the rescue group were significantly higher than the 25 % sap solution and negative control group. Femur length was also analysed (Fig. 2 (C)), and we found that mice treated with commercial spring water alone, or a 25 % sap solution, had significantly shorter femurs than normal mice. The femur lengths of the groups that received 50 % or higher sap solutions were not significantly different from the normal group. The effect of sap on low-calcium diet-induced osteoporosis-like symptoms was further analysed by bone histomorphometry. The tibia was sectioned and stained with haematoxylin, and osteoid thickness was analysed (Fig. 2 (A)). The thickness of the negative group was significantly less than the normal group, and thickness gradually increased with increasing percentage sap solutions (Fig. 2 (B)). The tibia thicknesses of all of the low-calcium diet groups were significantly less than the normal group. Supplementation with 50 % or greater sap solutions or rescue solution improved tibia thickness compared to the negative control group. The present results suggested that 50 % or higher percentage sap solutions improve low-calcium-induced osteoporosis-like symptoms.

The effect of sap on calcium-processing gene expression

Based on the previous experiment, administration of a 50 % sap solution appeared to mitigate low-calcium diet-induced osteoporosis-like symptoms in mice. To begin to uncover the mechanism of the beneficial effect of sap on osteoporosis-like symptoms, we examined the expression of three duodenal and renal calcium-absorbing genes using real-time PCR. The expression of duodenal *CaBP-9k* was similar in mice that received sap, commercial spring water or a normal diet (Fig. 3 (A)). Mice that received the high calcium-containing rescue solution had significantly reduced expression levels of duodenal *CaBP-9k* compared with negative and normal dieted groups. The expression of duodenal *TRPV6* was increased in the sap solution groups, but was decreased in the rescue group when compared with negative and normal dieted mice. In the kidney, the *CaBP-9k* expression levels of the




Fig. 1. The effect of *Acer mono* sap on osteoporosis-like symptoms assessed using bone radiography. Femurs were obtained from mice and analysed as described in the Materials and methods. (A), Representative femurs of each of the indicated treatment groups. (B, C) Bone density at the distal (B) and proximal (C) heads of all femurs (five per group). Values are means of duplicate measurements of all samples with their standard errors depicted by vertical bars. Data are expressed as a percentage of the density of a normal femur. D, distal head of femur; P, proximal head of femur; Negative, low-calcium diet plus commercial spring water; Normal, normal diet plus commercial spring water; Rescue, low-calcium diets plus high-calcium solution. Mean values were significantly different from those of the normal group: * $P < 0.05$. Mean values were significantly different from those of the negative control group: † $P < 0.05$.

sap solution and control mice were higher than those of the mice in the negative and rescue groups, and *CaBP-9k* expression was down-regulated in the negative and rescue groups compared to the normal group (Fig. 3 (B)). The expression of renal *TRPV6* was not altered in any of the low-calcium diet groups. However, renal *TRPV5* expression was decreased in the negative, sap solution and rescue groups compared to the normal group. These distinct patterns of regulation of duodenal and renal calcium-processing gene expression suggested that the mechanism that improves osteoporosis-like symptoms in the sap solution group involves not only augmentation of calcium levels, but also enhanced activity of the calcium active transport system.

The effect of the sap on serum calcium and parathyroid hormone levels

Serum calcium concentration was used to monitor changes in calcium homeostasis. The serum calcium concentrations of the negative group and the 25% sap solution group were lower than the normal group, while in the 50% or higher sap solution and rescue groups, serum calcium concentration was similar to normal levels (Table 2). The present result was consistent with the bone data given earlier. Serum PTH

levels are also an indicator of osteoporosis^(4,9), as high PTH levels indicate that calcium ions are being released from the bone into serum to maintain calcium homeostasis. The negative and sap solution groups exhibited elevated serum PTH levels, while the rescue group had a significantly lower PTH level compared to the normal group.

Discussion

Maple syrup is a sweetener made from the sap of maple trees. It is often eaten with pancakes, waffles or French toast, and is used as an ingredient in baking and the preparation of desserts. *A. mono* is a kind of maple tree found in eastern Asia, and its sap can be ingested directly, without any processing. The sap of *A. mono* consists primarily of water, and contains minerals and sugars, similar to the sap of maple⁽¹⁾. In the current study, we examined the pharmacological effect of the sap of *A. mono*. This sap is believed to be beneficial to bone health because of its high mineral content. We first examined the mineral content of sap, and compared it to commercial spring water. We showed that potassium, calcium and magnesium were present at high levels in sap. The level of calcium ions was 37 times higher, and magnesium was 3.9 times higher in the sap of *A. mono* than in commercial spring water. We hypothesized




Fig. 2. The effect of *Acer mono* sap on osteoporosis-like symptoms assessed using bone histomorphometry. Tibias were removed from mice and processed as described in the Materials and methods. (A), Representative tibias of the indicated treatment groups. (B) Osteoid thickness taken at the middle of the bone for all femurs (five per group). \leftrightarrow , Measured thickness; space between the arrows represents the bone marrow. Values are means of duplicate measurements of all samples with their standard errors depicted by vertical bars. Data are expressed as a percentage the tibia thickness of the normal group. (C), Femur length analysed by radiography. Values are means of duplicate measurements of all samples with their standard errors depicted by vertical bars. Negative, low-calcium diet plus commercial spring water; Normal, normal diet plus commercial spring water; Rescue, low-calcium diets plus high-calcium solution. Mean values were significantly different from those of the normal group: * $P < 0.05$. Mean values were significantly different from those of the negative control group: † $P < 0.05$.

that it was the high calcium content of sap that prevented or improved the osteoporosis-like symptoms.

It is well known that calcium is essential to attain maximum bone mass^(10,11). Reduced calcium intake results in osteoporosis in man and in many other animal species⁽¹²⁾. Clinical and experimental nutritional secondary hyperparathyroidism rapidly develops in horses, dogs and cats fed diets high in phosphate and low in calcium⁽¹²⁾. Young rats and mice fed diets low in calcium develop osteoporosis-like symptoms, and ageing mice fed a high-calcium diet exhibit accelerated bone resorption with increased levels of dietary phosphate⁽¹²⁾. Similarly, ageing rats fed a diet with adequate calcium content developed progressively more severe osteoporosis as dietary phosphate was elevated⁽¹²⁾. Adult dogs fed diets high in phosphate but adequate levels of calcium developed rapid bone loss resembling osteoporosis, whereas young dogs developed severe nutritional secondary hyperparathyroidism characterized by fibrous osteodystrophy⁽¹²⁾. In the current study, we examined

whether the high calcium content of the sap of *A. mono* would affect bone density when given as a supplement to a low-calcium diet. Although we demonstrated that administration of solutions of increasing percentages of sap increase bone density and osteoid thickness in a mouse model of induced osteoporosis-like symptoms, the sap supplement was less efficient in improving or preventing the symptoms when compared with a high-calcium solution or a normal diet. The present results suggested that in addition to supplying calcium ions, there are other factors involved in the beneficial effect of sap on osteoporosis-like symptoms. The other factor might be potassium which was present at 17-fold higher concentration in sap than in commercial spring water. Potassium influences calcium homeostasis, particularly the urinary conservation and excretion of calcium⁽¹³⁾. Low-potassium diets increase urinary calcium losses while high-potassium diets reduce it⁽¹³⁾. High-potassium intake ameliorates bone resorption in high-salt diets and is associated with a higher baseline bone




Fig. 3. The effect of *Acer mono* sap on the expression of calcium-processing genes. Duodenal (A; transient receptor potential vanilloid 6 (TRPV6), calbindin-D9k (CaBP-9k)) and renal (B; TRPV6, transient receptor potential vanilloid 5 (TRPV5), CaBP-9k) mRNA were prepared from mice of the indicated treatment groups (five per group) as described in the Materials and methods. Relative gene expression of CaBP-9k, TRPV5 and TRPV6 was examined using real-time PCR. Values are means of duplicate measurements of all samples with their standard errors depicted by vertical bars. Data are expressed as a percentage of the expression of *HPRT1*, an internal control gene. Negative, low-calcium diet plus commercial spring water; Normal, normal diet plus commercial spring water; Rescue, low-calcium diet plus high-calcium solution. Mean values were significantly different from those of the normal group: * $P < 0.05$. Mean values were significantly different from those of the negative control group: † $P < 0.05$.

mineral density and lower levels of bone loss^(14,15). Taken together, the present results suggest that the sap of *A. mono* improves low-calcium diet-induced osteoporosis-like symptoms via augmentation of mineral levels.

To clarify the mechanism of the sap's effect on osteoporosis-like symptoms, we examined the expression of calcium active transport genes. Based on the results of previous studies^(9,16–18), we examined the expression of TRPV5, TRPV6 and CaBP-9k. TRPV5 and TRPV6 are calcium channels found in the apical membranes of intestinal and renal epithelial cells, and have been proposed as the mediators of calcium uptake during trans-cellular calcium transport⁽¹⁹⁾. They are highly related proteins expressed mainly in cells involved in calcium absorption or re-absorption in the duodenum and kidney⁽²⁰⁾. TRPV6 is distributed in the duodenum, jejunum, ileum and kidney, and in exocrine tissues such as the pancreas, prostate, mammalian gland and sweat gland^(19,21,22). Hormonal vitamin D-induced expression of TRPV5 occurs in rabbit renal epithelial cells, and TRPV5 has been identified in human, rat and mouse kidney. Dietary calcium also regulates TRPV6 mRNA

expression of duodenocytes and renal distal convoluted tubules⁽⁷⁾. In the current study, we demonstrated that duodenal TRPV6 mRNA expression is induced by sap solutions, independent of the dose of sap. Renal TRPV6 was not influenced by any of the dietary beverage conditions. TRPV5 mRNA was induced by sap and the high-calcium rescue solution, but the level of expression was not significantly higher than normal levels. CaBP-9k is a cytosolic protein with a high affinity for calcium ions^(23,24). In mammals, a number of tissues and organs, including the intestine, uterus, placenta, kidney and bone, express CaBP-9k^(16–18,25–31). Intestinal CaBP-9k is involved in intestinal calcium absorption and is regulated at the transcriptional and post-transcriptional levels by 1,25-dihydroxyvitamin D₃ (1,25-(OH)₂D₃), the hormonal form of vitamin D, in rodents^(32–34). It has also been shown that the expression of duodenal CaBP-9k may be linked to 1,25-dihydroxycholecalciferol in man⁽¹⁷⁾. Renal CaBP-9k is expressed at distal convoluted tubules, which are believed to facilitate calcium re-absorption⁽³⁵⁾. In the current study, we showed that duodenal CaBP-9k mRNA levels were not altered by sap.

Table 2. Serum calcium and parathyroid hormone (PTH) analysis‡
(Mean values with their standard errors)

Treatment	Serum calcium levels (mg/l)		Serum PTH (pg/ml)	
	Mean	SEM	Mean	SEM
Negative	70.1*	10.1	41.78*	9.97
Sap of <i>Acer mono</i>				
25 %	72.3*	8.2	35.07*	4.34
50 %	86.6	5.1	52.70*	17.04
100 %	90.2	13.2	46.98*	9.00
Rescue	86.6	6.8	5.41*†	1.30
Normal	91.4	4.5	14.20†	5.63

Mean values were significantly different from those of the normal group: * $P < 0.05$.
Mean values were significantly different from those of the negative control group:
† $P < 0.05$.

‡ For details of procedures, see Materials and methods.

However, sap solution increased renal *CaBP-9k* mRNA expression. The high calcium-containing rescue solution significantly decreased duodenal *CaBP-9k* and *TRPV6* transcription. In addition, it has previously been shown that high calcium intake attenuates *CaBP-9k* and *TRPV6* gene expression in mice⁽³⁶⁾. Thus, while sap appeared to regulate duodenal *TRPV6* and renal *CaBP-9k* expression, this effect was different from that of the rescue solution. The present result indicated that the underlying mechanism of regulation of these genes by sap may not involve calcium ions. If the high calcium content of sap was involved in the regulation of *CaBP-9k* and *TRPV6*, one would also expect the effect of sap on the expression patterns of these genes to be similar to previous studies^(9,36).

Generally, a calcium-deficient diet decreases serum calcium concentration and accelerates PTH secretion⁽³⁷⁾. In the current study, we demonstrated that the serum calcium levels of mice that received a 50 % or higher sap solution were similar to mice that were fed a normal diet. The present results suggest that the amount of calcium in a 50 % sap solution is sufficient to maintain normal calcium homeostasis. PTH plays a major role in regulating calcium homeostasis. PTH regulates the conversion of 25-hydroxyvitamin D to the active metabolite 1,25-(OH)₂D₃. PTH also activates dihydropyridine-sensitive channels that mediate calcium entry. It has also been suggested that microtubule-dependent exocytosis stimulated by PTH is required for the activation of calcium channels and calcium influx⁽³⁸⁾. 1,25-(OH)₂D₃ inhibits PTH secretion and parathyroid cell proliferation through the action of the vitamin D receptor^(39,40). It has also been shown that PTH secretion and bone mineralization correlate with calcium levels, rather than the activation state of the 1,25-(OH)₂D₃ receptor signalling pathway^(41,42). In the current study, serum PTH was up-regulated by a low-calcium diet compared to the normal group; however, the mice in the rescue group showed significantly lower serum PTH levels. The present results suggest that sap improves osteoporosis-like symptoms via a different mechanism than the high calcium-containing solution, similar to the regulation of the calcium-processing genes.

In summary, the positive effect of sap on osteoporosis-like symptoms was evaluated by measuring serum calcium

concentrations, femur density and length, and by a histological examination. Serum calcium levels were significantly lower in mice that received a low-calcium diet supplemented with commercial spring water, and in the 25 % sap solution group compared to the normal group, but not in the 50 and 100 % sap solution groups, or the rescue group. Femur density and length were significantly reduced in the negative and 25 % sap solution groups. The present results imply that a 50 % or higher sap solution could ameliorate osteoporosis-like symptoms induced by a low-calcium diet. The results of the histological analysis correlated with the bone density results. In addition, we examined the expression of calcium-processing genes in the duodenum and kidney, and showed that duodenal *TRPV6* and renal *CaBP-9k* were up-regulated by sap in a dose-independent manner. We propose that the sap of *A. mono* may improve low-calcium diet-induced osteoporosis-like symptoms by augmenting mineral levels in the body.

Acknowledgements

This work was supported by the Ministry of Education and Human Resources Development (MOE), the Ministry of Commerce, Industry and Energy (MOCIE) and the Ministry of Labor (MOLAB) through the fostering project of the Laboratory of Excellency. In addition, the authors appreciate a grant provided by Korea Forest Research Institute. G.-S. L., H.-S. B., I.-G. C., H.-Y. K. and E.-B. J. designed the research; G.-S. L., H.-S. B., M.-H. K., B.-M. L., S.-H. K., E.-M. J., K.-S. G., H.-J. J. and H.-J. L. performed the research; G.-S. L., H.-S. B. and E.-B. J. analysed the data; and G.-S. L. and E.-B. J. wrote the paper. The authors declare no conflicts of interest.

References

1. Angeles GBJ, Bond JS, Boyer TJ, *et al.* (2004) The cohesion-tension theory. *New Phytol* **163**, 451–452.
2. Kunkel G (1984) *Plants for Human Consumption*. Koenigstein: Koeltz Scientific Books.
3. Chopra RN, Nayar SL & Chopra IC (1986) *Glossary of Indian Medicinal Plants*. New Delhi: Council of Scientific and Industrial Research.
4. Pettifor JM, Marie PJ, Sly MR, du Bruyn DB, Ross F, Isdale JM, de Klerk WA & van der Walt WH (1984) The effect of differing dietary calcium and phosphorus contents on mineral metabolism and bone histomorphometry in young vitamin D-replete baboons. *Calcif Tissue Int* **36**, 668–676.
5. Diepens RJ, den Dekker E, Bens M, Weidema AF, Vandewalle A, Bindels RJ & Hoenderop JG (2004) Characterization of a murine renal distal convoluted tubule cell line for the study of transcellular calcium transport. *Am J Physiol Renal Physiol* **286**, F483–F489.
6. Hoenderop JG, Nilius B & Bindels RJ (2003) Epithelial calcium channels: from identification to function and regulation. *Pflügers Arch* **446**, 304–308.
7. Van Cromphaut SJ, Rummens K, Stockmans I, Van Herck E, Dijcks FA, Ederveen AG, Carmeliet P, Verhaeghe J, Bouillon R & Carmeliet G (2003) Intestinal calcium transporter genes are upregulated by estrogens and the reproductive cycle through vitamin D receptor-independent mechanisms. *J Bone Miner Res* **18**, 1725–1736.

8. Lee GS & Jeung EB (2007) Uterine TRPV6 expression during the estrous cycle and pregnancy in a mouse model. *Am J Physiol* **293**, E132–E138.
9. Lee GS, Lee KY, Choi KC, Ryu YH, Paik SG, Oh GT & Jeung EB (2007) A phenotype of a calbindin-D9k gene-knockout is compensated for by the induction of other calcium-transporter genes in a mouse model. *J Bone Miner Res* **22**, 1968–1978.
10. Bunker VW (1994) The role of nutrition in osteoporosis. *Br J Biomed Sci* **51**, 228–240.
11. Tranquilli AL, Lucino E, Garzetti GG & Romanini C (1994) Calcium, phosphorus and magnesium intakes correlate with bone mineral content in postmenopausal women. *Gynecol Endocrinol* **8**, 55–58.
12. Anderson MP, Hunt RD, Griffiths HJ, McIntyre KW & Zimmerman RE (1977) Long-term effect of low dietary calcium:phosphate ratio on the skeleton of *Cebus albifrons* monkeys. *J Nutr* **107**, 834–839.
13. Nieves JW (2005) Osteoporosis: the role of micronutrients. *Am J Clin Nutr* **81**, 1232S–1239S.
14. Harrington M & Cashman KD (2003) High salt intake appears to increase bone resorption in postmenopausal women but high potassium intake ameliorates this adverse effect. *Nutr Rev* **61**, 179–183.
15. Tucker KL, Hannan MT & Kiel DP (2001) The acid–base hypothesis: diet and bone in the Framingham Osteoporosis Study. *Eur J Nutr* **40**, 231–237.
16. Lee GS, Choi KC & Jeung EB (2006) Glucocorticoids differentially regulate expression of duodenal and renal calbindin-D9k through glucocorticoid receptor-mediated pathway in mouse model. *Am J Physiol* **290**, E299–E307.
17. Walters JR, Howard A, Lowery LJ, Mawer EB & Legon S (1999) Expression of genes involved in calcium absorption in human duodenum. *Eur J Clin Invest* **29**, 214–219.
18. Wissenbach U & Niemeyer BA (2007) TRPV6. In *Transient Receptor Potential (TRP) Channels. Handbook of Experimental Pharmacology*, vol. 179, pp. 221–234 [V Flockerzi and B Nilius, editors]. Berlin: Springer.
19. den Dekker E, Hoenderop JG, Nilius B & Bindels RJ (2003) The epithelial calcium channels, TRPV5 & TRPV6: from identification towards regulation. *Cell Calcium* **33**, 497–507.
20. Song Y, Peng X, Porta A, Takanaga H, Peng JB, Hediger MA, Fleet JC & Christakos S (2003) Calcium transporter 1 and epithelial calcium channel messenger ribonucleic acid are differentially regulated by 1,25 dihydroxyvitamin D3 in the intestine and kidney of mice. *Endocrinology* **144**, 3885–3894.
21. van Abel M, Hoenderop JG, van der Kemp AW, van Leeuwen JP & Bindels RJ (2003) Regulation of the epithelial Ca²⁺ channels in small intestine as studied by quantitative mRNA detection. *Am J Physiol Gastrointest Liver Physiol* **285**, G78–G85.
22. Weber K, Erben RG, Rump A & Adamski J (2001) Gene structure and regulation of the murine epithelial calcium channels ECaC1 and 2. *Biochem Biophys Res Commun* **289**, 1287–1294.
23. Christakos S, Gabriellides C & Rhoten WB (1989) Vitamin D-dependent calcium binding proteins: chemistry, distribution, functional considerations, and molecular biology. *Endocr Rev* **10**, 3–26.
24. Kumar R, Wieben E & Beecher SJ (1989) The molecular cloning of the complementary deoxyribonucleic acid for bovine vitamin D-dependent calcium-binding protein: structure of the full-length protein and evidence for homologies with other calcium-binding proteins of the troponin-C superfamily of proteins. *Mol Endocrinol* **3**, 427–432.
25. An BS, Choi KC, Kang SK, Lee GS, Hong EJ, Hwang WS & Jeung EB (2003) Mouse calbindin-D9k gene expression in the uterus during late pregnancy and lactation. *Mol Cell Endocrinol* **205**, 79–88.
26. An BS, Kang SK, Shin JH & Jeung EB (2002) Stimulation of calbindin-D9k mRNA expression in the rat uterus by octylphenol, nonylphenol and bisphenol. *Mol Cell Endocrinol* **191**, 177–186.
27. Choi KC, Leung PC & Jeung EB (2005) Biology and physiology of Calbindin-D9k in female reproductive tissues: involvement of steroids and endocrine disruptors. *Reprod Biol Endocrinol* **3**, 66.
28. Hong EJ, Choi KC & Jeung EB (2004) Induction of calbindin-D9k messenger RNA and protein by maternal exposure to alkylphenols during late pregnancy in maternal and neonatal uteri of rats. *Biol Reprod* **71**, 669–675.
29. Lee GS, Kim HJ, Jung YW, Choi KC & Jeung EB (2005) Estrogen receptor alpha pathway is involved in the regulation of Calbindin-D9k in the uterus of immature rats. *Toxicol Sci* **84**, 270–277.
30. Nguyen TH, Lee GS, Ji YK, Choi KC, Lee CK & Jeung EB (2005) A calcium binding protein, calbindin-D9k, is mainly regulated by estrogen in the pituitary gland of rats during estrous cycle. *Brain Res Mol Brain Res* **141**, 166–173.
31. Yun SM, Choi KC, Kim IH, An BS, Lee GS, Hong EJ, Oh GT & Jeung EB (2004) Dominant expression of porcine Calbindin-D9k in the uterus during a luteal phase. *Mol Reprod Dev* **67**, 251–256.
32. Darwish HM & DeLuca HF (1992) Identification of a 1,25-dihydroxyvitamin D3-response element in the 5'-flanking region of the rat calbindin D-9k gene. *Proc Natl Acad Sci U S A* **89**, 603–607.
33. Roche C, Bellaton C, Pansu D, Miller A 3rd & Bronner F (1986) Localization of vitamin D-dependent active Ca²⁺ transport in rat duodenum and relation to CaBP. *Am J Physiol* **251**, G314–G320.
34. Wasserman RH & Fullmer CS (1989) On the molecular mechanism of intestinal calcium transport. *Adv Exp Med Biol* **249**, 45–65.
35. Peng JB, Chen XZ, Berger UV, Vassilev PM, Brown EM & Hediger MA (2000) A rat kidney-specific calcium transporter in the distal nephron. *J Biol Chem* **275**, 28186–28194.
36. Van Cromphaut SJ, Dewerchin M, Hoenderop JG, Stockmans I, Van Herck E, Kato S, Bindels RJ, Collen D, Carmeliet P, Bouillon R & Carmeliet G (2001) Duodenal calcium absorption in vitamin D receptor-knockout mice: functional and molecular aspects. *Proc Natl Acad Sci U S A* **98**, 13324–13329.
37. Kato S, Mano T, Kobayashi T, Yamazaki N, Himeno Y, Yamamoto K, Itoh M, Harada N & Nagasaka A (2002) A calcium-deficient diet caused decreased bone mineral density and secondary elevation of estrogen in aged male rats – effect of menatrenone and elcatonin. *Metabolism* **51**, 1230–1234.
38. Bacskaï BJ & Friedman PA (1990) Activation of latent Ca²⁺ channels in renal epithelial cells by parathyroid hormone. *Nature* **347**, 388–391.
39. Carling T, Rastad J, Szabo E, Westin G & Akerstrom G (2000) Reduced parathyroid vitamin D receptor messenger ribonucleic acid levels in primary and secondary hyperparathyroidism. *J Clin Endocrinol Metab* **85**, 2000–2003.
40. Demay MB, Kiernan MS, DeLuca HF & Kronenberg HM (1992) Sequences in the human parathyroid hormone gene that bind the 1,25-dihydroxyvitamin D3 receptor and mediate transcriptional repression in response to 1,25-dihydroxyvitamin D3. *Proc Natl Acad Sci U S A* **89**, 8097–8101.
41. Li YC, Bolt MJ, Cao LP & Sitrin MD (2001) Effects of vitamin D receptor inactivation on the expression of calbindins and calcium metabolism. *Am J Physiol* **281**, E558–E564.
42. Song CZ, Tian X & Gelehrter TD (1999) Glucocorticoid receptor inhibits transforming growth factor-beta signaling by directly targeting the transcriptional activation function of Smad3. *Proc Natl Acad Sci U S A* **96**, 11776–11781.