

Editorial

Lichenology in Africa

In the past three years *The Lichenologist* has published c. 90 papers but only four of these were based on African material. Two of these were physiological papers co-authored by one of us (RB), whereas the other two were taxonomic and described new species from South Africa and Kenya. This contrasts dramatically with the 31 papers based on European material. Given the relative sizes of the continents and the diversity of habitats in Africa – from Mediterranean biomes in the north and far south, through deserts, tropical rainforests and savannah to temperate biomes in the south, and from sea level to alpine regions – this is an exceedingly poor return. Even within Africa, the distribution is uneven with three of the four papers being based on South African material, one on Kenyan, and none on western or northern material.

The history of lichenology in Africa dates back at least as far as Linnaeus. Johann König, a pupil of Linnaeus, visited Cape Town on his way to India in 1768 and made several collections, including the type collection of *Lichen chrysophthalmus* (Linnaeus 1771), which we now know as *Teloschistes chrysophthalmus* [as *chrysophthalmus*]. However, the first collector to spend any time collecting lichens in southern Africa was Carl Pehr Thunberg between 1772–1775 who, although based in Cape Town, travelled extensively in the interior (Thunberg 1794). In the following 150 years, a number of local collectors became active, including Miss [Olivia?] Armstrong and Peter MacOwen, while European visitors such as Josef Brunthaler and Friedrich Wilms made extensive expeditions into the interior. Their collections were, of course, taken back to Europe and deposited in European herbaria (LUND, G, W, ZT, etc) and many were described as new species by Johannes Müller (1885–1888), Edvard Vainio (e.g. Vainio 1901, 1926), Alexander Zahlbruckner (e.g. Zahlbruckner 1926, 1932, 1936) and especially Ernst Stizenberger in his *Lichenaea Africana* (Stizenberger 1890, 1891). These early collectors were primarily botanists who also collected other groups and more details of their activities can be found on the Biographical Database of Southern African Science (<http://www.s2a3.org.za/bio/Main.php>).

In the early 20th century Paul Andries van der Byl collected lichens extensively (van der Byl 1933a, b, 1935a, b), and in 1950 Ethel Doidge included lichens in her monumental compendium of southern African fungi (Doidge 1950). Subsequently, between 1956–1991, Ove Almborn issued six fascicles (150 collections) of an exsiccate, *Lichenes Africani*, and also published several other papers on African lichens (Almborn 1966, 1987, 1988, 1989). However, the first person based in South Africa to concentrate on the lichens of southern Africa was Franklin Brusse of SANBI, Pretoria who, between 1984–1994, published c. 40 papers describing taxonomic novelties from South Africa and Namibia. Although his main interest was in *Parmelia* s. lat., describing numerous new

species and making many new combinations, he also described many new species and nine new genera across a wide range of families (e.g. Brusse 1985a, b, 1987a, b, c, d, 1988a, b, 1994). More recently, one of us (AF) has published an updated checklist of South African lichens (Fryday 2015).

Luciana Zedda and co-workers have carried out some important work on terricolous lichen communities in Namibia and NW South Africa (e.g. Zedda & Rambold 2009; Zedda *et al.* 2009, 2011a, b) but otherwise the ecology of southern African lichens is poorly studied.

Lichenological exploration in eastern Africa started in the mid-1800s, with collectors mainly comprised of explorers, missionaries and European visitors with an interest in nature. The earliest collector in the region is perhaps W. G. Schimper who collected in Ethiopia in the 1840s. Other personalities who made small lichen collections from different parts of East Africa in the late part of the 19th century were Marquess Antinori, O. Beccari, J. Hannington, J. M. Hildebrandt, R. L. von Hohnel, C. H. E. W. Holst, H. H. Johnstone, W. Last, H. Meyer and G. Thomson, who made collections in Tanzania and Zanzibar. Prince Heinrich of Liechtenstein, Dr Pospischill and A. K. Ritter collected in Kenya and Tanzania and G. F. Scott Elliot collected in the Ruwenzori Mountains in Uganda. These collections were reported in several works (Baglietto 1875; Krempelhuber 1877; Stirton 1877–1878; Jatta 1882; Müller 1885–1888, 1890, 1893, 1894; Stein 1889; Steiner 1897; Vainio 1898).

In the early 1900s, collectors comprised both amateur and professional botanists. Amongst these were J. Brunthaler, B. Fink, L. Hauman, G. Lindau, R. A. Maas Geesteranus, Józef Motyka, Vicomte de Poncins, R. Pichi Sermolli, Professor Senni, B. Schroder, R. Wettstein and the Reverend Fathers of Consolata Missions. Results of these collections were published in several papers (Jatta 1908, 1909; Lindau 1911; Hue 1916; Zahlbruckner 1926, 1932; Zahlbruckner & Hauman 1936; Cengia Sambo 1937, 1938, 1939; Maas Geesteranus 1955; Motyka 1961) and later summarized in treatments of tropical lichens from Africa by Carroll Dodge (1953, 1956, 1957, 1959, 1964, 1971).

In the 1970s, Hildur Krog and Dougal Swinscow collected intensively in East Africa, their collections culminating in the publication of *Macrolichen Flora of East Africa* (Swinscow & Krog 1988). Subsequent excursions by other lichenologists greatly increased the knowledge of the East African lichen biota (Sérusiaux 1978, 1979, 1981, 1984a, b, 1989; Ahti *et al.* 1987; Almborn 1989; Elix 2002; Lücking & Kalb 2002; Jørgensen 2003; Kalb 2004, 2007, 2008; Alstrup & Aptroot 2005; Killmann & Fischer 2005; Sérusiaux & Diederich 2005; Frisch *et al.* 2006; Sérusiaux *et al.* 2006; Bock *et al.* 2007; Yeshitela 2008; Archer *et al.* 2009; Frisch & Tibell 2010). For the last two decades one of us (PK) has been collecting in Kenya and has updated the Kenyan lichen checklist, which will soon be published.

Cite this article: Fryday AM, Beckett RP and Kirika PM (2022) Lichenology in Africa. *Lichenologist* 54, 227–230. <https://doi.org/10.1017/S0024282922000329>



Elsewhere on the continent, North African lichenology has a long history, dating back to 1798 in Algeria. Checklists have recently been prepared for Algeria (Amrani *et al.* 2015, 2018) and Tunisia (Seaward *et al.* 2021) and there have been some inventories carried out in Morocco (Mohamed *et al.* 2021) and Egypt (Hegazy 2012). Unfortunately, recent references to the ecology, physiology or taxonomy of West African lichens are less easy to locate.

There have been very few studies on the ecophysiology and physiology of African lichens. Even the potential for using African lichens medicinally has received little attention (Adenubi *et al.* 2022). This is probably because there are relatively few physiologists that are familiar with lichenology on the staff of African universities, a problem that also extends into other subject areas. Furthermore, in Africa there are relatively few truly unique (physiologically speaking) habitats that would tempt lichenologists from the rest of the world to visit for research purposes. However, within southern Africa there are vast areas dominated by soil crust communities, within which lichens often play an important role. Thanks to the studies of Lucinda Zedda, Burkhart Büdel and their colleagues, progress has been made on characterizing these ecologically important systems (see references above, and Büdel *et al.* (2009)), but disappointingly their physiology remains to be investigated. Within this 'biome' there is a rather exceptional sub-type, that of the 'desert fog' lichens. Outside southern Africa, these communities are found only in a small number of places, namely California, the Peruvian coast and the Negev Desert in Israel. Otto Lange and his co-workers (Lange *et al.* 1991, 2006, 2007) elegantly demonstrated the remarkable ability of some Namibian desert 'fog' lichens to achieve net photosynthesis with non-liquid water. This work was continued by Maphangwa *et al.* (2012a), who initiated discussion on the important question of the likely impact of climate change on these lichens (Maphangwa *et al.* 2012b).

For the last 30 years, one of us (RB) has studied various aspects of the stress physiology of lichens. Although much of this work could have been carried out using lichens from anywhere in the world, in recent years work has focused on lichens from the Afrotropical vegetation. The study of Mkhize *et al.* (2022: this issue) compared the 'quenching' characteristics of photosynthesis in 'sun' and 'shade' collections of the same species of Afrotropical lichens. Results suggest that Afrotropical lichen photobionts can flexibly adjust the amount and type of their quenching.

It is to be hoped that in future more scientists will be prompted to study lichen physiology in Africa. What makes lichens special, and arguably separates them from the vast majority of other photosynthetic organisms, is their desiccation tolerance, and this holds innovative potential. Once the molecular mechanisms have been elucidated, technologies can be used to transfer genes for stress tolerance to crop plants (Farrant & Hilhorst 2022). These studies are of great importance, since Africa is predicted to bear the brunt of future climate change which will no doubt increase abiotic stress and challenge the security of food production.

The present volume is an attempt to bring African lichens and lichenology to the attention of the wider community, while at the same time promoting the study of lichens and lichenology within the continent itself. In this issue, Kaasalainen *et al.* study the diversity of the genus *Peltigera* on Mt Kilimanjaro in Tanzania and report eight species including the new species, *P. alkalicola*; Kirika *et al.* show that the pantropical species *Canoparmelia texana* consists of two independent lineages, both of which occur in

Kenya; Medeiros & Lutzoni provide a modern treatment of the *Graphidaceae* genera with hyaline ascospores in South Africa, describing one new species, *Allographa oldayana*, making two new combinations and reporting 11 other species as new to South Africa; Nadel & Clerc provide notes on the genus *Usnea* from the islands of São Tomé and Príncipe in tropical West Africa, reporting 15 taxa, of which two, *U. beckeri* and *U. longiciliata*, are described as new to science; Temu *et al.* investigate the molecular, morphological and chemical variation of the *Usnea pectinata* aggregate in Tanzania, and São Tomé and Príncipe; Farkas & Muhoro present an identification key for parmelioid taxa in Kenya; Mkhize *et al.* investigate the adaptations of photosynthesis in sun and shade in populations of some Afrotropical lichens (see above) and Kitara *et al.* examine the distribution pattern and host tree specificity of *Lobaria pulmonaria* in tropical montane forest landscapes in Tanzania. Finally, Aptroot reviews a book on foliicolous lichens and their lichenicolous fungi in Macaronesia and Atlantic Europe authored by van den Boom.

To the best of our knowledge, there are no research groups in Africa (or anywhere else for that matter) focused on the systematics or ecology of African lichens, but we know that interest exists because of our own personal experience and the number of posts on social media requesting help with identifications. However, the lack of field guides and local experts means that getting started is an almost impossible task. Additionally, most of the hundreds of species described by Stizenberger, Zahlbruckner and others have not been reported since and are known only from their type collection, and these collections are in European herbaria. Consequently, anyone attempting to find the correct name for an unknown collection is left with the thankless task of trawling through the protologues of numerous possible taxa, which invariably lack the detailed information necessary to make a positive identification.

In spite of the recent activity in southern and eastern Africa, as elsewhere on the continent, large parts of these regions still remain underexplored and the lichen biota urgently requires revision to fully understand its diversity and biogeography, especially with the use of molecular tools. However, as can be seen from the contributions to this volume, the potential for exciting new discoveries on the continent is enormous and we hope this special issue will be a springboard to future work.

References

- Adenubi OT, Famuyide IM, McGaw LJ and Eloff JN (2022) Lichens: an update on their ethnopharmacological uses and potential as sources of drug leads. *Journal of Ethnopharmacology* **298**, 115657.
- Ahti T, Krog H and Swinscow TDV (1987) New or otherwise interesting *Cladonia* species in East Africa. *Annales Botanici Fennici* **24**, 85–94.
- Almborn O (1966) Revision of some lichen genera in southern Africa I. *Botaniska Notiser* **119**, 70–112.
- Almborn O (1987) Lichens at high altitudes in southern Africa. *Bibliotheca Lichenologica* **25**, 401–417.
- Almborn O (1988) Some distribution patterns in the lichen flora of South Africa. *Monographs in Systematic Botany from the Missouri Botanical Garden* **25**, 429–432.
- Almborn O (1989) Revision of the lichen genus *Teloschistes* in central and southern Africa. *Nordic Journal of Botany* **8**, 521–537.
- Alstrup V and Aptroot A (2005) Pyrenocarpous lichens from Tanzania and Kenya. *Cryptogamie, Mycologie* **26**, 265–271.
- Amrani S, Nacer A, Noureddine N-E and Seaward MRD (2015) Lichenological exploration of Algeria: historical overview and annotated bibliography, 1799–2013. *Willdenowia* **45**, 15–34.

- Amrani S, Seaward MRD, Sipman HJM and Feuerer T** (2018) Lichenological exploration of Algeria II: checklist of lichenized, lichenicolous and allied fungi. *Herzogia* **31**, 817–892.
- Archer AW, Elix JA, Fischer E, Killmann D and Sérusiaux E** (2009) The lichen genus *Pertusaria* (Ascomycota) in Central Africa (Congo/Kivu, Rwanda and Burundi) and western Kenya. *Nova Hedwigia* **88**, 309–333.
- Baglietto F** (1875) Lichenes in regione Bogos Abyssinia e septentrionalis lecti ab O. Beccari. *Nuovo Giornale Botanico Italiano* **7**, 239–254.
- Bock C, Hauck M and Fischer E** (2007) The lichen flora of Rwanda: an annotated checklist. *Willdenowia* **37**, 563–575.
- Brusse F** (1985a) *Corynecystis*, a new lichen genus from the Karoo, South Africa. *Bothalia* **15**, 552–553.
- Brusse F** (1985b) *Glyphopeltis* (Lecideaceae), a new lichen genus from southern Africa. *Lichenologist* **17**, 267–268.
- Brusse F** (1987a) *Psathyrophlyctis*, a new lichen genus from southern Africa. *Bothalia* **17**, 182–184.
- Brusse FA** (1987b) *Coronoplectrum*, a new lichen genus from the Namib Desert, South West Africa/Namibia. *Mycotaxon* **28**, 131–135.
- Brusse FA** (1987c) *Echidnocymbium* (Biatoraceae), a new lichen genus from southern Africa. *Mycotaxon* **29**, 173–176.
- Brusse FA** (1987d) *Schistoplaca* (Lichenes, Biatoraceae), a new lichen genus. *Mycotaxon* **29**, 245–249.
- Brusse F** (1988a) *Lithoglypha*, a new lichen genus from Clarens Sandstone. *Bothalia* **18**, 89–96.
- Brusse F** (1988b) *Schizodiscus*, a new porpidioid lichen genus from Drakensberg. *Bothalia* **18**, 94–96.
- Brusse FA** (1994) A remarkable new lichen genus *Catarrhospora* (Ascomycotina, Porpidiaceae), from Cape Floral Kingdom, South Africa. *Mycotaxon* **52**, 501–512.
- Büdel B, Darienko T, Deutschewitz K, Dojani S, Friedl T, Mohr K, Salisch M, Reisser W and Weber B** (2009) Southern African biological soil crusts are ubiquitous and highly diverse in drylands, being restricted by rainfall frequency. *Microbial Ecology* **57**, 229–247.
- Cengia Sambo M** (1937) Lichenes Africae Orientalis Italicae. *Nuovo Giornale Botanico Italiano* **44**, 456–483.
- Cengia Sambo M** (1938) Licheni del Kenia e del Tanganica raccolti dai Rev. Padri della consolata. *Nuovo Giornale Botanico Italiano* **45**, 364–387.
- Cengia Sambo M** (1939) Licheni dell'A.O.I. [Africa Orientale Italiana]. Raccolti dal Console Prof. Senni e dal Centurione ing. Giordano ufficiali della Milizia Naz. Forestale di Addis Abeba nel 1937. *Nuovo Giornale Botanico Italiano* **46**, 437–455.
- Dodge CW** (1953) Some lichens from tropical Africa. *Annals of the Missouri Botanical Garden* **40**, 271–401.
- Dodge CW** (1956) Some lichens of tropical Africa. II. *Usnea*. *Annals of the Missouri Botanical Garden* **43**, 381–396.
- Dodge CW** (1957) Some lichens of tropical Africa. II. *Usnea* (Continued). *Annals of the Missouri Botanical Garden* **44**, 1–76.
- Dodge CW** (1959) Some lichens of tropical Africa. III. *Parmeliaceae*. *Annals of the Missouri Botanical Garden* **46**, 39–193.
- Dodge CW** (1964) Some lichens of tropical Africa. IV. *Dermatocarpaceae* to *Pertusariaceae*. *Beihefte zur Nova Hedwigia* **12**, 1–282.
- Dodge CW** (1971) Some lichens of tropical Africa. V. *Lecanoraceae* to *Physciaceae*. *Beihefte zur Nova Hedwigia* **38**, 1–225.
- Doidge EM** (1950) The South African fungi and lichens to the end of 1945. *Bothalia* **5**, 1–1094.
- Elix JA** (2002) New species of *Xanthoparmelia* (lichenized *Ascomycotina*, *Parmeliaceae*) from Africa. *Lichenologist* **34**, 283–291.
- Farrant JM and Hilhorst H** (2022) Crops for dry environments. *Current Opinion in Biotechnology* **74**, 84–91.
- Frisch A** (2006) Contributions towards a new systematics of the lichen family *Thelotrema* I. The lichen family *Thelotrema* in Africa. *Bibliotheca Lichenologica* **92**, 3–370.
- Frisch A and Tibell L** (2010) New data on crustose mazaediate lichens from tropical Africa. *Bibliotheca Lichenologica* **104**, 323–332.
- Fryday AM** (2015) A new checklist of lichenized, lichenicolous and allied fungi reported from South Africa. *Bothalia* **45**, a148.
- Hegazy TA** (2012) A new lichen record at Ain Quraychat, Siwa Oasis, Egypt. *Catrina* **7**, 79–85.
- Hue A-M** (1916) Lichenes Poncinsianos in Africa tropico-orientali anglica a Vicecomite de Poncins anno 1912 lectos elaboravit. *Memoire de la Societé Botanique de France* **28**, 1–24.
- Jatta A** (1882) Licheni africani raccolti nello Scioa dal Marchese Antinori. *Nuovo Giornale Botanico Italiano* **14**, 169–175.
- Jatta A** (1908) Species novae in excelsis Ruwenzori in expeditione Ducis Aprutii lectae. *Annali di Botanica* **6**, 407–409.
- Jatta A** (1909) Licheni del Ruwenzori. In Roccati A (ed.), *Il Ruwenzori, Parte Scientifica Vol. 1., Zoologia e Botanica*. Milano: Ulrico Hoepli, pp. 537–547.
- Jørgensen PM** (2003) Notes on African *Pannariaceae* (lichenized ascomycetes). *Lichenologist* **35**, 11–20.
- Kalb K** (2004) New or otherwise interesting lichens II. *Bibliotheca Lichenologica* **88**, 301–329.
- Kalb K** (2007) New or otherwise interesting lichens III. *Bibliotheca Lichenologica* **95**, 297–316.
- Kalb K** (2008) New or otherwise interesting lichens. IV. *Sauteria* **15**, 239–248.
- Killmann D and Fischer E** (2005) New records for the lichen flora of Rwanda, East Africa. *Willdenowia* **35**, 193–204.
- Krempelhuber A** (1877) Neue Beiträge zu Afrika's Flechten-Flora. *Linnaea* **41**, 135–144.
- Lange OL, Meyer A, Ullmann I and Zellner H** (1991) Microclimate conditions, water content and photosynthesis of lichens in the coastal fog zone of the Namib Desert: measurements in the fall. *Flora* **185**, 133–266.
- Lange OL, Green TGA, Melzer B, Meyer A and Zellner H** (2006) Water relations and CO₂ exchange of the terrestrial lichen *Teloschistes capensis* in the Namib fog desert: measurements during two seasons in the field and under controlled conditions. *Flora* **201**, 268–280.
- Lange OL, Green TGA, Melzer B, Meyer A and Zellner H** (2007) Water relations and carbon dioxide exchange of epiphytic lichens in the Namib fog desert. *Flora* **202**, 479–487.
- Lindau G** (1911) Lichenes. In Mildbraed J (ed.), *Wissenschaftliche Ergebnisse der Deutschen Zentral-Afrika-Expedition 1907–1908 unter Führung Adolf Friedrichs, Herzog zu Mecklenburg. II, Botanik*. Leipzig: Klinckschardt & Biermann, pp. 101–110.
- Linnaeus C** (1771) *Mantissa Plantarum II*. Holmiae: Laurentius Salvius.
- Lücking R and Kalb K** (2002) New species and further additions to the foliicolous lichen flora of Kenya (East Africa), including the first lichenicolous *Aulaxina* (*Ostropales: Gomphillaceae*). *Botanical Journal of the Linnean Society* **139**, 171–180.
- Maas Geesteranus RA** (1955) Some lichenological observations in Kenya. *Webbia* **11**, 519–523.
- Maphangwa KW, Musil CF, Raitt L and Zedda L** (2012a) Differential interception and evaporation of fog, dew and water vapour and elemental accumulation by lichens explain their relative abundance in a coastal desert. *Journal of Arid Environments* **82**, 71–80.
- Maphangwa KW, Musil CF, Raitt L and Zedda L** (2012b) Experimental climate warming decreases photosynthetic efficiency of lichens in an arid South African ecosystem. *Oecologia* **169**, 257–268.
- Mkhize KGW, Minibayeva F and Beckett RP** (2022) Adaptions of photosynthesis in sun and shade in populations of some Afromontane lichens. *Lichenologist* **54**, 319–329.
- Mohamed E, Mohammed L and Abdelhay EG** (2021) Study of lichens of the Moroccan Atlantic Coast Safi-Essaouira: bioindication of climate disturbances and nitrogen pollution. *International Journal of Science and Research* **10**, 696–703.
- Motyka J** (1961) *Usnea* a R. A. Maas Geesteranus in Africa orientali et australi anno 1949 collectae. *Persoonia* **1**, 415–431.
- Müller J** (1885–1888) Lichenologische Beiträge XXI–XXX. *Flora (Regensburg)* **68–71**.
- Müller J** (1890) Lichenes Africae tropico-orientalis. *Flora (Regensburg)* **73**, 334–347.
- Müller J** (1893) Lichenes africani. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* **19**, 505–510.
- Müller J** (1894) Lichenes usambarenses. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* **20**, 238–298.
- Seaward MRD, Khadhri A and Amrani S** (2021) Checklist of the lichens and lichenicolous fungi of Tunisia. *Herzogia* **34**, 400–427.

- Sérusiaux E** (1978) Contribution à l'étude des lichens du Kivu (Zaïre), du Rwanda et du Burundi II. Espèces nouvelles de lichens foliicoles. *Lejeunia* **90**, 1–18.
- Sérusiaux E** (1979) Contribution à l'étude des lichens du Kivu (Zaïre), du Rwanda et du Burundi IV. Le genre *Stereocaulon*. *Revue Bryologique et Lichénologique* **45**, 1–43.
- Sérusiaux E** (1981) Contribution à l'étude des lichens du Kivu (Zaïre), du Rwanda et du Burundi VI. Les genres *Coccocarpia* Pers. et *Lobaria* (Schreb.) Hoffm. *Cryptogamie, Bryologie, Lichénologie* **2**, 461–469.
- Sérusiaux E** (1984a) Contribution à l'étude des lichens du Kivu (Zaïre), du Rwanda et du Burundi VIII. New and interesting species of parmeliaceous lichens. *Bryologist* **87**, 1–11.
- Sérusiaux E** (1984b) New species or interesting records of foliicolous lichens. *Mycotaxon* **20**, 283–306.
- Sérusiaux E** (1989) *Echinoplaca furcata*, a new species of foliicolous lichen (*Gomphillaceae*) from Rwanda. *Mycotaxon* **35**, 237–242.
- Sérusiaux E and Diederich P** (2005) *Lambinonia*, a new genus for *Melanconium strigulae*, a lichenicolous hyphomycete on foliicolous species of *Strigula*. *Lichenologist* **37**, 499–506.
- Sérusiaux E, Fischer E and Killmann D** (2006) *Nyungwea*, a new genus of lichen with goniocyst-producing stipes from Rwanda and Uganda (E Africa). *Lichenologist* **38**, 115–121.
- Stein B** (1889) Ueber afrikanische Flechten. *Jahres-Bericht der Schlesischen Gesellschaft für Vaterländische Cultur* **66**, 133–142.
- Steiner J** (1897) Flechten aus Britisch-Ostafrika. *Sitzungsberichte, Akademie der Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Klasse* **106**, 207–234.
- Stirton J** (1877–1878) On certain lichens belonging to the genus *Parmelia*. *Scottish Naturalist* **4**, 200–203.
- Stizenberger E** (1890) Lichenaea Africana. *Bericht über die Thätigkeit der St. Gallischen, Naturwissenschaftlichen Gesellschaft* **1888–89**, 105–249.
- Stizenberger E** (1891) Lichenaea Africana. *Bericht über die Thätigkeit der St. Gallischen, Naturwissenschaftlichen Gesellschaft* **1889–90**, 133–268.
- Swinscow TDV and Krog H** (1988) *Macrolichens of East Africa*. London: British Museum (Natural History).
- Thunberg CP** (1794) *Prodromus Plantarum Capensium, quas in Promontorio Bonae Spei Africes Annis 1772–1775 Collegit*. Upsaliae: Edman.
- Vainio EA** (1898) Lichenes a G. F. Scott-Elliot in vicinis montis Ruwenzori in Africa centrali annis 1893–94 collecti. *Beiblatt zur Hedwigia* **37**, 39–44.
- Vainio E** (1901) Lichenes. In *Catalogue of the African Plants Collected by Dr Friedrich Welwitsch in 1853–61, Vol. 2, pt. 2 Cryptogamia*. London: British Museum (Natural History), pp. 396–463.
- Vainio E** (1926) Lichenes Africani novi. *Annales Universitatis Fennicae Åboensis, Series A* **2**(3), 1–33.
- van der Byl PA** (1933a) Korsmosse van die Unie van Suid-Afrika. I. Familie *Roccellaceae*. II. Familie *Teloschistaceae*. *Annals of the University of Stellenbosch* **11A** (6), 1–18.
- van der Byl PA** (1933b) Korsmosse van die Unie van Suid-Afrika. III. Familie *Cladoniaceae*. *Annals of the University of Stellenbosch* **11A** (4), 1–14.
- van der Byl PA** (1935a) Korsmosse van die Unie van Suid-Afrika. IV. Die geslag *Ramalina*. *Annals of the University of Stellenbosch* **13A** (1), 1–18.
- van der Byl PA** (1935b) Korsmosse van die Unie van Suid-Afrika. V. Familie *Collemaaceae*. *Annals of the University of Stellenbosch* **13A** (4), 1–13.
- Yeshitela K** (2008) *Effects of anthropogenic disturbance on the diversity of foliicolous lichens of East Africa: Goder (Ethiopia), Budongo (Uganda) and Kakamega (Kenya)*. Ph.D. thesis, University of Koblenz-Landau.
- Zahlbruckner A** (1926) Afrikanische Flechten (Lichenes). *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* **60**, 468–552.
- Zahlbruckner A** (1932) Lichenes in Africa lecti. *Annales de Cryptogamie Exotique* **5**, 198–275.
- Zahlbruckner A** (1936) Neue Flechten, XII. *Annales Mycologici* **34**, 159–179.
- Zahlbruckner A and Hauman L** (1936) Les lichens des hautes altitudes au Ruwenzori. *Mémoires de l'Institut Royal Colonial Belge, Section des Sciences Naturelles et Médicales* **5**, 1–31.
- Zedda L and Rambold G** (2009) Diversity and ecology of soil lichens in the Knersvlakte (South Africa). *Bryologist* **112**, 19–29.
- Zedda L, Schultz M and Rambold G** (2009) Diversity of epiphytic lichens in the savannah biome of Namibia. *Herzogia* **22**, 153–164.
- Zedda L, Kong S-M and Rambold G** (2011a) Morphological groups as a surrogate for soil lichen biodiversity in Southern Africa. *Bibliotheca Lichenologica* **106**, 391–408.
- Zedda L, Gröngroft A, Schultz M, Petersen A, Mills A and Rambold G** (2011b) Distribution patterns of soil lichens across the principal biomes of southern Africa. *Journal of Arid Environments* **75**, 215–220.

Alan M. Fryday, Richard P. Beckett and Paul M. Kirika