



A rare case of human taeniasis caused by *Taenia saginata* with species undetermined cysticercosis

Jie Hou¹ , Weilin Chen¹, Rong Chen², Chunlei He³, Ying Ma¹ and Junyan Qu⁴

Research Article

Cite this article: Hou J, Chen W, Chen R, He C, Ma Y, Qu J (2023). A rare case of human taeniasis caused by *Taenia saginata* with species undetermined cysticercosis. *Parasitology* **150**, 240–247. <https://doi.org/10.1017/S003118202200169X>

Received: 29 July 2022
Revised: 9 November 2022
Accepted: 1 December 2022
First published online: 19 December 2022

Keywords:

Human cysticercosis; *Taenia asiatica*; *Taenia saginata*; *Taenia solium*; taeniasis

Authors for correspondence:

Ying Ma,
E-mail: majiyang72@hotmail.com;
Junyan Qu,
E-mail: qujunyan15647@wchscu.cn

¹Department of Laboratory Medicine, West China Hospital, Sichuan University, Chengdu, Sichuan, China; ²Department of Clinical Laboratory, The First People's Hospital of Shuangliu District, Chengdu/West China (Airport) Hospital, Sichuan University, Chengdu, Sichuan, China; ³West China School of Medicine, Sichuan University, Chengdu, Sichuan, China and ⁴Center of Infectious Disease, West China Hospital, Sichuan University, Chengdu, China

Abstract

Taeniasis and cysticercosis, which are caused by *Taenia saginata*, *Taenia solium* and *Taenia asiatica*, are zoonotic parasitic infections with a significant disease burden worldwide. There is consensus amongst experts that *T. saginata* is a common tapeworm that causes taeniasis in humans as opposed to cysticercosis. This case study of a middle-aged Tibetan man conducted in 2021 challenges the prevailing notion that *T. saginata* exclusively causes taeniasis and not cysticercosis by documenting symptoms and laboratory studies related to both taeniasis and multiple cysticercosis. The patient's medical record with the symptoms of taeniasis and cysticercosis was reviewed, and the tapeworm's proglottids and cyst were identified from the patient by morphological evaluation, DNA amplification and sequencing. The patient frequently experienced severe headaches and vomiting. Both routine blood screenings and testing for antibodies against the most common parasites were normal. After anthelmintic treatment, an adult tapeworm was found in feces, and medical imaging examinations suggested multiple focal nodules in the brain and muscles of the patient. The morphological and molecular diagnosis of the proglottids revealed the Cestoda was *T. saginata*. Despite the challenges presented by the cyst's morphology, the molecular analysis suggested that it was most likely *T. saginata*. This case study suggests that *T. saginata* infection in humans has the potential to cause human cysticercosis. However, such a conclusion needs to be vetted by accurate genome-wide analysis in patients with *T. saginata* taeniasis associated with cysts. Such studies shall provide new insights into the pathogenicity of *T. saginata*.

Introduction

The zoonotic infection taeniasis is most commonly caused by *Taenia saginata*, *Taenia solium* and *Taenia asiatica*, which are associated with ingesting raw or undercooked, infected beef meat, pork meat or pork viscera, respectively (Ito *et al.*, 2003; Eichenberger *et al.*, 2020). The life cycle of pathogenic *Taenia* species involves humans as definitive hosts, who harbour the adult stage in their small intestines. Cattle (*T. saginata*) or pigs (*T. solium* and *T. asiatica*) can act as intermediate hosts, who hold the larval stages as cysts that dwell themselves under the skin and in the muscles, nervous system or visceral organs such as the liver of the hosts (Qian *et al.*, 2020). Tapeworm eggs or gravid proglottids are passed in feces, and the eggs can survive for days to months in the environment (Jansen *et al.*, 2021). Humans can also serve as accidental intermediate hosts when exposed to embryonated eggs (this role is normally fulfilled by *T. solium*) released by themselves (autoinfestation) or by another tapeworm carrier living in close contact with the subject or involved in contaminated food and water, causing infection in various organs of the human body with the larvae also known as cysticerci (human cysticercosis) (Garcia *et al.*, 2003; Garcia, 2018; Alroy and Gilman, 2020). While *T. saginata* is responsible for a significant number of cases of taeniasis in humans, it is not thought to be the aetiological agent of cysticercosis in humans because they do not appear to be an accidental intermediate host for this parasite (Garcia *et al.*, 2020). It is not known whether *T. asiatica* causes cysticercosis in humans (Eom *et al.*, 2020).

Twelve cases of *T. saginata* cysticercosis in humans had been described in the literature before 1972, and these diagnoses were based on autopsy and morphological examination by hookless scoleces (De Rivas, 1937; Tanasescu and Repciuc, 1939; Asenjo and Bustamente, 1950; Niiio, 1950; Bacigalupo, 1956; Abuladze, 1964; Goldsmid, 1966; Pawlowski and Schultz, 1972). In addition, *T. saginata* infection in adults can occur simultaneously with human cysticercosis (Pawlowski and Schultz, 1972). The infection of humans with the larvae (cysticerci) of *T. saginata* appears to be an exceptional situation (Pawlowski and Schultz, 1972). Medical imaging testing, clinical/exposure criteria, serologic testing, histological pathology and molecular identification all help in the diagnosis of human cysticercosis (Del Brutto *et al.*, 2017). Due to the lack of molecular methods readily available in most hospital settings, proper species identification in all reported cases of human cysticercosis is impossible.

© The Author(s), 2022. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Numerous molecular procedures have been developed to identify *Taenia* spp. responsible for human taeniasis, using mitochondrial and ribosomal DNA, repetitive DNA sequences and/or genes encoding relevant antigens as specimens (Flores *et al.*, 2018). The *cytochrome c oxidase I* (COX1) gene is a widely used DNA barcoding marker because the rapid rate of evolution allows for the differentiation of not just closely related species, such as the *Taenia* genus, but also phylogeographic groups within a single species (Hebert *et al.*, 2003; Galimberti *et al.*, 2012; Zhang *et al.*, 2014). Evidence suggests the HDP2 segment is a ribosomal DNA sequence useful for species-specific molecular diagnosis of human intestinal taeniasis (Gonzalez *et al.*, 2000; Flores *et al.*, 2018). In this case study, we identified the case of a middle-aged Tibetan man with taeniasis owing to *T. saginata* and cysticercosis that had impacted his brain and muscles diagnosed by morphological and molecular methods. We also used COX1 to create a phylogenetic tree of *Taenia*.

Materials and methods

Clinical examinations

The electronic medical records of the study subject were retrospectively reviewed, and the following demographic and clinical information was collected: age, gender, past medical history, physical examinations, diagnosis, underlying conditions, drug use, iatrogenic procedures, laboratory tests, computed tomography (CT), enhanced magnetic resonance imaging (MRI), ultrasound scan and clinical outcomes.

Morphological evaluation

Two obvious cysts were detected in the muscles under the right anterior thoracic wall and right midaxillary line through palpation and ultrasound scan, respectively. The cyst biopsy of the right midaxillary line and the proglottids from the patient after deworming treatment was collected and observed morphologically. The gravid proglottids and cystic lesions were fixed in a 10% formalin solution, stained with the hydrochloric acid-carmin solutions and observed under a microscope for pathological diagnosis.

Amplification and sequencing

A portion of the cyst isolated from the right midaxillary line and proglottids was stored in 70% ethanol at -20°C until subsequent molecular diagnosis. Total genomic DNA was extracted by using the *Trelief*TM Animal Genomic DNA Kit (Tsingke Biotechnology, Beijing, China) in accordance with the manufacturer's instructions. Mitochondrial DNA was analysed by polymerase chain reaction (PCR) targeting COX1 using species-specific forward primers Tsag_COX1/F for *T. saginata* (positions 322–348) and a common reverse primer Tae_COX1/R for *Taenia* (positions 1148–1129) (Yamasaki *et al.*, 2004), cestode forward primers JB3 (positions 1–24) and reverse primers JB4.5 (positions 444–421) (Bowles *et al.*, 1992; Bowles and McManus, 1994; Tappe *et al.*, 2016). The HDP2 fragment was amplified with the forward primer PTs7S35F1 and the reverse primer PTs7S35R1 (Gonzalez *et al.*, 2000). Details of the PCR primers and the reaction conditions are provided in Table 1. PCRs were run in a 30- μL reaction mixture composed of 15- μL I-5TM 2 \times High-Fidelity Master Mix (Tsingke Biotechnology), 1- μL DNA template, 10 pmol of each primer and 12- μL double-distilled water. The thermocycler parameters of the PCR amplification were as follows: 98 $^{\circ}\text{C}$ for 3 min; 39 cycles composed of denaturation at 98 $^{\circ}\text{C}$ for 10 s, annealing according to the primers and elongation at 72 $^{\circ}\text{C}$ for 15 s. After

Table 1. Primer pairs used for PCRs

Order	Target	Primer names	Orientation	Positions	Sequences (5'–3')	PCR products (bp)	Annealing	Samples	BLAST per cent identity ^a
1	COX1	Tsag_COX1/F	Forward	322–348	TTGATTCCCTTCGATGGCTTTCTTTTG	827	60 $^{\circ}\text{C}$ for 30 s	Proglottids cyst ^b	<i>Taenia saginata</i> 99.27–100%
		Tae_COX1/R	Backward	1148–1129	GACATAACATAATGAAATG				<i>Taenia asiatica</i> 96.25–96.37%
2	COX1	JB3	Forward	1–24	TTTTTTGGGCATCCTGAGGTTTAT	444	51 $^{\circ}\text{C}$ for 10 s	Proglottids cyst ^c	<i>T. saginata</i> 99.24–100%
		JB4.5	Backward	444–421	TAAAGAAAGAACATAATGAAATG				
3	HDP2	PTs7S35F1	Forward	1–22	CAGTGGCATAGCAGAGGAGGAA	599	57 $^{\circ}\text{C}$ for 25 s	Cyst	<i>T. saginata</i> 99.67–99.83%
		PTs7S35R1	Backward	599–577	GGACGAAGAATGGAGTTGAAGGT				<i>T. asiatica</i> 95.66–99.67%
									<i>Taenia solium</i> 92.99–97.33%

^aThe per cent identity was obtained from the NCBI BLAST webpage and the organism was limited to the *Taenia* (taxid: 6202) genus and the query coverage was 100%.

^bOnly proglottids were successfully amplified.

^cBoth proglottids and cysts were successfully amplified.

Table 2. Reference COX1 sequences of *Taenia* species in GenBank

<i>Taenia</i> species	GenBank accession no.
<i>T. asiatica</i>	NC_004826.2
<i>T. saginata</i>	NC_009938.1
<i>T. solium</i>	NC_004022.1
<i>Taenia hydatigena</i>	NC_012896.1
<i>Taenia multiceps</i>	NC_012894.1
<i>Taenia pisiformis</i>	NC_013844.1
<i>Taenia crassiceps</i>	NC_002547.1
<i>Taenia arctos</i>	NC_024590.1
<i>Taenia crocutae</i>	NC_024591.1
<i>Taenia regis</i>	NC_024589.1
<i>Taenia laticollis</i>	NC_021140.1
<i>Taenia madoquae</i>	NC_021139.1
<i>Taenia martis</i>	NC_020153.1
<i>Taenia ovis</i>	NC_021138.1
<i>Taenia serialis</i>	NC_021457.1
<i>Taenia twitchelli</i>	NC_021093.1

39 cycles, the reaction was terminated with an elongation step at 72°C for 5 min, followed by a final hold at 4°C. The amplicons were observed using a 1.5% agarose gel under ultraviolet light and then purified, after which Sanger sequencing methods were carried out at Tsingke Biotechnology or Sangon Biotech (Shanghai, China). The sequenced amplicons were compared to the GenBank database entries using NCBI BLAST (Johnson et al., 2008), wherein the organism was limited to *Taenia* (taxid: 6202) and the query coverage was 100%.

Phylogenetic analysis

Multiple sequence alignment was constructed with Clustal W, Clustal Omega and SnapGene software. Phylogenetic analysis of the sequences of COX1 identified in this study and the reference sequences of *Taenia* species available in the GenBank database (Table 2) were constructed using the maximum-likelihood method by MAGA X. The bootstrap values were calculated with 2000 replications, and the nucleotide substitution of the Tamura–Nei model was adapted.

Results

Case presentation

A middle-aged Tibetan man from China was admitted to our hospital with a 7-year history of recurring severe headaches and vomiting. The headache had started with vomiting, nausea, dizziness and chest tightness, especially on the right side, without any apparent predisposing cause. Weekly, he had 2 or 3 episodes of non-projectile vomiting without bile or coffee-like particles, and the headaches lasted for nearly a day each time. There were no other symptoms, such as fever, chills, convulsions, seizures, increased intracranial pressure or other discomforts. He had not been treated for these signs and symptoms before admitting to our hospital in December 2020. After being fully diagnosed in our hospital, his comorbidities included hypertension, fatty liver, liver insufficiency, paranasal sinusitis and intestinal bacterial infection. The immunosuppression was not found. Additionally, he spent a significant amount of time in a pastoral area that is an

epidemic province for taeniasis, although no tapeworm infections were diagnosed before his admission to our hospital. He used to ingest raw beef, which might have led to the ingestion of the cysticerci of *T. saginata*. Moreover, as far as pork consumption is concerned, his family's eating habit, like most Han Chinese, were eating it cooked, ruling out the possibility of related parasites.

The routine laboratory blood tests, including eosinophil and lymphocyte count, were completely normal apart from the elevated inflammation markers (C-reactive protein and neutrophil count). *Schistosomes*, *Clonorchis sinensis*, *Echinococcus granulosus* and *Toxoplasma gondii* antibodies were negative in his serum, as were *Cryptococcus* capsular antigen and next-generation sequencing results from cerebrospinal fluid. Multiple intracranial nodules affecting the supratentorial and infratentorial cerebral parenchyma were shown in detail on CT and MRI of the head, indicating possible intracranial parasitic infection (Fig. 1). Ultrasound scan confirmed the presence of 2 palpable and soft masses located in muscles, which were approximately 19 × 8 × 15 mm³ under the right chest wall and 26 × 10 × 19 mm³ under the right midaxillary line (Fig. 2). A subsequent biopsy of the mass (Fig. 3) showed larval-like tissue, peripheral fibrous tissue hyperplasia, lymphocytic infiltration and hyaline degeneration. The patient was probably diagnosed with taeniasis and cysticercosis and treated with oral albendazole (400 mg, twice daily) over 2 weeks. Hydrocortisone 50 mg was provided 2 days after the first albendazole treatment to counteract any potential negative effects on the central nervous system. After only 2 days of this antiparasitic treatment, the adult tapeworm was eliminated through the patient's feces (Fig. 3). When compared with the first MRI (half a month before antiparasitic treatment), the second MRI (half a month after antiparasitic treatment) demonstrated a slightly smaller focus (Fig. 1). The headache and vomiting resolved, and the patient remained symptom free over a 3-month follow-up period.

Morphological observation

The adult tapeworm was milky white, flat and long like a belt, thin and translucent, and the main segments are shown in Fig. 3. The uterine branches in the gravid proglottids were regular, and each side had approximately 15–30 branches (Fig. 3). The slide of the cyst showed no obvious typical features; however, some presumed deciduous hooks were found in the same slide, leading to indistinguishable morphology (Fig. 3).

Molecular identification

The proglottid from the patient was successfully amplified and sequenced using 2 pairs of the primers of the target COX1. The cyst was amplified with JB3 and JB4.5 primers, and the nuclear DNA HDP2 target fragment was also effectively amplified and sequenced. The 827-bp amplified products of proglottid using the *T. saginata* species-specific forward primer Tsag_COX1/F and *Taenia* common reverse primer Tae_COX1/R were 99.27–100% identical to *T. saginata* and 96.25–96.37% equivalent to *T. asiatica* without other species (Fig. 4). BLAST results indicated that the per cent identity of the 396-bp PCR products of the proglottid and cyst samples, which had trimmed JB3 and JB4.5 primer sequences, was 99.24–100% identical with *T. saginata* (Fig. 4). However, the 599 bp of the HDP2 target sequence revealed 99.67–99.83% identity with *T. saginata*, 95.66–99.67% with *T. asiatica* and 92.99–97.33% with *T. solium* under the same BLAST conditions (Fig. 4). Additional information regarding sequence alignment is provided in the Supplementary material. Based on the COX1 phylogenetic analysis (Fig. 5), the case is most likely *T. saginata*, next *T. asiatica* and finally *T. solium*.




Fig. 1. MRI shows that the largest cystic nodule was located next to the left ventricular triangle: (A–C) half a month before antiparasitic treatment and (D–F) half a month after antiparasitic treatment.




Fig. 2. Ultrasound scan shows hypoechoic mass containing cysticercus with calcification in the muscles of (A, B) right anterior thoracic wall ($19 \times 8 \times 15 \text{ mm}^3$) and (C, D) right midaxillary line ($26 \times 10 \times 19 \text{ mm}^3$).




Fig. 3. Tapeworm materials from the patient: (A) subcutaneous mass; (B) cyst; (C) adult tapeworm; (D) unstained proglottids; (E) a stained proglottid and (F) cystic lesion with hydrochloric acid-carmin staining showing presumed deciduous hooks (black arrows).

Discussion

Taeniasis coexisting with cysticercosis in humans is uncommon, especially for *T. saginata* infection. New insights into the pathogenicity and life cycle of *T. saginata* may be drawn from the case of the Tibetan patient with *Taenia* intestinal infection and cysticercosis (brain, muscles) reported in this paper. The life cycle of *Taenia*, involving the pathogenic mechanism of *Taenia* infection, is significant for ascertaining the disease classification (taeniasis or cysticercosis), epidemiological context, disease control and prevention (Garcia *et al.*, 2020). Current medical evidence confirmed the proven diagnosis of human taeniasis caused by *T. saginata* and suggested the diagnosis of muscular cysticercosis and neurocysticercosis caused by the larval stage of uncertain *Taenia* species, which may belong to *T. saginata*, *T. asiatica* or hybrids of the 2 sibling species (Okamoto *et al.*, 2010; Yamane *et al.*, 2012). Studies have also reported that rare *Taenia* species (*Taenia crassiceps*, *Taenia serialis* and *Taenia martis*) also cause human cysticercosis, confirmed by molecular diagnostic tests (Tappe *et al.*, 2016; Mueller *et al.*, 2020). The potential pathogenic mechanism of the larval stage of these species might be similar to *T. solium*, wherein embryos hatch in the small intestine and then may invade and spread haematogenously to the brain or muscle and other tissues (Garcia *et al.*, 2020).

Three common *Taenia* species can be differentiated by examination of their morphological characteristics, such as the scolex, mature and gravid proglottids in the adult stage, and the scolex in the larval stage (Eom *et al.*, 2020). However, the differential diagnosis between them is challenging when their morphology is not so typically visible. As far as morphological characteristics were concerned, the gravid proglottids (Fig. 3) from this patient were the same as *T. saginata* or *T. asiatica*, not *T. solium*, while the morphological evaluation of cysticercosis, that showed the presence of detached hooks (Fig. 3), referred to *T. asiatica* and *T. solium* rather than *T. saginata*. However, the traditional

morphological taxonomy has some inherent limitations leading to possible false-species identification and can neglect cryptic or pseudocryptic species. That is the reason why integrated taxonomy approaches, wherein molecular studies combine detailed morphological information, are important in helping characterize pairing at the individual level resulting in the perfect characterization of cryptic biodiversity (Hebert *et al.*, 2003; Laakmann *et al.*, 2020).

The most common method for molecular identification of *Taenia* tapeworms has been PCR coupled with the sequencing of the amplified PCR product (Ale *et al.*, 2014). Other assays for the differential diagnosis of *Taenia* include multiplex PCR (Nunes *et al.*, 2003; Yamasaki *et al.*, 2004; Jeon *et al.*, 2009; Ng-Nguyen *et al.*, 2017), PCR coupled with restriction fragment length polymorphism analysis (Gonzalez *et al.*, 2004), random-amplified polymorphic DNA analysis (Jeon *et al.*, 2009), loop-mediated isothermal amplification (Nkouawa *et al.*, 2009) and matrix-assisted laser desorption ionization-time-of-flight mass spectrometry (Wendel *et al.*, 2021). Different target sequences of *Taenia* genomic DNA, including mitochondrial DNA (e.g. COX1, CYTb, valine tRNA, NADH, 12S rDNA gene) (Yamasaki *et al.*, 2002; Okamoto *et al.*, 2010; Roelfsema *et al.*, 2016), ribosomal DNA (i.e. 28S rRNA, 5.8S rRNA and ITS rRNA gene) (von Nickisch-Roseneck *et al.*, 1999; Praet *et al.*, 2013) and nuclear DNA (HDP1, HDP2, cathepsin L-like cysteine peptidase, Ag2 gene) (Gonzalez *et al.*, 2000; Gonzalez *et al.*, 2004; Flores *et al.*, 2018), have been used as markers in molecular diagnosis. In particular, the mitochondrial COX1 gene that has been used in the study is a universally accepted DNA barcoding marker for assessing the genetic variation and evolutionary biology of Metazoa, including helminth parasites (Hebert *et al.*, 2003; Galimberti *et al.*, 2012; Zhang *et al.*, 2014; Rostami *et al.*, 2015; Mioduchowska *et al.*, 2018; Eberle *et al.*, 2020). Molecular characterization in our study based on COX1 and HDP2 indicated that the cyst seemed to be more inclined to *T. saginata* or *T. asiatica*. The analysis of HDP2 sequences in cysticercus samples ruled out the possibility of *T.*




Fig. 4. Sequence alignment analysis of (A) COX1 based on Tsag_COX1/F and Tae_COX1/R primers, (B) COX1 based on JB3 and JB4.5 primers and (C) HDP2 sequence.




Fig. 5. Phylogenetic tree was constructed based on the COX1 sequences of *Taenia* spp. using the maximum-likelihood method.

solium, but failed to reliably differentiate *T. saginata* from *T. asiatica* or their hybrids (Okamoto et al., 2010). High homologies were found between *T. saginata* and *T. asiatica* on mitochondrial *COX1* gene, as shown by the phylogenetic dendrogram (Fig. 5) which is in line with a prior study showing a 4.6% difference between the whole mitochondrial genome sequence of *T. saginata* and *T. asiatica* (Jeon et al., 2007). However, the BLAST results of the cyst were inconsistent with its morphology, which encompassed dropping hooks in the slide (Fig. 3). These disagreements in morphology and molecular identification suggested the possibility of hybrids of *T. saginata* and *T. asiatica*.

For studies involving DNA sequences, whole-genome sequencing has become the standard method for collecting relevant data (Eberle et al., 2020). Unfortunately, due to the scarcity of high-quality cyst samples, we have not been able to sequence the entire cyst genome. However, our study suggested that patients with taeniasis accompanied by cysticercosis may need whole-genome analysis to further ascertain whether *T. saginata* is also responsible for causing human cysticercosis. Despite the widespread availability of molecular identification methods, often tested on proglottids' samples, the sensitivity of detection in cyst samples may be lowered due to DNA degradation within cysts (Figueiredo et al., 2019). Many researchers have reported various mitochondrial/nuclear gene discordances in specimens, indicating potential hybrids between *T. asiatica* and *T. saginata* (Nkouawa et al., 2009; Okamoto et al., 2010; Yamane et al., 2012; Ale et al., 2014). Therefore, a molecular sequence-based species characterization that relies on information encoded by a single gene of the mitochondrial genome can be deceptive. Hence, mitochondrial, ribosomal or nuclear marker-based multilocus sequence analysis of genetic heterogeneity within the *Taenia* spp. could be the more effective routine molecular identification in clinical laboratories (Pajuelo et al., 2015). Therefore, it would be interesting to develop an easy, simple-to-handle and highly sensitive, multilocus sequences-based molecular diagnostic method for specimen identification of *Taenia* for validation in clinical settings.

In conclusion, the cause of human cysticercosis by non-*solium* *Taenia* species, such as *T. saginata* or *T. asiatica*, remains unclear. Correct diagnosis can be aided by utilizing whole-genome analysis and molecular identification approaches that focus on multilocus sequences. In order to better understand the comprehensive epidemiology and total life cycle of *Taenia*, the limitations of the existing molecular diagnostic applications require more research for the correct diagnosis and prevention of the disease.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S003118202200169X>

Data availability. The data and materials information used and analyzed in the current study are available from the first author (Jie Hou) or corresponding author (Ying Ma) upon reasonable request.

Acknowledgements. We thank the patient and his family.

Author's contributions. Conceptualization: Jie Hou, Ying Ma and Junyan Qu; data curation: Jie Hou and Weilin Chen; formal analysis: Jie Hou, Weilin Chen and Rong Chen; investigation: Jie Hou, Weilin Chen and Chunlei He; methodology: Jie Hou; project administration: Ying Ma and Jie Hou; resources: Ying Ma and Junyan Qu; software: Jie Hou; supervision: Ying Ma and Junyan Qu; validation: Ying Ma and Junyan Qu; visualization: Jie Hou; writing – original draft: Jie Hou; writing – review and editing: Ying Ma and Junyan Qu.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of interest. The authors declare there are no conflicts of interest.

Ethical standards. The patient's oral informed consent was obtained by phone, and written informed consent was waived with the approval of the West China Hospital, Sichuan University Biomedical Ethics Committee (approval number: 2022-472) because we met the criteria for an informed consent waiver.

References

- Abuladze KI (1964) *Principles of Cestology*, Vol. IV. Moscow: Nauka, p. 530.
- Ale A, Victor B, Praet N, Gabriël S, Speybroeck N, Dorny P and Devleeschauwer B (2014) Epidemiology and genetic diversity of *Taenia asiatica*: a systematic review. *Parasites & Vectors* 7, 45.
- Alroy KA and Gilman RH (2020) Tapeworm infections. In Ryan Edward T., Hill David R., Solomon Tom, Aronson Naomi E. and Endy Timothy P. (eds), *Hunter's Tropical Medicine and Emerging Infectious Diseases*. London: Elsevier, pp. 932–940.
- Asenjo A and Bustamente E (1950) Die neurochirurgische behandlung der cysticerkose. *DMW-Deutsche Medizinische Wochenschrift* 75, 1180–1183.
- Bacigalupo JADAB A, (1956) La *Taenia suginata* puede producir cysticercus bovis en el hombre? *Prensa Medica Argentina* 43, 1052–1054.
- Bowles J and McManus DP (1994) Genetic characterization of the Asian *Taenia*, a newly described taeniid cestode of humans. *American Journal of Tropical Medicine and Hygiene* 50, 33–44.
- Bowles J, Blair D and McManus DP (1992) Genetic variants within the genus *Echinococcus* identified by mitochondrial DNA sequencing. *Molecular and Biochemical Parasitology* 54, 165–173.
- Del Brutto OH, Nash TE, White Jr AC, Rajshekhkar V, Wilkins PP, Singh G, Vasquez CM, Salgado P, Gilman RH and Garcia HH (2017) Revised diagnostic criteria for neurocysticercosis. *Journal of the Neurological Sciences* 372, 202–210.
- De Rivas D (1937) *Cysticercus bovis* in man. In Papers on helminthology published in communication of the 30 year jubileum of K. J. Skriabin.
- Eberle J, Ahrens D, Mayer C, Niehuis O and Misof B (2020) A plea for standardized nuclear markers in metazoan DNA taxonomy. *Trends in Ecology & Evolution* 35, 336–345.
- Eichenberger RM, Thomas LF, Gabriel S, Bobic B, Devleeschauwer B, Robertson LJ, Saratsis A, Torgerson PR, Braae UC, Dermauw V and Dorny P (2020) Epidemiology of *Taenia saginata* taeniosis/cysticercosis: a systematic review of the distribution in east, southeast and south Asia. *Parasites & Vectors* 13, 234.
- Eom KS, Rim HJ and Jeon HK (2020) *Taenia asiatica*: historical overview of taeniasis and cysticercosis with molecular characterization. *Advances in Parasitology* 108, 133–173.
- Figueiredo BNS, RA LI, Sato M, da Silva CF, Pereira-Junior RA, Chigusa Y, Kawai S and Sato MO (2019) Occurrence of bovine cysticercosis in two regions of the state of Tocantins-Brazil and the importance of pathogen identification. *Pathogens (Basel, Switzerland)* 8(2), 66. doi: 10.3390/pathogens8020066
- Flores MD, Gonzalez LM, Hurtado C, Motta YM, Dominguez-Hidalgo C, Merino FJ, Perteguer MJ and Garate T (2018) HDP2: a ribosomal DNA (NTS-ETS) sequence as a target for species-specific molecular diagnosis of intestinal taeniasis in humans. *Parasites & Vectors* 11, 117.
- Galimberti A, Romano DF, Genchi M, Paoloni D, Vercillo F, Bizzarri L, Sasseria D, Bandi C, Genchi C, Ragni B and Casiraghi M (2012) Integrative taxonomy at work: DNA barcoding of taeniids harboured by wild and domestic cats. *Molecular Ecology Resources* 12, 403–413.
- Garcia HH (2018) Neurocysticercosis. *Neurologic Clinics* 36, 851–864.
- Garcia HH, Gonzalez AE, Evans CA, Gilman RH and Cysticercosis Working Group in Peru (2003) *Taenia solium* cysticercosis. *Lancet (London, England)* 362, 547–556.
- Garcia HH, Gonzalez AE and Gilman RH (2020) *Taenia solium* cysticercosis and its impact in neurological disease. *Clinical Microbiology Reviews* 33(3), e00085–19. doi: 10.1128/CMR.00085-19
- Goldsmid J (1966) Two unusual cases of cysticercosis in man in Rhodesia. *Journal of Helminthology* 40, 331–336.
- Gonzalez LM, Montero E, Harrison LJ, Parkhouse RM and Garate T (2000) Differential diagnosis of *Taenia saginata* and *Taenia solium* infection by PCR. *Journal of Clinical Microbiology* 38, 737–744.
- Gonzalez LM, Montero E, Morakote N, Puente S, Diaz De Tuesta JL, Serra T, Lopez-Velez R, McManus DP, Harrison LJ, Parkhouse RM and Garate T (2004) Differential diagnosis of *Taenia saginata* and *Taenia saginata asiatica* taeniasis through PCR. *Diagnostic Microbiology and Infectious Disease* 49, 183–188.

- Hebert PD, Cywinska A, Ball SL and deWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings. Biological Sciences* **270**, 313–321.
- Ito A, Nakao M and Wandra T (2003) Human taeniasis and cysticercosis in Asia. *Lancet (London, England)* **362**, 1918–1920.
- Jansen F, Dorny P, Gabriël S, Dermauw V, Johansen MV and Trevisan C (2021) The survival and dispersal of *Taenia* eggs in the environment: what are the implications for transmission? A systematic review. *Parasites & Vectors* **14**, 88.
- Jeon HK, Kim KH and Eom KS (2007) Complete sequence of the mitochondrial genome of *Taenia saginata*: comparison with *T. solium* and *T. asiatica*. *Parasitology International* **56**, 243–246.
- Jeon HK, Chai JY, Kong Y, Waikagul J, Insisiengmay B, Rim HJ and Eom KS (2009) Differential diagnosis of *Taenia asiatica* using multiplex PCR. *Experimental Parasitology* **121**, 151–156.
- Johnson M, Zaretskaya I, Raytselis Y, Merezukh Y, McGinnis S and Madden TL (2008) NCBI BLAST: a better web interface. *Nucleic Acids Research* **36**, W5–W9.
- Laakmann S, Blanco-Bercial L and Cornils A (2020) The crossover from microscopy to genes in marine diversity: from species to assemblages in marine pelagic copepods. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **375**, 20190446.
- Mioduchowska M, Czyz MJ, Goldyn B, Kur J and Sell J (2018) Instances of erroneous DNA barcoding of metazoan invertebrates: are universal cox1 gene primers too 'universal'? *PLoS ONE* **13**, e0199609.
- Mueller A, Förch G, Zustin J, Muntau B, Schuldt G and Tappe D (2020) Case report: molecular identification of larval *Taenia martis* infection in the pouch of Douglas. *American Journal of Tropical Medicine and Hygiene* **103**, 2315–2317.
- Ng-Nguyen D, Stevenson MA, Dorny P, Gabriel S, Vo TV, Nguyen VT, Phan TV, Hii SF and Traub RJ (2017) Comparison of a new multiplex real-time PCR with the Kato Katz thick smear and copro-antigen ELISA for the detection and differentiation of *Taenia* spp. in human stools. *PLoS Neglected Tropical Diseases* **11**, e0005743.
- Niño FL (1950) Cisticercosis humana en la Republica Argentina. Estudio de ma nueva observación. *Prensa Medica Argentina* **37**, 3040–3044.
- Nkouawa A, Sako Y, Nakao M, Nakaya K and Ito A (2009) Loop-mediated isothermal amplification method for differentiation and rapid detection of *Taenia* species. *Journal of Clinical Microbiology* **47**, 168–174.
- Nunes CM, Lima LG, Manoel CS, Pereira RN, Nakano MM and Garcia JF (2003) *Taenia saginata*: polymerase chain reaction for taeniasis diagnosis in human fecal samples. *Experimental Parasitology* **104**, 67–69.
- Okamoto M, Nakao M, Blair D, Anantaphruti MT, Waikagul J and Ito A (2010) Evidence of hybridization between *Taenia saginata* and *Taenia asiatica*. *Parasitology International* **59**, 70–74.
- Pajuelo MJ, Eguiluz M, Dahlstrom E, Requena D, Guzman F, Ramirez M, Sheen P, Frace M, Sammons S, Cama V, Anzick S, Bruno D, Mahanty S, Wilkins P, Nash T, Gonzalez A, Garcia HH, Gilman RH, Porcella S, Zimic M and Cysticercosis Working Group in Peru (2015) Identification and characterization of microsatellite markers derived from the whole genome analysis of *Taenia solium*. *PLoS Neglected Tropical Diseases* **9**, e0004316.
- Pawlowski Z and Schultz MG (1972) Taeniasis and cysticercosis (*Taenia saginata*). *Advances in Parasitology* **10**, 269–343.
- Praet N, Verweij JJ, Mwape KE, Phiri IK, Muma JB, Zulu G, van Lieshout L, Rodriguez-Hidalgo R, Benitez-Ortiz W, Dorny P and Gabriel S (2013) Bayesian modelling to estimate the test characteristics of coprology, coproantigen ELISA and a novel real-time PCR for the diagnosis of taeniasis. *Tropical Medicine & International Health* **18**, 608–614.
- Qian MB, Xiao N, Li SZ, Abela-Ridder B, Carabin H, Fahrion AS, Engels D and Zhou XN (2020) Control of taeniasis and cysticercosis in China. *Advances in Parasitology* **110**, 289–317.
- Roelfsema JH, Nozari N, Pinelli E and Kortbeek LM (2016) Novel PCRs for differential diagnosis of cestodes. *Experimental Parasitology* **161**, 20–26.
- Rostami S, Salavati R, Beech RN, Babaei Z, Sharbatkhorri M and Harandi MF (2015) Genetic variability of *Taenia saginata* inferred from mitochondrial DNA sequences. *Parasitology Research* **114**, 1365–1376.
- Tanasescu I and Repciuc E (1939) Ein fall von cysticercus bovis im unterhautgewebe des Menschen. *Virchows Archiv für Pathologische Anatomie und Physiologie und für klinische Medizin* **304**, 555–558.
- Tappe D, Berkholtz J, Mahlke U, Lobeck H, Nagel T, Haeupler A, Muntau B, Racz P and Poppert S (2016) Molecular identification of zoonotic tissue-invasive tapeworm larvae other than *Taenia solium* in suspected human cysticercosis cases. *Journal of Clinical Microbiology* **54**, 172–174.
- von Nickisch-Rosenegk M, Silva-Gonzalez R and Lucius R (1999) Modification of universal 12S rDNA primers for specific amplification of contaminated *Taenia* spp. (Cestoda) gDNA enabling phylogenetic studies. *Parasitology Research* **85**, 819–825.
- Wendel TP, Feucherolles M, Rehner J, Poppert S, Utzinger J, Becker SL and Sy I (2021) Evaluating different storage media for identification of *Taenia saginata* proglottids using MALDI-TOF mass spectrometry. *Microorganisms* **9**(1), 82. doi: 10.3390/microorganisms9102006
- Yamane K, Suzuki Y, Tachi E, Li T, Chen X, Nakao M, Nkouawa A, Yanagida T, Sako Y, Ito A, Sato H and Okamoto M (2012) Recent hybridization between *Taenia asiatica* and *Taenia saginata*. *Parasitology International* **61**, 351–355.
- Yamasaki H, Nakao M, Sako Y, Nakaya K, Sato MO, Mamuti W, Okamoto M and Ito A (2002) DNA differential diagnosis of human taeniid cestodes by base excision sequence scanning thymine-base reader analysis with mitochondrial genes. *Journal of Clinical Microbiology* **40**, 3818–3821.
- Yamasaki H, Allan JC, Sato MO, Nakao M, Sako Y, Nakaya K, Qiu D, Mamuti W, Craig PS and Ito A (2004) DNA differential diagnosis of taeniasis and cysticercosis by multiplex PCR. *Journal of Clinical Microbiology* **42**, 548–553.
- Zhang G, Chen J, Yang Y, Liu N, Jiang W, Gu S, Wang X and Wang Z (2014) Utility of DNA barcoding in distinguishing species of the family Taeniidae. *Journal of Parasitology* **100**, 542–546.