

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1240>

#### EV0911

### A Study of empathy in Romanian general nursing students

F. Romosan\*, R.S. Romosan, A.M. Romosan  
"Victor Babes" University of Medicine and Pharmacy, Neuroscience,  
Timisoara, Romania

\* Corresponding author.

**Introduction** Empathy is of most importance in the medical field. The ability to comprehend and connect with the emotional state of another person is essential for establishing a successful interaction between patients and health care professionals.

**Objectives** The aim of this study was to assess empathic response in general nursing (GN) students.

**Methods** The study was conducted between 2015–2016 on 75, sixth-year Romanian GN undergraduates from the Timisoara "Victor Babes" university of medicine and pharmacy. To evaluate the level of empathy, we used the empathy quotient (EQ), a 60-item self-report inventory with 40 questions empathy-related and 20 filler questions.

**Results** We included in this study 12 (16%) males and 63 (84%) females, with a mean age of 23.48 years (SD = 2.17). Female GN students had significantly higher EQ mean scores than the general female population. Compared to female students, males obtained significantly lower EQ mean scores. There were no significant differences between male and female students regarding age distribution.

**Conclusions** Female GN students showed greater empathy than their male colleagues, as well as than the general female population. These results support the theory on women's understanding of others mental states, vital to the provision of a worthy nursing care.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1241>

#### EV0912

### Cognitive emotional regulation in Romanian general medicine students

R.S. Romosan\*, A.M. Romosan, V.R. Enatescu, I. Papava,  
C. Giurgi-Oncu

"Victor Babes" University of Medicine and Pharmacy, Neuroscience,  
Timisoara, Romania

\* Corresponding author.

**Introduction** During medical school, students experience significant amounts of stress. Since certain emotion regulatory strategies are known to be maladaptive, the way in which students are capable to regulate their emotions becomes very important, because it can affect their physical and mental welfare.

**Objectives** The purpose of this study was to assess cognitive emotion regulation strategies in Romanian general medicine (GM) students.

**Methods** The study was conducted between 2015–2016 on 86 sixth-year Romanian GM undergraduates from the Timisoara "Victor Babes" university of medicine and pharmacy. In order to identify the cognitive emotion regulation strategies (or cognitive coping strategies) that students use after experiencing negative situations or life events we used the Romanian version of the cognitive emotion regulation questionnaire (CERQ).

**Results** The sample consisted of 30 (34.9%) males and 56 (65.1%) females, with ages ranging between 24 and 31 years (mean age = 24.97 years, SD = 1.74). Compared to female students, males obtained significantly lower mean scores in "umination" ( $t = -2.84$ ,

$P = 0.005$ , 95% CI = -1.64; -0.29), "positive refocusing" ( $t = -2.09$ ,  $P = 0.037$ , 95% CI = -1.42; -0.04) and "catastrophizing" ( $t = -3.17$ ,  $P = 0.002$ , 95% CI = -1.31; -0.3). Both male and female GM students had significantly higher mean scores in "blaming others" than their respectively gender-related general population.

**Conclusions** Results of this study suggest that GM students, when facing stressful or negative events, are more inclined in using "blaming others" as a coping strategy. Female students seem to be more inclined than males to use "rumination", "catastrophizing" and "positive refocusing" as cognitive coping mechanisms.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1242>

#### EV0913

### Antipsychotic-induced hyperprolactinemia

S. Khouadja\*, R. Ben Soussia, A. Bouallagui, I. Marrag, S. Younes,  
M. Nasr

University Hospital, Psychiatry, Mahdia, Tunisia

\* Corresponding author.

**Introduction** As antipsychotic agents are increasingly used, many patients are at risk for antipsychotic-induced hyperprolactinemia.

**Aims of the study** Estimate the prevalence of hyperprolactinemia in patients treated by a single antipsychotic and identify the risk factors for its occurrence.

**Methods** This is a prospective study carried out at the psychiatric department of psychiatry of university hospital of Mahdia during 24 months. We have included all patients with a follow up and treated by single antipsychotic for at least 12 weeks. A pituitary MRI has been requested for patients with a prolactin level higher than 100 ng/ml.

**Results** We have collected 92 patients. Hyperprolactinemia was found in 34.8% of patients among which 7.6% have had prolactin level greater than 150 ng/ml. Pituitary MRI have revealed 2 cases of macro-adenoma. The decrease of the antipsychotic doses has significantly improved prolactin levels. The switch of antipsychotic with another less inducing hyperprolactinemia has significantly decreased prolactin levels. 7 factors were correlated significantly to hyperprolactinemia: sex (female), substance use, presence of side effects, combination of psychotropic drugs, atypical antipsychotics, type of antipsychotic: Haloperidol and amisulpride, antipsychotic dose greater than 1000 mg Chlorpromazine equivalent.

**Conclusion** Hyperprolactinemia must be carefully identified, through a pre-therapeutic assessment and monitoring of patients.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1243>

#### EV0914

### Hyperinsulinism as evidence of munchausen syndrome by proxy: A case report

#### A case report

S. Khouadja\*, S. Younes, S. Fathallah, R. Ben Soussia, L. Zarrouk,  
M. Nasr

University Hospital, Psychiatry, Mahdia, Tunisia

\* Corresponding author.

**Introduction** Hyperinsulinism is one of the most important causes of hypoglycemia. Rarely, drug toxicity can be a reason. In the context of Munchausen syndrome by proxy (MSBP), toxicity usually occurs in children due to drug administration by a parent or caregiver.

**Methods** we report a case of a 2-year-old girl with hyperinsulinemic hypoglycaemia due to insulin injections by her mother.