

Results: The E-sport intervention consists of two fields “in-game” and “over-game” In-game concerns the practice of gaming Over-game concerns the interventions organization Power in each field is driven by specific values and access to certain competencies Power in-game was equally open to patients and professionals leading to symmetric power relations Power over-game was open to professionals only leading to asymmetrical power relations Professionals may allow power distribution to patients during gameplay, while still retaining the overall power over the intervention

Conclusions: It is possible to balance between care-and-custody in forensic psychiatry. This study provides important insights to guide further practice.

Keywords: forensic psychiatry; qualitative study; care-custody; relations

EPP0686

Dangerousness assessment in psychiatric inpatients suffering from psychotic disorders

S. Khoudja^{1,2*}, R. Melki², C. Ben Taleb² and L. Zarrouk²

¹Psychiatry, University Hospital Of Mahdia., Mahdia, Tunisia and

²Psychiatry, University Hospital Of Mahdia, Mahdia, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1012

Introduction: Dangerousness is a state in which a person is likely to commit violent acts.

Objectives: Describe the socio-demographic and clinical characteristics of psychiatric inpatients hospitalized in the locked unit and suffering from schizophrenia or other psychotic disorders and to assess their dangerousness.

Methods: This is a cross-sectional study carried out in the locked unit of psychiatric department of the University Hospital of Mahdia during one year. We have collected data of patients diagnosed with schizophrenia or other psychotic disorders according to DSM 5. Psychometric assessment was done using the BPRS, the PANSS, the VRAG and the HCR-20 scales.

Results: We have included 173 patients. The average age was 36 years with a sex ratio of 9. The majority of our patients were unmarried and of a low economic level. Alcohol and cannabis consumption was found in 7.6% and in 5.7% of cases respectively. A history of incarceration was found in 79% of cases. Homicide was the infraction the most committed in 8% of cases. 71.2% of patients had an anterior hospitalization in the locked unit. Aggressiveness and instability were the main indication for hospitalization. The diagnosis was schizophrenia in 84% of cases. Patients were treated with classic antipsychotic drugs in 55.8% of cases. Non-adherence to treatment was reported in 33% of cases. The average score of psychometric scales were BPRS = 21.4; VRAG = - 4.87 and HCR-20 = 17± 0.87.

Conclusions: Our study showed comparable assessments for dangerousness with the literature. Evaluating dangerousness should represent the first step of the therapeutic process.

Keywords: psychiatry; dangerousness; schizophrénia; Psychostic disorders

EPP0687

Agresion and violent behaviour risk assessment

G. Marinho^{1*} and S. Almeida²

¹Clinica 6, CHPL, Lisbon, Portugal and ²Serviço De Psiquiatria, Hospital Prisional S. João Deus, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1013

Introduction: After discharge from forensic psychiatric hospital, rates of violent reoffending are reported to range from 2% to 8% per year in high income countries. Risk assessment informs decisions around admission to and discharge from secure psychiatric hospital and contributes to treatment and supervision Current approaches to assess violence risk in secure hospitals are resource intensive, limited by accuracy and authorship bias. Forensic Violence Oxford (FoVOx) was developed using all forensic psychiatric patients in Sweden, based on the largest forensic psychiatric sample to date, and has the advantage of using routinely available data, which are less liable to bias than interview-based measures.

Objectives: Literature review on the Forensic Psychiatry and Violence Oxford (FoVOx) tool.

Methods: Pubmed and Google Scholar search

Results: The 12 items within the FoVOx tool are sex, age, previous violent crime, previous serious violent crime, primary discharge diagnosis, drug use disorder at point of hospitalization or discharge, any lifetime drug use disorder, alcohol use disorder at point of hospitalization or discharge, personality disorder at discharge, employment at admission, five or more prior inpatient episodes, and whether current length of stay has exceeded one year.

Conclusions: The FoVOx tool is scalable, quick, free to use and available online. Its use could enable clinicians to provide a reasonably accurate risk assessment in a brief and cost-effective way, and free up time to focus on clinical care and risk management rather than risk assessment.

Keywords: violence; risk assessment; forensic psychiatry

EPP0688

What can we learn about perpetrators of domestic and familial homicide and their involvement with mental health services from domestic homicide reviews?

P. Macinnes

Ioppn, King's College London, London, United Kingdom

doi: 10.1192/j.eurpsy.2021.1014

Introduction: The evidence that supports an association between domestic violence and abuse (DVA) perpetration and mental disorders is increasing. Since 2011, authorities in England and Wales have been required to conduct Domestic Homicide Reviews (DHRs) into deaths caused by violence, abuse or neglect of individuals aged 16 or over, by a family member or a current or ex-partner.

Objectives: The aim of the study is to describe the characteristics of perpetrators of domestic homicide in a sample of DHR reports in which the perpetrator was known to mental health services in the 12 months before the offence. This sample will undergo qualitative framework analysis as part of another study conducted by the authors.

Methods: The researchers compiled a list of DHRs available online and randomly sampled 168 reports; in 20 of those reports, the perpetrators were under the care of mental health services in the 12 months prior to the offence. We have applied descriptive statistics to report on the sample characteristics.

Results: The common mental illnesses diagnosed amongst perpetrators were depression (20%), anxiety (15%) and schizophrenia

(15%). 25% of the perpetrators didn't have a psychiatric diagnosis and 70% had a history of self-harm and suicidal ideation. Substance use was prevalent (60%). Half of the homicides involved children.

Conclusions: This report describes the demographic and mental health characteristics of a sample of perpetrators of domestic homicides. Further research is needed into the patterns of mental health service use by DVA perpetrators in order to improve identification and risk management.

Conflict of interest: No significant relationships.

EPP0691

Elderly criminal patients in Razi psychiatric hospital of tunis

H. Jemli¹, R. Jomli^{2,3*}, H. Rym³, I. Bouslama¹, A. Ben Cheikh Ahmed¹ and F. Nacef²

¹Psychiatry Department A, Razi Hospital, manouba, Tunisia;

²Psychiatry A Department, Razi Hospital, Manouba, Tunisia and

³Avicenne, manouba psychiatry, manouba, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1015

Introduction: Criminality has become of increasing concern in the practice of psychiatry. However, violence among elderly psychiatric patients is an underestimated and understudied phenomenon.

Objectives: The aim of the study is to identify differences in the socio-demographic, clinical and criminological profiles between elderly criminals under treatment for psychiatric disorders and those not known to have mental disorders prior to the criminal offense in Tunisia.

Methods: We present a retrospective study on twenty male criminal mental patients, aged sixty years or older, who were hospitalized in the Forensic Psychiatry Department of Razi Hospital during 18 years, following a dismissal for insanity under Article 38 of the Penal Code and Article 29 of Law 92/83 on Mental Health.

Results: Prevalence was higher among elderly criminals without a known psychiatric history (2.42% versus 1.98%). The average age was roughly the same, around 73 years old. Neurological and cardiovascular histories were the most common in both groups. The criminal act was indicative of dementia in 8 cases. Criminal history was more frequent in elderly patients with a personal psychiatric history (55.5% versus 18.2%). Patients whose act was revelatory of their mental disorder committed more violent crimes (63.7% versus 44.4%) using blunt objects (71.4% versus 0%). The victim most often belonged to the aggressor's family, particularly the spouse (87.5%).

Conclusions: Screening for criminal risk factors in the elderly, early diagnosis of mental disorders and a comprehensive therapeutic project are necessary to prevent the risk of violent behaviour.

Keywords: Elderly; forensic psychiatry; criminality

EPP0692

Tidier. e-sport; a recovery oriented intervention in forensic psychiatry.

L. Sørensen^{1,2*}, H. Kennedy^{3,4}, B. Jensen², M. Terkildsen², R. Poulsen², M. Josefsen² and A. Di Lieto²

¹Department Of Clinical Medicine, Aarhus University- Health, Aarhus N, Denmark; ²Department Of Forensic Psychiatry, Aarhus University Hospital Psychiatry, Aarhus N, Denmark; ³Department Of Psychiatry, Trinity College- Dublin University, Dublin, Ireland and ⁴National Forensic Mental Health Service, Central Mental Hospital Dundrum, Dundrum, Ireland

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1016

Introduction: Recently video gaming, have attracted considerable attention for its possible beneficial therapeutic effects, the possibility for testing behavior in safe artificial environments and as a tool for professionals and patients to build specific competencies for the everyday life. Also, a substantial amount of research suggests that videogaming might improve the participants social and cognitive skills and emotional regulation. There is little or no evidence that videogaming increases long term aggression or leads to physical aggression. At a medium secure forensic psychiatric in-patient ward, the patients and staff engage in weekly E – Sport sessions (primarily counterstrike) to further the recovery process.

Objectives: To provide a standardized description of how E-sport is organized and used in the recovery process among forensic psychiatric patients.

Methods: The Template for Intervention Description and Replication (TIDieR) checklist and guide is widely used to in health research to describe interventions in clinical trials and other health research contexts. By use of TIDieR we describe a newly developed E-sport intervention, in which staff members and patients in a medium secure forensic psychiatric ward engage in weekly E-Sport sessions (primarily counterstrike) to improve patient-staff relationship.

Results: The E-sport intervention is detailed by use of the 12 TIDieR items and practical experiences and insights will be described.

Conclusions: This standardized and detailed description of how is used in a recovery-oriented process in forensic psychiatry can be used for future studies that wishes to implement the intervention or for research studies replicating the treatment.

Conflict of interest: No significant relationships.

EPP0693

Factors related to the dangerousness of psychiatric inpatients

M. Kacem^{1*}, S. Khoudja², S. Brahim³, A. Chaouch² and L. Zarrouk⁴

¹Department Of Psychiatry, University hospital of mahdia, chebba, Tunisia; ²Psychiatry, University Hospital Of Mahdia, Mahdia, Tunisia;

³Department Of Psychiatry, University Hospital of Mahdia, chebba, Tunisia and ⁴Department Of Psychiatry, University Hospital Of Mahdia, Tunisia., Psychiatry, Mahdia, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1017

Introduction: Mental illness may explain some acting outs, but it does not necessarily lead to a dangerous attitude.

Objectives: Describe the socio-demographic, clinical and therapeutic characteristics of patients considered dangerous and to identify the determinants of psychiatric dangerousness.