

Dietary fish oil and DHA down-regulate antigen-activated CD4⁺ T-cells while promoting the formation of liquid-ordered mesodomains

Wooki Kim^{1*}, Rola Barhoumi², David N. McMurray^{3,4} and Robert S. Chapkin^{2,3,4*}

¹Department of Food Science and Biotechnology, Kyung Hee University, Seocheon-dong, Gibeunggu, Yonginsu, Gyeonggi-do, Republic of Korea

²Department of Veterinary Integrative Biosciences, Texas A&M University, College Station, TX 77843-4458, USA

³Program in Integrative Nutrition and Complex Diseases, 321 Kleberg Center, MS 2253, Texas A&M University, College Station, TX 77843-2253, USA

⁴Department of Microbial and Molecular Pathogenesis, Texas A&M University System Health Science Center, College Station, TX, USA

(Submitted 16 October 2012 – Final revision received 2 July 2013 – Accepted 3 July 2013 – First published online 20 August 2013)

Abstract

We have demonstrated previously that *n*-3 PUFA endogenously produced by *fat-1* transgenic mice regulate CD4⁺ T-cell function by affecting the formation of lipid rafts, liquid-ordered mesodomains in the plasma membrane. In the present study, we tested the effects of dietary sources of *n*-3 PUFA, i.e. fish oil (FO) or purified DHA, when compared with an *n*-6 PUFA-enriched maize oil control diet in DO11.10 T-cell receptor transgenic mice. Dietary *n*-3 PUFA were enriched in CD4⁺ T-cells, resulting in the increase of the *n*-3:*n*-6 ratio. Following antigen-specific CD4⁺ T-cell activation by B-lymphoma cells pulsed with the ovalbumin 323–339 peptide, the formation of liquid-ordered mesodomains at the immunological synapse relative to the whole CD4⁺ T-cell, as assessed by Laurdan labelling, was increased ($P < 0.05$) in the FO-fed group. The FO diet also suppressed ($P < 0.05$) the co-localisation of PKC θ with ganglioside GM1 (monosialotetrahexosylganglioside), a marker for lipid rafts, which is consistent with previous observations. In contrast, the DHA diet down-regulated ($P < 0.05$) PKC θ signalling by moderately affecting the membrane liquid order at the immunological synapse, suggesting the potential contribution of the other major *n*-3 PUFA components of FO, including EPA.

Key words: *n*-3 PUFA: DHA: Fish oil: Lipid rafts: CD4⁺ T-cells

Immune responses are required for host protection against infections and neoplasms. However, in order to prevent immune-related disorders, such as autoimmune and chronic inflammatory diseases, the balance between the activation and suppression of immune responses is regulated very tightly by complex immune cell types in higher organisms. Among those immune cell types, CD4⁺ helper T-cells play a pivotal role in regulating chronic inflammation in various disease models, including, but not limited to, chronic colitis^(1,2), atherosclerosis⁽³⁾, rheumatoid arthritis^(4,5) and chronic eosinophilic lung inflammation⁽⁶⁾.

Immune functions are affected by the diet. Dietary fish oil (FO) enriched with *n*-3 PUFA exhibits anti-inflammatory properties in both *in vitro* and *in vivo* experimental models^(7,8). DHA (22:6 Δ 4,7,10,13,16,19) and EPA (20:5 Δ 5,8,11,14,17) are the major *n*-3 PUFA found in dietary fish oil⁽⁹⁾, which have been studied for their immunoregulatory properties.

Furthermore, clinical and epidemiological studies have supported the favourable role of *n*-3 PUFA on T-cell-mediated chronic inflammatory diseases^(10,11). These immunomodulatory effects of *n*-3 PUFA appear to be mediated by a variety of mechanisms, including the inhibition of pro-inflammatory eicosanoid production, the antagonism of NF- κ B and the modulation of lipid rafts (reviewed previously in Kim *et al.*^(12,13)). Our laboratory has also demonstrated that *n*-3 PUFA decrease the accumulation of CD4⁺ T-cells in the colonic lamina propria during dextran sodium sulphate-induced colitis, thereby suppressing colitis-associated colon cancer^(14,15).

Previous studies have focused on CD4⁺ T-cells activated polyclonally by either concanavalin A, monoclonal antibodies specific to CD3 and CD28, or phorbol ester (phorbol-12-myristate-13-acetate) and ionophore (ionomycin). However, since these treatments potentially activate T-cells by multidirectional signalling

Abbreviations: AA, arachidonic acid; ALA, α -linoleic acid; FO, fish oil diet; GM1, monosialotetrahexosylganglioside; GP, generalised polarisation; IS, immunological synapse; MO, maize oil diet; OVA, ovalbumin 323–339.

* **Corresponding authors:** W. Kim, email kimw@khu.ac.kr; R. S. Chapkin, email r-chapkin@tamu.edu

on the cell surface, their interpretation is complicated. In order to mimic antigen-induced clustering of proteins on the T-cell surface, researchers have used superantigens and cholera toxin B subunit, which also have limitations since they do not accurately reflect the response of T-cells to nominal antigens. In contrast, antigen-specific T-cell activation, in which polar immunological synapses (IS) to antigen-presenting cells are formed, is more physiological. Antigen-specific T-cell receptor transgenic mice of the DO11.10 strain harbour CD4⁺ T-cells, which respond to the ovalbumin 323–339 (OVA) epitope. Due to the genetic deletion of recombination activating gene 2 (*RAG2*), the antigen specificity of those transgenic T-cells are maintained during positive and negative selection in the thymus⁽¹⁶⁾.

Heterogeneous liquid-ordered mesodomains of the plasma membrane, i.e. lipid rafts, effectively sort signalling proteins during CD4⁺ T-cell antigen receptor activation according to their affinity for raft domains, leading to the assembly of stable, larger rafts at the contact site of T-cells/antigen-presenting cells, i.e. IS^(17,18). Interestingly, lipid raft size appears to be tightly regulated, since oversized lipid raft domains negatively affect T-cell signalling⁽¹⁹⁾. Using *fat-1* transgenic mice in which endogenous *n*-3 PUFA are generated and targeted to the T-cell membrane^(20,21), we have demonstrated that *n*-3 PUFA modulate lipid raft composition and down-regulate CD4⁺ T-cell signalling⁽²²⁾. The *fat-1* transgenic mouse model has an advantage in that fatty acid loss to the culture medium during the functional *ex vivo* assessment of CD4⁺ T-cells is prevented⁽²¹⁾. However, it remains to be determined whether dietary supplementation with *n*-3 PUFA can recapitulate these findings. Therefore, in the present study, we fed DO11.10-Rag2^{-/-} T-cell receptor transgenic mice a semi-purified diet containing either fish oil (enriched with DHA and EPA), purified DHA alone, or control maize oil (devoid of *n*-3 PUFA) at physiologically relevant levels⁽²³⁾. Here we report the effects of dietary FO and DHA on antigen-induced lipid raft formation and subsequent T-cell signalling.

Materials and methods

Mice and diets

All experimental procedures using laboratory animals were approved by the Texas A&M University Laboratory Animal Care and Use Committee. DO11.10 Rag2^{-/-} T-cell receptor transgenic mice were purchased from Taconic Farms, bred and maintained at Texas A&M University. At the onset of the study, sex- and age (8–16 weeks)-matched mice were fed a semi-purified control diet containing 5% maize oil (MO) (Dyets Company) by weight during the 7 d acclimatisation period, followed by a 2-week feeding period with either the same MO control diet, a FO diet (1% MO + 4% FO; Omega Protein Corporation) or a DHA diet (4.05% MO + 0.95% DHASCO[®] (44.9% purity); Martek). All the diets met the Nutrition Research Council's nutrition requirements⁽²⁴⁾ containing identical levels of major (fat 5%, protein 20%, carbohydrate 42%, starch 22% and fibre 6%) and minor (vitamins 3.5%, minerals 1%, methionine 0.35% and choline chloride 0.2%) ingredients and differed only in the sources

of fat as described above. The diets were stored at –20°C and provided fresh daily in order to avoid oxidation. Food and drinking-water were provided *ad libitum*.

CD4⁺ T-cell purification and Laurdan labelling

Mice were euthanised by CO₂ asphyxiation. Spleen was removed and placed in complete Roswell Park Memorial Institute (RPMI)-1640 medium with 25 mM-HEPES (Irvine Scientific), supplemented with 5% fetal bovine serum (FBS; HyClone), 10⁵ U penicillin/l (60 mg/l) and 100 mg streptomycin/l (Irvine Scientific), 2 mM-L-glutamine (Gibco) and 10 μM-2-mercaptoethanol (Sigma). CD4⁺ T-cells (>90% pure as determined by surface staining of CD3 and CD4, data not presented) were isolated from the spleen by a magnetic microbead positive selection method (Miltenyi Biotec) according to the manufacturer's recommendations⁽²²⁾. Purified CD4⁺ T-cells were labelled with Laurdan (Invitrogen) for lipid raft visualisation as described previously⁽²²⁾. Briefly, 5 μmol/l Laurdan was prepared in serum-free RPMI medium and incubated with 2 × 10⁶ cells/ml for 30 min at 37°C. The cells were subsequently washed and resuspended in serum-free Leibovitz's medium (Irvine Scientific).

Analysis of fatty acid composition

Total lipids were extracted from the diets or CD4⁺ T-cells by the method of Folch *et al.*⁽²⁵⁾. Briefly, lipids were extracted by the mixture of chloroform and methanol (2:1, v/v), followed by methylation with 6% (v/v) hydrogen chloride in methanol. Fatty acid methyl esters were extracted using hexane and 0.1 M-potassium chloride and analysed by capillary GC (Fig. 1) as described previously⁽²⁶⁾.

Antigen-specific stimulation and lipid raft assessment

For antigen-specific activation, A-20 B-lymphoma cells (ATCC) at a logarithmic growth phase were incubated with 25 μg/ml of OVA peptide (Quality Controlled Biochemicals, custom synthesised, peptide sequence: ISQAVHAAHAEINEAGR) for 4 h at 37°C in complete RPMI medium. After incubation, excess OVA was removed by washing with PBS, and OVA-pulsed B cells and Laurdan-labelled CD4⁺ T-cells were mixed at a 1:2 ratio. Cell mixtures were seeded onto poly-L-lysine-precoated chambered cover glass slides (2 × 10⁶ CD4⁺ T-cells/chamber; Sigma-Aldrich) in serum-free Leibovitz's medium. After 30 min of incubation at 37°C, Laurdan-labelled plasma membranes were visualised by two-photon microscopy (Zeiss LSM 510 META NLO) with a 40 × objective 1.3 numerical aperture (NA) oil at wavelengths of 400–460 and 470–530 nm. The Coherent Chameleon femtosecond-pulsed Ti:Sapphire laser was set at an excitation wavelength of 770 nm. All images were converted to 8-bit/channel TIFF format and were processed using Adobe Photoshop CS3[®]. The mean intensity of each colour channel was measured in areas of interest either at the contact regions of the IS by drawing an oval at the T-cell membrane proximal to the B-lymphoma cell or on the same whole CD4⁺ T-cell by drawing a polygon around




Fig. 1. (A) Formation of liquid-ordered mesodomains was quantified as general polarisation (GP) values at the immunological synapse (■) or the whole CD4⁺ T-cell (□). DO11.10 T-cell receptor transgenic CD4⁺ T-cells were co-cultured with ovalbumin-pulsed B-lymphoma cells (ten to twelve cells per mouse from four mice per diet were examined to obtain a total of forty to forty-eight observations). Values are means, with standard errors represented by vertical bars. *Mean value was significantly different from that of the whole cell ($P < 0.05$). (B) Increase in membrane liquid order at the immunological synapse relative to the whole CD4⁺ T cell was assessed by calculating the fold increase of GP values. Fold increase was expressed as $GP_{\text{immunological synapse}}/GP_{\text{whole cell}}$. ^{a,b}Mean values with unlike letters were significantly different ($P < 0.05$). MO, maize oil diet; FO, fish oil diet.

the cell as described previously⁽²²⁾. The fold increase of generalised polarisation (GP) values, which indicate the liquid-ordered state of the membrane^(22,27,28), was calculated as $GP_{\text{contact region}}/GP_{\text{whole cell}}$.

Assessment of the co-localisation of PKC θ with lipid rafts

Isolated CD4⁺ T-cells were mixed with OVA-pulsed A-20 B-lymphoma cells (2:1) and seeded onto poly-L-lysine-precoated chambered cover glass slides. After a 30 min incubation period, cells were fixed in 4% formaldehyde for 20 min, rinsed with PBS and incubated with 10 mM-glycine in PBS for 10 min at room temperature to quench excess aldehyde groups. Cell membranes were permeabilised by exposure to 0.2% Triton X-100 in PBS for 5 min at room temperature, followed by washing with PBS. The cells were subsequently covered with blocking solution (1% bovine serum albumin/0.1% Na₂S₂O₈ in PBS) and incubated at 4°C overnight with a primary rabbit antibody specific to PKC θ (Santa Cruz Biotechnology). After washing with PBS, cells were incubated with a secondary Alexa Fluor[®] 568 goat antibody specific to rabbit IgG (Molecular Probes) and monosialotetrahexosylganglioside (GM1)-specific cholera toxin B subunit

conjugated with FITC (Sigma-Aldrich) for lipid raft visualisation. Following serial ethanol dehydration steps, samples were mounted onto glass slides with ProLong Antifade reagent (Molecular Probes). Fluorescence images were acquired by confocal microscopy to determine the co-localisation of PKC θ with GM1 using a Bio-Rad Radiance 2000MP multiphoton system (63× objective 1.4 NA water). Cell images were captured in 8-bit TIFF format, and the co-localisation of PKC θ with GM1 at the IS was calculated by Pearson's correlation using Image J software (NIH).

Statistics

All data were tested for normality followed by one-way ANOVA with Tukey's *post hoc* test using GraphPad Prism version 5.00 for Windows (GraphPad Software, Inc.). Differences in data at $P < 0.05$ were considered statistically significant.

Results

Incorporation of dietary n-3 PUFA into CD4⁺ T-cells

The fatty acid composition of the diets, expressed in mol% of individual fatty acids, was assessed by GC⁽²⁶⁾ and is presented in Table 1. As expected, n-3 PUFA, i.e. α -linoleic acid (ALA, 18:3n-3), EPA (20:5n-3) and docosapentaenoic acid (22:5n-3), were enriched in the FO diet, at the expense of oleic acid (18:1n-9) and linoleic acid (18:2n-6) when compared with the MO control diet. DHA (22:6n-3) was present in both the FO (5.55 (SEM 0.50) mol%) and DHA (4.09 (SEM 0.36) mol%) diets, and this difference was statistically significantly ($P < 0.05$). Consequently, the total n-3 PUFA content was highest in the FO diet (18.03 (SEM 1.11) mol%) followed by the DHA (4.63 (SEM 0.41) mol%) and MO (0.67 (SEM 0.05)) diets ($P < 0.05$ between the groups). During the 2-week dietary intervention, body weights were recorded at days 0, 7 and 14 and did not differ significantly between the groups (data not shown). In order to determine the enrichment of dietary fatty acids in CD4⁺ T-cells, total fatty acid profiles in purified splenic CD4⁺ T-cells were analysed

Table 1. Fatty acid composition of the diets

(Mean values with their standard errors, n 4 mice)

	MO (mol%)		FO (mol%)		DHA (mol%)	
	Mean	SEM	Mean	SEM	Mean	SEM
14:0	ND		9.14 ^b	0.11	3.29 ^a	0.04
16:0	12.33 ^a	0.05	24.90 ^b	0.59	13.74 ^a	0.04
16:1n-7	ND		10.65 ^b	0.14	0.36 ^a	0.01
18:0	2.11 ^b	0.09	4.20 ^a	0.02	1.89 ^b	0.01
18:1n-9	28.96 ^b	0.03	15.74 ^a	0.17	27.55 ^b	0.06
18:2n-6	55.93 ^c	0.18	17.34 ^a	0.13	48.54 ^b	0.44
18:3n-3	0.67 ^a	0.05	1.22 ^b	0.11	0.54 ^a	0.05
20:5n-3	ND		10.52	0.37	ND	
22:5n-3	ND		0.74	0.18	ND	
22:6n-3	ND		5.55 ^b	0.50	4.09 ^a	0.36
Σ n-3	0.67 ^a	0.05	18.03 ^c	1.11	4.63 ^b	0.41

MO, maize oil diet; FO, fish oil diet; ND, not detectable.

^{a,b,c}Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

Table 2. Fatty acid analysis of CD4⁺ T-cells following the dietary intervention

 (Mean values with their standard errors, *n* 4 mice)

	MO (mol%)		FO (mol%)		DHA (mol%)	
	Mean	SEM	Mean	SEM	Mean	SEM
14:0	3.23 ^a	0.37	3.82 ^a	0.43	9.15 ^b	0.61
16:0	41.35	0.55	36.82	5.81	44.42	2.10
16:1 <i>n</i> -7	0.14	0.12	0.31	0.04	0.14	0.11
18:0	29.41	1.50	33.35	0.47	22.39	6.00
18:1 <i>n</i> -9	6.50	0.84	5.72	1.56	4.19	1.18
18:1 <i>n</i> -7	0.73	0.02	0.78	0.19	0.29	0.24
18:2 <i>n</i> -6	7.96	0.55	6.45	2.30	4.78	0.87
18:3 <i>n</i> -3	5.05 ^b	2.43	5.25 ^b	1.99	0.80 ^a	0.65
20:4 <i>n</i> -6	5.63 ^b	1.32	4.51 ^b	0.28	1.98 ^a	1.62
20:5 <i>n</i> -3	ND		0.86	0.11	ND	
22:5 <i>n</i> -3	ND		0.47	0.03	ND	
22:6 <i>n</i> -3	ND		1.66 ^a	0.53	5.54 ^b	2.01
Σ <i>n</i> -3	5.05	2.43	7.38	2.51	6.35	3.36
Σ <i>n</i> -6	13.59 ^b	0.76	10.97 ^{a,b}	2.01	6.76 ^a	2.48
<i>n</i> -3: <i>n</i> -6	0.37 ^a	0.20	0.67 ^b	0.11	0.94 ^c	0.16

MO, maize oil diet; FO, fish oil diet; ND, not detectable.

^{a,b,c} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

(Table 2). CD4⁺ T-cells from FO-fed mice exhibited comparable ALA (FO 5.25 (SEM 1.99) *v.* MO 5.05 (SEM 2.43) mol%) and arachidonic acid (AA, 20:4*n*-6) (FO 4.51 (SEM 0.28) *v.* MO 5.63 (SEM 1.32) mol%) when compared with the MO control group, while EPA (0.86 (SEM 0.11) mol%), docosapentaenoic acid (0.47 (SEM 0.03) mol%) and DHA (1.66 (SEM 0.53)%) were detected in CD4⁺ T-cells from the FO group but not in the cells from the MO control group. In contrast, the cells from mice fed the DHA diet contained trace amounts of ALA (0.80 (SEM 0.65) mol%) and AA (1.98 (SEM 1.62) mol%) when compared with the MO (ALA 5.05 (SEM 2.43) and AA 5.63 (SEM 1.32) mol%) and FO (ALA 5.25 (SEM 1.99) and AA 4.51 (SEM 0.28) mol%) dietary groups, while DHA (5.54 (SEM 2.01) mol%) was highly enriched ($P < 0.05$). Total amounts of *n*-3 PUFA in CD4⁺ T-cells were not significantly different between the dietary groups, but the cells from mice fed the DHA diet exhibited significantly reduced ($P < 0.05$) levels of *n*-6 PUFA when compared with the MO group (6.76 (SEM 2.48) *v.* 13.59 (SEM 0.76) mol%, respectively). Consequently, the membrane ratio of *n*-3:*n*-6 was enhanced most by DHA feeding (0.94 (SEM 0.16)), followed by FO (0.67 (SEM 0.11)) and MO control (0.37 (SEM 0.20)) feeding ($P < 0.05$).

Increase in membrane liquid order at the immunological synapse following fish oil and DHA feeding

The formation of liquid-ordered mesodomains at the IS in CD4⁺ T-cells was assessed by Laurdan labelling as described previously^(22,28). Following antigen-specific IS formation and T-cell activation, DO11.10 T-cell receptor transgenic CD4⁺ T-cells from the MO control diet-fed mice exhibited an insignificant increase in membrane liquid order at the IS as determined by GP values (whole cell 0.18 (SEM 0.01) *v.* IS 0.20 (SEM 0.01), $P > 0.05$) (Fig. 1(A)). In contrast, GP values at the IS when compared with those of the whole T-cell for

FO-fed (whole cell 0.16 (SEM 0.01) *v.* IS 0.22 (SEM 0.01), $P < 0.05$) and DHA-fed groups (whole cell 0.17 (SEM 0.01) *v.* IS 0.20 (SEM 0.01)) were increased. In order to investigate whether GP values at the IS relative to the whole cell were significantly different among the dietary groups, we further calculated the fold increase by normalising GP_{immunological synapse} to GP_{whole cell}. Using this relative index, the FO-fed mice exhibited a significant increase ($P < 0.05$) in liquid-ordered mesodomains at the IS as expressed by the fold increase of GP values at the IS relative to the whole cell (FO 1.40 (SEM 0.11) *v.* MO 1.10 (SEM 0.06); Fig. 1(B)). These data indicate that membrane enrichment with *n*-3 PUFA following FO or DHA feeding accelerated the formation of liquid-ordered mesodomains during antigen-specific T-cell activation.

Suppressed co-localisation of PKCθ with GM1 by the fish oil- and DHA-containing diets

In order to determine whether enhanced formation of liquid-ordered mesodomains further alters CD4⁺ T-cell function, i.e. cellular signalling, the conjugates of CD4⁺ T-cells and antigen-presenting cells (OVA-pulsed B-lymphoma cells) were fixed on a glass slide followed by immunostaining for PKCθ (an essential T-cell signalling protein) and cholera toxin B subunit that binds to ganglioside GM1 (a lipid raft marker) (Fig. 2(A)). Subcellular immunofluorescence was captured by confocal microscopy and the co-localisation of the markers was determined at the IS by drawing an oval around the contact site between the T-cells and antigen-presenting cells. As determined by Pearson's coefficient, CD4⁺ T-cells from mice fed either the FO (0.37 (SEM 0.04)) or DHA (0.38 (SEM 0.04)) diet exhibited suppressed ($P < 0.05$) co-localisation of PKCθ with lipid rafts at the IS when compared with the MO control diet (0.48 (SEM 0.04)) (Fig. 2(B)).

Discussion

The anti-inflammatory effects of *n*-3 PUFA on CD4⁺ T-cells have been studied extensively by us and others^(7,21,22,24,29–32,33,34–37). The manipulation of lipid rafts, liquid-ordered mesodomains of the plasma membrane, by dietary fatty acids has emerged as one of the molecular targets of *n*-3 PUFA^(13,38,39–41). We have demonstrated previously that endogenous *n*-3 PUFA produced by *fat-1* transgenic mice accumulated in the plasma membrane and up-regulated the formation of lipid rafts in CD4⁺ T-cells at the IS⁽²²⁾. Subsequently, the function of CD4⁺ T cells, as assessed by the localisation of signalling proteins and cell proliferation, was inhibited by *n*-3 PUFA⁽²²⁾. Interestingly, Rockett *et al.*⁽⁴²⁾ recently demonstrated that dietary fish oil increased lipid raft size in B cells, indicating that *n*-3 PUFA-induced formation of lipid rafts is not a cell type-specific phenomenon.

In the present study, we investigated the effects of dietary *n*-3 PUFA on the formation of liquid-ordered mesodomains, which are thought to be the building blocks of lipid rafts, using a biologically relevant dose of fish oil. In addition, purified DHA, a major component of fish oil thought to mediate anti-inflammatory properties, was tested alone at levels




Fig. 2. (A) Representative immunofluorescence images for the analysis of the co-localisation of PKC θ with monosialotetrahexosylganglioside (GM1). Purified CD4⁺ T-cells and ovalbumin-pulsed B-lymphoma cells were co-incubated, fixed, permeabilised and labelled with a PKC θ -specific antibody and Alexa Fluor[®] 568 conjugated with a secondary antibody and GM1-specific fluorescein isothiocyanate–cholera toxin B subunit (CTX). The co-localisation of fluorescence signals was assessed at the immunological synapse. (B) The co-localisation of PKC θ at the immunological synapse, expressed as Pearson's coefficient (twelve to fifteen cells per mouse from four mice per diet were examined to obtain a total of forty-eight to fifty-seven observations). * Mean value was significantly different from that of the MO control diet ($P < 0.05$). MO, maize oil diet; FO, fish oil diet.

approximating fish oil supplementation. It should be noted that the other highly bioactive *n*-3 PUFA, i.e. EPA, is also present in the FO, but not the DHA, diet. Following a 2-week dietary intervention, T-cell membrane fatty acids were significantly enriched with *n*-3 PUFA. By comparing fatty acid profiles of the experimental diets (Table 1) and purified CD4⁺ T-cells (Table 2), it was noted that the profound difference in the amount of linoleic acid in the diets did not affect the small pool of linoleic acid in CD4⁺ T-cells. However, the membrane content of AA, a metabolite of linoleic acid and a precursor to pro-inflammatory eicosanoids, was decreased in the DHA group, indicative of an alternative pathway by which DHA controls inflammatory responses. As expected, *n*-3 PUFA such as ALA, EPA, docosapentaenoic acid and DHA were enriched in CD4⁺ T-cells following the consumption of the FO diet. In contrast, DHA deposition following DHASCO[®] (DHA only) feeding was greater than that in the CD4⁺ T-cells of mice fed the FO diet. Consequently, the total amounts of *n*-3 PUFA in CD4⁺ T-cells from mice fed the FO and DHA diets were not significantly different, even though EPA was enriched only in the cells from the FO dietary group. Since the different dietary sources of *n*-3 PUFA resulted in unique cellular *n*-3:*n*-6 ratios, we further tested the effect of *n*-3 PUFA on CD4⁺ T-cell function.

We have reported previously that CD4⁺ T-cells from *fat-1* transgenic mice, which were enriched endogenously with *n*-3 PUFA, exhibited enhanced lipid raft formation at the IS⁽²²⁾. We now report that the same effect was obtained by altering the lipid composition of the diet to include either FO or DHA (Fig. 1(A)), thus demonstrating both the robustness of the *n*-3 PUFA effect and its relevance for assessing the impact of dietary supplementation with *n*-3 PUFA. With regard to the magnitude of the diet-induced changes in GP values, Kaiser *et al.*⁽⁴³⁾ have reported that the difference in GP values between liquid-ordered large unilamellar vesicles and a mixed large unilamellar vesicles was approximately

0.1 (comparing liquid-ordered and liquid-disordered membrane states). In addition, incubation of Jurkat cells with a 2:1 ratio of cholesterol:7-ketocholesterol resulted in a change in the GP value of approximately 0.12, with a corresponding 40% decrease in IL-2 secretion when compared with control cells⁽⁴⁴⁾. These data suggest that small changes in GP values are biologically relevant, i.e. associated with perturbations in cell function.

A further investigation of the effects of the diets on GP values at the IS revealed that DHA affected the liquid order at the IS differently with respect to the combination of EPA and DHA provided in the FO diet (Fig. 1(B)). Specifically, the fold increase of the GP values at the IS relative to the whole cell was significantly different in FO-fed mice, but not in DHA-fed group, when compared with the MO control. This difference between the FO and DHA diets may be attributed to the higher levels of *n*-3 PUFA in the FO diet when compared with the DHA diet (Table 1). In this regard, Williams *et al.*⁽⁴¹⁾ demonstrated that both DHA and EPA can be incorporated into lipid rafts, though the avidity of DHA was as twice as that of EPA.

The elevated GP ratio at the IS was previously reported to be linked to the suppression of T-cell signalling⁽²²⁾. In the present study, we demonstrated that the co-localisation of PKC θ with GM1 at the IS (Fig. 2(A)) was suppressed comparably by the FO and DHA diets (Fig. 2(B)), indicating an effect of dietary *n*-3 PUFA on the formation of liquid-ordered mesodomains and subsequent localisation of signalling proteins. Indeed, Hou *et al.*⁽³⁵⁾ recently reported that DHA inhibited actin remodelling in T-cells, resulting in a suppression of signalling cascades.

In conclusion, we demonstrated that *n*-3 PUFA from distinct dietary sources (FO *v.* DHA) can be integrated into antigen-activated CD4⁺ T-cells, resulting in the modulation of plasma membrane liquid order and PKC θ translocation to the IS. The previously unappreciated effect of EPA, which was present only in the FO diet, will be pursued in future studies.

Acknowledgements

The present study was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT and Future Planning (2013R1A1A1006288) to W. K., and by NIH grant CA129444 and the US Department of Agriculture CSREES Special Grant, 'Designing Foods for Health' 2010–34402-20875 to R. S. C. The NRF, NIH and USDA had no role in the design, analysis or writing of the manuscript.

The authors' contributions are as follows: W. K., D. N. M. and R. S. C. contributed to the study design, data analysis and interpretation, and manuscript preparation and revision; W. K. and R. B. contributed to the data acquisition.

None of the authors has any conflicts of interest.

References

- Westendorf AM, Templin M, Geffers R, *et al.* (2005) CD4⁺ T cell mediated intestinal immunity: chronic inflammation versus immune regulation. *Gut* **54**, 60–69.
- Coccia M, Harrison OJ, Schiering C, *et al.* (2012) IL-1beta mediates chronic intestinal inflammation by promoting the accumulation of IL-17A secreting innate lymphoid cells and CD4⁺ Th17 cells. *J Exp Med* **209**, 1595–1609.
- Koltsova EK, Garcia Z, Chodaczek G, *et al.* (2012) Dynamic T cell–APC interactions sustain chronic inflammation in atherosclerosis. *J Clin Invest* **122**, 3114–3126.
- van der Lubbe PA, Reiter C, Riethmuller G, *et al.* (1993) Anti-CD4 therapy in rheumatoid arthritis. *Clin Exp Rheumatol* **11**, Suppl. 8, S151.
- Zhang X, Nakajima T, Goronzy JJ, *et al.* (2005) Tissue trafficking patterns of effector memory CD4⁺T cells in rheumatoid arthritis. *Arthritis Rheum* **52**, 3839–3849.
- Doherty TA, Soroosh P, Broide DH, *et al.* (2009) CD4⁺ cells are required for chronic eosinophilic lung inflammation but not airway remodeling. *Am J Physiol Lung Cell Mol Physiol* **296**, L229–L235.
- Zhang P, Kim W, Zhou L, *et al.* (2006) Dietary fish oil inhibits antigen-specific murine Th1 cell development by suppression of clonal expansion. *J Nutr* **136**, 2391–2398.
- Orr SK, Trepanier MO & Bazinet RP (2012) *n*-3 Polyunsaturated fatty acids in animal models with neuroinflammation. *Prostaglandins Leukot Essent Fatty Acids* **88**, 97–103.
- Kris-Etherton PM, Harris WS & Appel LJ (2002) Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation* **106**, 2747–2757.
- Adams LS, Phung S, Wu X, *et al.* (2008) White button mushroom (*Agaricus bisporus*) exhibits antiproliferative and proapoptotic properties and inhibits prostate tumor growth in athymic mice. *Nutr Cancer* **60**, 744–756.
- Rangel-Huerta OD, Aguilera CM, Mesa MD, *et al.* (2012) Omega-3 long-chain polyunsaturated fatty acids supplementation on inflammatory biomarkers: a systematic review of randomised clinical trials. *Br J Nutr* **10**, Suppl. 7, S159–S170.
- Kim W, McMurray DN & Chapkin RS (2009) Chemotherapeutic properties of *n*-3 polyunsaturated fatty acids – old concepts and new insights. *Immunol Endocr Metab Agents Med Chem* **9**, 38–44.
- Kim W, Khan NA, McMurray DN, *et al.* (2010) Regulatory activity of polyunsaturated fatty acids in T-cell signaling. *Prog Lipid Res* **49**, 250–261.
- Jia Q, Lupton JR, Smith R, *et al.* (2008) Reduced colitis-associated colon cancer in Fat-1 (*n*-3 fatty acid desaturase) transgenic mice. *Cancer Res* **68**, 3985–3991.
- Monk JM, Kim W, Callaway E, *et al.* (2012) Immunomodulatory action of dietary fish oil and targeted deletion of intestinal epithelial cell PPARdelta in inflammation-induced colon carcinogenesis. *Am J Physiol Gastrointest Liver Physiol* **302**, G153–G167.
- Robertson JM, Jensen PE & Evavold BD (2000) DO11.10 and OT-II T cells recognize a C-terminal ovalbumin 323–339 epitope. *J Immunol* **164**, 4706–4712.
- Meiri KF (2005) Lipid rafts and regulation of the cytoskeleton during T cell activation. *Philos Trans R Soc Lond B Biol Sci* **360**, 1663–1672.
- Jury EC, Flores-Borja F & Kabouridis PS (2007) Lipid rafts in T cell signalling and disease. *Semin Cell Dev Biol* **18**, 608–615.
- Nicolau DV Jr, Burrage K, Parton RG, *et al.* (2006) Identifying optimal lipid raft characteristics required to promote nanoscale protein–protein interactions on the plasma membrane. *Mol Cell Biol* **26**, 313–323.
- Kang JX, Wang J, Wu L, *et al.* (2004) Transgenic mice: fat-1 mice convert *n*-6 to *n*-3 fatty acids. *Nature* **427**, 504.
- Fan YY, Kim W, Callaway E, *et al.* (2008) Fat-1 transgene expression prevents cell culture-induced loss of membrane *n*-3 fatty acids in activated CD4⁺T-cells. *Prostaglandins Leukot Essent Fatty Acids* **79**, 209–214.
- Kim W, Fan YY, Barhoumi R, *et al.* (2008) *n*-3 Polyunsaturated fatty acids suppress the localization and activation of signaling proteins at the immunological synapse in murine CD4⁺T cells by affecting lipid raft formation. *J Immunol* **181**, 6236–6243.
- Sugano M (1996) Characteristics of fats in Japanese diets and current recommendations. *Lipids* **31**, S283–S286.
- Fan YY, McMurray DN, Ly LH, *et al.* (2003) Dietary (*n*-3) polyunsaturated fatty acids remodel mouse T-cell lipid rafts. *J Nutr* **133**, 1913–1920.
- Folch J, Lees M & Sloane Stanley GH (1957) A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* **226**, 497–509.
- Chapkin RS & Carmichael SL (1990) Effects of dietary *n*-3 and *n*-6 polyunsaturated fatty acids on macrophage phospholipid classes and subclasses. *Lipids* **25**, 827–834.
- Gaus K, Chklovskaja E, Fazekas de St Groth B, *et al.* (2005) Condensation of the plasma membrane at the site of T lymphocyte activation. *J Cell Biol* **171**, 121–131.
- Gaus K, Zech T & Harder T (2006) Visualizing membrane microdomains by Laurdan 2-photon microscopy. *Mol Membr Biol* **23**, 41–48.
- Fowler KH, Chapkin RS & McMurray DN (1993) Effects of purified dietary *n*-3 ethyl esters on murine T lymphocyte function. *J Immunol* **151**, 5186–5197.
- Costa Rosa LF, Safi DA & Guimaraes AR (1996) The effect of *N*-3 PUFA rich diet upon macrophage and lymphocyte metabolism and function. *Biochem Mol Biol Int* **40**, 833–842.
- Sasaki T, Kanke Y, Kudoh K, *et al.* (1999) Effects of dietary docosahexaenoic acid on surface molecules involved in T cell proliferation. *Biochim Biophys Acta* **1436**, 519–530.
- Chapkin RS, Arrington JL, Apanasovich TV, *et al.* (2002) Dietary *n*-3 PUFA affect TcR-mediated activation of purified murine T cells and accessory cell function in co-cultures. *Clin Exp Immunol* **130**, 12–18.
- Ly LH, Smith R 3rd, Chapkin RS, *et al.* (2005) Dietary *n*-3 polyunsaturated fatty acids suppress splenic CD4⁽⁺⁾ T cell function in interleukin (IL)-10(–/–) mice. *Clin Exp Immunol* **139**, 202–209.

34. Yog R, Barhoumi R, McMurray DN, *et al.* (2010) *n*-3 Polyunsaturated fatty acids suppress mitochondrial translocation to the immunologic synapse and modulate calcium signaling in T cells. *J Immunol* **184**, 5865–5873.
35. Hou TY, Monk JM, Fan YY, *et al.* (2012) *n*-3 Polyunsaturated fatty acids suppress phosphatidylinositol 4,5-bisphosphate-dependent actin remodelling during CD⁴⁺T-cell activation. *Biochem J* **443**, 27–37.
36. Monk JM, Jia Q, Callaway E, *et al.* (2012) Th17 cell accumulation is decreased during chronic experimental colitis by (*n*-3) PUFA in Fat-1 mice. *J Nutr* **142**, 117–124.
37. Shaikh SR, Jolly CA & Chapkin RS (2012) *n*-3 Polyunsaturated fatty acids exert immunomodulatory effects on lymphocytes by targeting plasma membrane molecular organization. *Mol Aspects Med* **33**, 46–54.
38. Chapkin RS, Wang N, Fan YY, *et al.* (2008) Docosahexaenoic acid alters the size and distribution of cell surface microdomains. *Biochim Biophys Acta* **1778**, 466–471.
39. Shaikh SR (2012) Biophysical and biochemical mechanisms by which dietary *N*-3 polyunsaturated fatty acids from fish oil disrupt membrane lipid rafts. *J Nutr Biochem* **23**, 101–105.
40. Turk HF & Chapkin RS (2012) Membrane lipid raft organization is uniquely modified by *n*-3 polyunsaturated fatty acids. *Prostaglandins Leukot Essent Fatty Acids*.
41. Williams JA, Batten SE, Harris M, *et al.* (2012) Docosahexaenoic and eicosapentaenoic acids segregate differently between raft and nonraft domains. *Biophys J* **103**, 228–237.
42. Rockett BD, Teague H, Harris M, *et al.* (2012) Fish oil increases raft size and membrane order of B cells accompanied by differential effects on function. *J Lipid Res* **53**, 674–685.
43. Kaiser HJ, Lingwood D, Levental I, *et al.* (2009) Order of lipid phases in model and plasma membranes. *Proc Natl Acad Sci U S A* **106**, 16645–16650.
44. Rentero C, Zech T, Quinn CM, *et al.* (2008) Functional implications of plasma membrane condensation for T cell activation. *PLoS One* **3**, e2262.