

## Serum 25-hydroxyvitamin D levels showed strong seasonality but lacked association with vitamin D intake in 3-year-old Japanese children

Emiko Ando<sup>1,2</sup>, Naho Morisaki<sup>1\*</sup>, Keiko Asakura<sup>3</sup>, Satoshi Sasaki<sup>2</sup>, Takeo Fujiwara<sup>4</sup> and Reiko Horikawa<sup>5</sup>

<sup>1</sup>Department of Social Medicine, National Center for Child Health and Development, 2-10-1, Okura, Setagaya-ku, Tokyo 157-8535, Japan

<sup>2</sup>Department of Social and Preventive Epidemiology, School of Public Health, The University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>3</sup>Department of Environmental and Occupational Health, School of Medicine, Toho University, 5-21-16, Omori-Nishi, Ota-ku, Tokyo 143-8580, Japan

<sup>4</sup>Department of Global Health Promotion, Tokyo Medical and Dental University, 1-5-45, Yushima, Bunkyo-ku, Tokyo 113-8519, Japan

<sup>5</sup>Department of Endocrinology, National Center for Child Health and Development, 2-10-1, Okura, Setagaya-ku, Tokyo 157-8535, Japan

(Submitted 9 February 2018 – Final revision received 23 May 2018 – Accepted 12 July 2018 – First published online 29 August 2018)

### Abstract

Current guidelines provide a universal recommendation on vitamin D intake to prevent insufficiency. However, the relative influence of food, UVB and other factors on serum 25-hydroxyvitamin D (25(OH)D) insufficiency has been poorly investigated in preschool children. We assessed serum 25(OH)D quantities and their association with vitamin D intake using a brief-type self-administered diet history questionnaire for children aged 3–6 years (BDHQ3y), outdoor playing time and background UVB radiation level among 574 36-month-old Japanese children living at latitude 35°N. The average serum 25(OH)D concentration was 23.5 (sd 6.1) ng/ml, and 170 (29.6%) children had vitamin D insufficiency (<20 ng/ml) despite high consumption of fish. Multiple logistic regression adjusting for social factors showed that when background UVB radiation level was <15 kJ/m<sup>2</sup> (monthly average), there was a 1.89 (95% CI 1.31, 2.74) times higher risk of vitamin D insufficiency, to which vitamin D intake nor time spent outdoors were significantly associated. ANOVA showed that the contribution of the variability in vitamin D intake on the variability of serum 25(OH)D level was 1.8% of that of UVB exposure. The correlation between vitamin D intake and serum 25(OH)D level was not stronger when limited to measurements in winter. We found that nearly 30% of 3-year-old Japanese children had vitamin D insufficiency despite high consumption of fish and living at relatively low latitude. We failed to observe an association between vitamin D intake and the risk of vitamin D insufficiency. This may be due to the extremely limited access to vitamin D-fortified food and supplements for children in Japan.

**Key words:** Vitamin D: Food: Serum 25-hydroxyvitamin D: UVB: Preschool children: Japan: Fish

Vitamin D is an essential micronutrient for growth, development and adequate puberty onset in children<sup>(1–3)</sup>. Historically, a low serum level of 25-hydroxyvitamin D (25(OH)D) has been associated with growth failure and skeletal deformation<sup>(2,3)</sup>. Recent studies also showed that even without such clinical symptoms, low levels of serum 25(OH)D may delay maturation of the immune system<sup>(4)</sup> and increase the risk of future asthma<sup>(5)</sup>, respiratory infections<sup>(5–9)</sup>, auto-immune diseases and metabolic disorders<sup>(10,11)</sup>. Thus, the high prevalence of sub-clinical vitamin D deficiency (even without clinical symptoms), otherwise called vitamin D insufficiency, is considered a public health problem<sup>(3,5,12)</sup>.

Serum 25(OH)D is influenced by both dietary intake of vitamin D and exposure to sunlight<sup>(2,13)</sup>. Current recommendations (by the American Academy of Paediatrics (AAP) 2014, Institute of Medicine (IOM) 2011)<sup>(14,15)</sup> state that children 1 year old and older should receive at least 600 IU (15 µg) of vitamin D daily through diet or supplements regardless of levels of sun exposure as awareness of the adverse effects of UVB exposure on health, such as the increased risk of future skin cancer, has risen<sup>(3,4,16)</sup>. Recent studies showed that serum 25(OH)D status differed also by other factors such as age, latitude and skin colour<sup>(1,12)</sup>, suggesting that population-specific recommendations may be better than a universal recommendation for preventing vitamin D deficiency.

**Abbreviations:** 25(OH)D, 25-hydroxyvitamin D; BDHQ, brief-type self-administered diet history questionnaire for children aged 3–6 years.

\* **Corresponding author:** N. Morisaki, email morisaki-n@nchcd.go.jp

Unfortunately, most of the studies of how food, UVB exposure and other factors influence vitamin D status in preschool children have been done in high-latitude Western countries (51–62°N) where prolonged periods of winter darkness reduce the skin's natural production of vitamin D<sup>(17–19)</sup> no studies have been conducted in lower latitude countries like Japan.

Furthermore, the extent of the influence of dietary vitamin D on serum 25(OH)D levels, that is, the relative contributions of vitamin D intake from food and UVB exposure on serum concentrations, has been poorly studied in preschool children. In adults, studies estimated that 15–42% of vitamin D derives from food such as fish, eggs and milk whereas the rest is synthesised by the skin through exposure to sunlight<sup>(5,20,21)</sup> (especially to UVB radiation in the range of 280–320 nm)<sup>(1–3,13)</sup>. Such knowledge may be useful for designing interventions to reduce vitamin D deficiency.

We, therefore, investigated factors associated with serum 25(OH)D status and evaluated the relative contributions of vitamin D intake from food and UVB exposure on serum 25(OH)D levels in a cohort of 3-year-old preschool children in Japan.

## Methods

### Subjects

Our study was based on the Seiiku Boshi cohort study, a prospective, hospital-based birth cohort, whose primary aim was to explore how antenatal factors influenced maternal health and the child's growth, function and risk of disease<sup>(22)</sup>. For this cohort, pregnant women attending their first antenatal visit at the National Center for Child Health and Development (NCCHD (35°N)) were recruited from 13 May 2010 to 28 November 2013 and were asked to complete a questionnaire which included several items on socio-demographic factors. Serial, onsite interviews of the respondents and their children were also conducted. Of 4164 women, 2310 consented to participate in the study. Most participants lived in or near Tokyo (35°N).

The 36-month follow-up interview was conducted monthly from July 2014 to October 2016 at the NCCHD and scheduled according to each child's birth month. A questionnaire including a brief-type self-administered diet history questionnaire for children aged 3–6 years (BDHQ3y) as well as a question on daily outdoor playtime was delivered to the participants 1 month before the interview and was submitted at the interview. Half of the participants were also randomly selected to undergo a blood test at the interview.

Among 1370 participants who were given the questionnaires and asked to take a blood test at the 36-month follow-up interview, serum 25(OH)D levels were measured in 721 (52.6%), among whom, 574 (79.6%) completed and returned both questionnaires.

The protocol of the cohort study (project number 417, approved on 2 August 2010) was approved by the Institutional Review Board at the NCCHD in Tokyo, Japan. Written informed consent was obtained from the mothers on behalf of their children.

### Brief-type self-administered diet history questionnaire for children aged 3–6 years

The BDHQ3y was developed to assess the dietary intake in Japanese children of 3–6 years of age during the preceding month<sup>(23)</sup>.

The BDHQ3y is four pages long and contains the following four sections: (1) intake frequency of fifty-seven food and non-alcoholic beverage items; (2) daily intake of rice, including type of rice (refined, unrefined or rice boiled with barley) and miso soup; (3) usual cooking methods and (4) general dietary habits. As the Japanese diet includes a wide range of fish species cooked in a variety of ways, the BDHQ3y orders fish into seven categories (squid, octopus, shrimp and shellfish; small fish with bones; canned tuna; dried fish (e.g. salted mackerel, salted salmon and dried horse mackerel); oily fish (e.g. sardine, mackerel, pacific saury, yellowtail, herring, eel and tuna); lean fish (such as salmon, trout, white fish, freshwater fish and bonito) and fish paste (such as fish sausage and boiled fish paste)). The BDHQ3y also asks about the frequency of the consumption of vitamin D-fortified foods or vitamin D supplements in the previous month but does not ask the respondent to name the food or supplement.

The BDHQ3y was constructed using the BDHQ, which was developed for adults and has been validated by previous studies<sup>(24–26)</sup>; however, the BDHQ3y excludes the section pertaining to the consumption of alcoholic beverages and coffee and includes a section pertaining to food items frequently consumed by children, such as yogurt drinks, French fries, chocolate and ketchup.

To assess the daily intake of sixty-six food items, the total energy content and forty-two selected nutrients based on responses to the BDHQ3y, we used an *ad hoc* computer algorithm<sup>(23,27)</sup>. The BDHQ3y does not collect data on the portion sizes of food consumed by the subjects and instead uses age-specific portion sizes of food and beverages based on previous studies<sup>(23,27)</sup>. A more detailed methodology for assessing portion size has been reported previously<sup>(23)</sup>. Crude values for the daily total intake of energy content and forty-two selected nutrients, including vitamin D, were calculated based on both the data reported on the BDHQ3y and the values given for the individual items on the *Standard Tables of Food Composition in Japan*<sup>(28)</sup>. In detail, the daily intake of vitamin D was calculated as the sum of vitamin D in selected food items, for example, cereals, vegetables except for green and yellow vegetables (including mushrooms and seaweeds), fish and shellfish, meat, eggs, dairy products, fats, snacks, beverages and seasonings<sup>(23)</sup>. Pearson and Spearman correlations between the BDHQ3y and a 3-d dietary record for vitamin D were 0.37 and 0.24, respectively<sup>(23)</sup>.

### Serum 25-hydroxyvitamin D concentration

Samples were centrifuged for 10 min at 3000 rpm to separate the sera from the pellet. The sera were stored at –20°C until measurement. The levels of 25(OH)D in the sera were estimated based on the 25(OH)D concentrations measured by competitive protein binding assay using a liquid scintillation counter developed by LSI Medience Corporation.

The tests were conducted by LSI Medience Corporation. The intra- and interassay coefficients of validation for 25(OH)D concentration measurement were 7.3–10.3 and 8.0–10.8%, respectively. We defined vitamin D insufficiency as a serum 25(OH)D concentration <20 ng/ml based on previous studies<sup>(2–4,12)</sup>.

### Assessment of outdoor playtime and background UVB level

The UVB exposure is influenced by both sunlight exposure time and the background UVB radiation level, which differs by season. Therefore, we collected information on outdoor playtime and background UVB radiation level at the time of the serum 25(OH)D assessment.

The questionnaire administered for the 36-month follow-up contained one question on regards to outdoor playtime: 'On average, how long does your child stay outside for the following activities (walking to kindergarten or nursery, walking to a park, walking to the supermarket or convenience store, walking in the park) per day?' The response choices were as follows: (1) 15 min, (2) 30 min, (3) 45 min, (4) 60 min and (5) more than 60 min. If a mother answered (5), she was next asked to indicate how many hours. None of the questions asked about the use of sunscreen.

For assessment of the background UVB radiation level, we used the monthly average of daily measurements reported by the Japan Meteorological Agency of the total UVB radiation for Tsukuba (36°N), a suburban town located approximately 75 km from Tokyo (35°N). Based on these reports, we created two categories: one based on whether UVB was above or below 15 kJ/m<sup>2</sup> and another based on whether UVB was above or below 10 kJ/m<sup>2</sup>. For all the years from 2011 to 2013, October–March had UVB <15 kJ/m<sup>2</sup>, and November–February had UVB <10 kJ/m<sup>2</sup>.

**Assessment of other covariates.** Socio-demographic factors including household income and the educational background of the parents were ascertained by the questionnaire administered at enrolment. Data were completed for maternal education and household income; however, thirty children had missing data on the father's education. We placed these children into the 'missing father's education status' group.

The BDHQ3y also asks about current height, weight as well as the response date. We calculated the BMI from self-reported height and weight. If the data were missing, they were complemented by measurements obtained at the 36-month visit.

**Statistical analysis.** First, we compared the average serum 25(OH)D level and the risk of vitamin D insufficiency based on tertiles of vitamin D intake from food, five categories pertaining to outdoor playtime and background UVB radiation level at the month of assessment, using test for trend.

Next, we assessed the association between vitamin D insufficiency and vitamin D intake from food, daily outdoor playtime, background UVB radiation level, household income and parental educational background using multiple logistic regressions. In addition, we assessed the association between serum 25(OH)D level and the factors listed above using multiple linear regression.

Third, we analysed the Pearson correlation coefficients between serum 25(OH)D level and vitamin D intake, outdoor playtime and season at the time of assessment in order to calculate the individual effects of food intake and sun exposure on serum 25(OH)D level. To assess the relative contribution of vitamin D intake from food, socio-economic factors and sun

exposure on serum 25(OH)D levels, we performed ANOVA using least squares to fit linear models, in which the relative contribution of each factor was calculated by dividing the partial sum of squares for each variable by the partial sum of squares of the full model.

Fourth, as the effect of vitamin D intake on serum 25(OH)D levels may be stronger among those with less UVB exposure, we repeated this analysis among subgroups of the sample using two different cut-offs for UVB exposure (<10 and <15 kJ/m<sup>2</sup>). In addition, we investigated whether vitamin D intake differed between children with and without vitamin D insufficiency among those with low UVB exposure (<10 or <15 kJ/m<sup>2</sup>).

The sample size for this study was determined by the feasibility of recruitment (expected number of participants of 3-year follow-up was *n* 800) and the estimated minimally detectable correlation was *r* 0.10 ( $\alpha=0.05$ ,  $\beta=0.20$ , required sample size *n* 783). The minimally detectable correlation of the final sample (*n* 574) was *r* 0.117<sup>(29)</sup>.

All statistical analysis were conducted using Stata version 14, and the results were represented as means and standard deviations. All tests were two-tailed, with a *P* values <0.05 considered as statistically significant.

### Results

In total, serum 25(OH)D measurements, valid BDHQ3y assessment of food intake and self-reported daily outdoor play time were obtained from 574 children (271 girls and 303 boys) at 36 months of age. All participants were free from disease including vitamin D-deficient rickets.

Characteristics of the study participants are shown in Table 1. The average serum 25(OH)D concentration was 23.5 (SD 6.1) ng/ml while the average vitamin D intake from food was 3.1 (SD 1.8) µg/d. The largest source of vitamin D was fish, with the daily vitamin D intake from fish being 2.4 (SD 1.7) µg/d. Overall, the proportion of children with vitamin D insufficiency was 29.6% (170 children), and 8.5% of children (forty-nine children) had a level below 15 ng/ml. The proportion of children with vitamin D insufficiency did not differ according to the quantity of vitamin D obtained from food.

Fig. 1 shows serum 25(OH)D concentration by levels of vitamin D and sun exposure (daily outdoor playtime and background UVB radiation level). The average serum 25(OH)D concentration did not differ between children according to the level of vitamin D intake from food (high: middle: low, *P*-trend=0.26) or outdoor play time (15 min: 30 min: 45 min: 60 min: more than 60 min, *P*-trend=0.14). On the other hand, serum 25(OH)D levels were significantly higher among measurements conducted when UVB was above 15 kJ/m<sup>2</sup> (April to September) compared with those conducted when UVB was below 15 kJ/m<sup>2</sup> (October to March) (*P*<0.001), as well as those conducted when UVB was above 10 kJ/m<sup>2</sup> (November to February) compared with months when UVB was above 10 kJ/m<sup>2</sup> (March to October) (*P*<0.001). Measurements in July and August showed the highest values while those in January showed the lowest values (online Supplementary Fig. S1).

Table 2 shows the association between vitamin D insufficiency and vitamin D intake from food, daily outdoor play time,

**Table 1.** Characteristics of study participants (Mean values and standard deviations; numbers and percentages)

Covariates	Total (n 574)		Serum 25(OH)D ≥ 20 ng/ml (n 404, vitamin D sufficiency)		Serum 25(OH)D < 20 ng/ml (n 170, vitamin D insufficiency)		P†
	Mean	SD	Mean	SD	Mean	SD	
Height (cm)	92.1	3.6	92.1	3.5	92.3	3.6	0.251
Weight (kg)	13.5	1.5	13.5	1.4	13.6	1.7	0.296
BMI (kg/m <sup>2</sup> )	15.9	1.4	15.9	1.3	16.0	1.7	0.552
BMI z-score‡	0.26	1.03	0.25	1.00	0.29	1.11	0.671
Serum 25(OH)D level (ng/ml)***	23.5	6.1	26.5	4.6	16.5	2.5	<0.001
Vitamin D intake from food (µg/d)	3.1	1.8	3.2	1.8	3.0	1.7	0.097
Fish	2.4	1.7	2.5	1.7	2.3	1.6	0.123
Milk (not fortified)	0.3	0.3	0.3	0.3	0.3	0.3	0.167
Eggs	0.2	0.1	0.2	0.1	0.2	0.1	0.262
		<i>n</i>	Population (%)	<i>n</i>	Vitamin D deficiency (%)§		
Sex							
Boys		303	52.8	80	26.4		
Girls		271	47.2	90	33.2		
Household income							
<4 million yen		40	7.0	14	35.0		
<8 million yen		184	32.1	61	33.2		
≥8 million yen		350	61.0	95	27.1		
Mother's education							
High school		96	16.7	34	35.4		
2 years of college		119	20.7	38	31.9		
University or graduate school		359	62.5	98	27.3		
Father's education							
Missing		30	5.2	11	36.7		
Under 2 years of college		101	18.6	34	33.7		
University		350	64.3	107	30.6		
Graduate school		93	17.1	18	19.4		
Vitamin D intake							
Low (<2.24 µg/d)		191	33.3	59	30.9		
Middle (2.24–3.43 µg/d)		191	33.3	66	34.6		
High (>3.43 µg/d)		192	33.4	45	23.4		
Daily outdoor play time (min)							
15		85	14.8	25	29.4		
30		126	22.0	45	35.7		
45		50	8.7	13	26.0		
60		193	33.6	57	29.5		
Over 60		120	20.9	30	25.0		
Monthly average of daily UVB radiation							
UVB >15 kJ/m <sup>2</sup>							
April–September		308	53.7	71	23.1		
UVB <15 kJ/m <sup>2</sup>							
October–March		266	46.3	99	37.2		
UVB >10 kJ/m <sup>2</sup>							
March–October		400	53.7	106	26.5		
UVB <10 kJ/m <sup>2</sup>							
November–February		174	46.3	64	36.8		

25(OH)D, 25-hydroxyvitamin D.

\*\*\* *P* ≤ 0.001.

† *P* value derived from Mann–Whitney *U* test comparing participants without vitamin D insufficiency to those with vitamin D insufficiency.


‡ BMI z-score was calculated by using the WHO Child Growth Standards (2007) as reference data.

§ Percentage with vitamin D insufficiency of the subgroup members.

background UVB radiation level, household income and parental educational background. In the adjusted model, background UVB radiation level showed the strongest association with vitamin D insufficiency, with measurements conducted in winter (when UVB <15 kJ/m<sup>2</sup>) having almost twice the odds of indicating vitamin D insufficiency compared with those obtained in summer when UVB >15 kJ/m<sup>2</sup> (OR 1.89, 95% CI 1.31, 2.74, *P* < 0.001). Compared with children whose intake was in the highest tertile (>3.43 µg/d), children whose intake

was in the middle tertile (2.24–3.43 µg/d) had 1.62 (95% CI 1.03, 2.57) times higher odds of vitamin D insufficiency, while those in the lowest tertile (<2.24 µg/d) having 1.32 (95% CI 0.83, 2.10) times higher odds of vitamin D insufficiency.

Table 3 shows the association between serum 25(OH)D concentration and vitamin D intake from food, daily outdoor playtime, background UVB radiation level, household income and parental educational background. In this model, vitamin D intake did not show a significant association with serum



**Fig. 1.** Serum 25-hydroxyvitamin D (25(OH)D) levels by vitamin D intake (a), daily outdoor playtime (b) and background UVB radiation level at measurement (c and d). \*  $P < 0.001$ . The box plots show the median and interquartile range (IQR); the whiskers represent 2.5 times the IQR above and below the median for each variable. Data not included between the whiskers are shown as dots. UVB cut-off is  $15 \text{ kJ/m}^2$  in (c) and  $10 \text{ kJ/m}^2$  in (d).  $P$  values calculated for non-parametric test for trend is 0.257 for (a) and 0.142 for (b).  $P$  value calculated for Mann–Whitney  $U$  test is  $P < 0.001$  for (c) and (d).

25(OH)D level after adjusting for other factors. Children whose intake was in the lowest tertile ( $<2.24 \mu\text{g/d}$ ) had on average  $-0.25$  (95% CI  $-1.45, 0.94$ ) ng/ml lower serum 25(OH)D levels than children whose intake was in the highest tertile ( $>3.43 \mu\text{g/d}$ ). Background UVB radiation level and the father's educational status were significantly associated with serum 25(OH)D concentration, with measurements conducted in winter (when  $\text{UVB} < 15 \text{ kJ/m}^2$ ) being on average  $-2.91$  (95% CI  $-3.89, -1.93$ ) ng/ml lower than those conducted in summer when  $\text{UVB} > 15 \text{ kJ/m}^2$ , and measurements in children missing data on paternal educational status were on average  $-3.90$  (95% CI  $-6.38, -1.43$ ) ng/ml lower than in those who completed graduate school.

Table 4 shows the Pearson correlations between serum 25(OH)D levels and vitamin D intake, outdoor playtime and background UVB radiation level. No significant correlation was observed between serum 25(OH)D levels and vitamin D intake ( $r 0.060, P=0.15$ ; scatter plot with fitted linear regression line shown in Fig. 2). Outdoor play time was also not significantly correlated with serum 25(OH)D levels ( $r 0.067, P=0.11$ ); however, background UVB radiation level was

positively correlated with serum 25(OH)D levels ( $r 0.262, P < 0.001$ ).

Next, we performed ANOVA to assess the relative contribution of vitamin D intake from food, socio-economic factors and sun exposure on serum 25(OH)D levels simultaneously because vitamin D intake was significantly correlated with outdoor play time ( $r 0.15, P < 0.001$ ), suggesting that socio-economic status was influencing both. The ANOVA showed that the largest contributor to variations in the serum 25(OH)D level was background UVB radiation (63%) followed by socio-economic factors (23%). The contributions of variability in vitamin D intake from food (1.2%) and variability in outdoor play time (2.7%) were both small. In this analysis, the contribution of variability in vitamin D intake from food on the variability of serum 25(OH)D level was 1.8% of that of UVB exposure.

In a subgroup analysis assessing whether the correlation between vitamin D intake and serum 25(OH)D levels was stronger among those with less UVB exposure (among measurements conducted when the background UVB radiation level was  $<15$  and  $<10 \text{ kJ/m}^2$ ), the correlation coefficient

**Table 2.** Associations between socio-economic status, vitamin D intake from food, daily outdoor play time and season with the likelihood of vitamin D insufficiency (25-hydroxyvitamin D <20 nmol/l)† (Odds ratios and 95% confidence intervals)

Covariates	Crude		Adjusted	
	OR	95% CI	OR	95% CI
Household income				
<4 million yen	1.45	0.72, 2.88	1.33	0.64, 2.74
<8 million yen	1.33	0.90, 1.96	1.34	0.89, 2.03
≥8 million yen		Ref.		Ref.
Mother's education				
High school	1.46	0.91, 2.36	1.34	0.78, 2.29
2 years of college	1.25	0.80, 1.96	1.21	0.75, 1.94
University or graduate school		Ref.		Ref.
Father's education				
Missing	2.41	0.98, 5.95	2.41	0.95, 6.16
Under 2 years of college	2.11*	1.09, 4.09	1.56	0.76, 3.16
University	1.83*	1.05, 3.22	1.53	0.86, 2.72
Graduate school		Ref.		Ref.
Vitamin D intake				
Low (<2.24 µg/d)	1.46	0.93, 2.30	1.32	0.83, 2.10
Middle (2.24–3.43 µg/d)	1.72*	1.10, 2.70	1.62*	1.03, 2.57
High (>3.43 µg/d)		Ref.		Ref.
Daily outdoor play time (min)				
15	1.25	0.67, 2.33	0.94	0.82, 1.07
30	1.67	0.96, 2.89		
45	1.05	0.50, 2.24		
60	1.26	0.75, 2.11		
Over 60		Ref.		
Monthly average of daily UVB radiation				
UVB >15 kJ/m <sup>2</sup>				
April–September		Ref.		Ref.
UVB <15 kJ/m <sup>2</sup>				
October–March	1.98***	1.38, 2.85	1.89***	1.31, 2.74

Ref., reference.

\*  $P < 0.05$ , \*\*\*  $P \leq 0.001$ .

† Adjusted for all covariates in this table.

remained low ( $r$  0.017 and 0.044) and statistically insignificant.

Fig. 3 shows the distribution of vitamin D intake among children with and without vitamin D insufficiency when the average monthly UVB radiation was below 15 and 10 kJ/m<sup>2</sup>. The values did not differ significantly. Similarly, the average vitamin D intake did not differ significantly between children with and without vitamin D insufficiency when the analysis was restricted to those with shorter outdoor playtime (data not shown).

### Discussion

This study indicated that in Japanese preschool children, the prevalence of vitamin D insufficiency was relatively high at 29.6%. Vitamin D intake from food did not have a significant association with the serum 25(OH)D level consistently regardless of the level of UVB exposure or other factors. Socio-economic factors and seasonality were stronger predictors of serum 25(OH)D levels.

Japanese (and other Asians) generally have been considered to have higher serum 25(OH)D levels as their diet includes more fish<sup>(30–34)</sup>, a very rich source of the nutrient<sup>(2,12,35)</sup>. However, interestingly, the prevalence of vitamin D insufficiency (29.6%) was not lower in our study than in those done among children in Western, higher latitude countries where less

fish is consumed (12–29%)<sup>(3,17–19,36–39)</sup>. This was true despite the fact that the children included in our study reported high levels of fish intake (at least three times the amount consumed among children in other countries)<sup>(18,19,40)</sup>. In addition, our subjects' average amount of vitamin D intake was similar to the value recently reported by the National Health and Nutrition Survey Japan in 2016 (3.7 (SD 3.8) µg/d, median: about 2.1 µg/d vitamin D intake by children 1–6 years)<sup>(41)</sup>. Our findings suggest that vitamin D insufficiency may be highly prevalent throughout Japan despite the relatively high consumption of vitamin D-rich foods, a finding supported by a recent, population-based study, which revealed that rickets caused by vitamin D deficiency has increased nationwide<sup>(42)</sup>.

This result may be partly due to certain lifestyle factors peculiar to Japan. Avoidance of excessive sun exposure has become more and more the rule, while access to vitamin D-fortified foods and vitamin D supplements for children remains very poor (vitamin D-fortified food itself is scarce in Japan, and there is no recommendation for vitamin D supplementation for children; vitamin D supplements for infants were introduced only in 2014)<sup>(42)</sup>. Oily fish such as sardine are the most vitamin D-rich oily fish in Japan, but in order to consume 15 µg (600 IU) of vitamin D per d<sup>(14,15)</sup> as recommended by the IOM and AAP, children need to eat 300 g of sardines every week. This may not be realistic for 3-year-old children.

**Table 3.** Associations between socio-economic status, vitamin D intake from food, daily outdoor play time and season with serum 25-hydroxyvitamin D level† (Mean differences (MD) and 95 % confidence intervals)

Covariates	Crude		Adjusted	
	MD	95 % CI	MD	95 % CI
Household income				
<4 million yen	-2.03*	-4.03, -0.02	-1.62	-3.62, 0.37
<8 million yen	-0.79	-1.88, 0.31	-0.98	-2.08, 0.11
≥8 million yen		Ref.		Ref.
Mother's education				
High school	-0.95	-2.33, 0.43	-0.61	-2.06, 0.83
2 years of college	0.16	-1.12, 1.43	0.21	-1.05, 1.47
University or graduate school		Ref.		Ref.
Father's education				
Missing	-3.95**	-6.52, -1.44	-3.90***	-6.38, -1.43
Under 2 years of college	-2.18*	-3.90, -0.46	-1.22	-2.99, 0.55
University	-1.58*	-2.97, -0.18	-0.96	-2.33, 0.41
Graduate school		Ref.		Ref.
Vitamin D intake				
Low (<2.24 µg/d)	-0.67	-1.90, 0.56	-0.25	-1.45, 0.94
Middle (2.24–3.43 µg/d)	-0.76	-1.99, 0.47	-0.44	-1.63, 0.76
High (>3.43 µg/d)		Ref.		Ref.
Daily outdoor play time (min)				
15	-0.75	-2.46, 0.95	0.25	-0.11, 0.60
30	-1.23	-2.77, 0.30		
45	-0.17	-2.19, 1.86		
60	-0.08	-1.48, 1.32		
Over 60		Ref.		
Monthly average of daily UVB radiation				
UVB >15 kJ/m <sup>2</sup>		Ref.		Ref.
April–September				
UVB <15 kJ/m <sup>2</sup>				
October–March	-3.01***	-3.99, -2.04	-2.91***	-3.89, -1.93

Ref., reference.

\*  $P < 0.05$ , \*\*  $P \leq 0.01$ ; \*\*\*  $P \leq 0.001$ .

† Adjusted for all covariates in this table.

**Table 4.** Correlations between vitamin D intake and sun exposure with serum 25-hydroxyvitamin D levels

	All		UVB <15 kJ/m <sup>2</sup> (n 266)		UVB <10 kJ/m <sup>2</sup> (n 174)	
	Correlation	P	Correlation	P	Correlation	P
Vitamin D intake from food (µg/d)	0.060	0.154	0.017	0.778	0.044	0.566
Daily outdoor play time (min/d)	0.067	0.110	0.007	0.908	0.013	0.867
Monthly average of daily UVB radiation (kJ/m <sup>2</sup> )	0.262	0.000	0.113	0.067	0.202	0.008

Serum 25(OH)D levels are influenced not only by dietary intake but also by vitamin D synthesised by the skin's exposure to UVB radiation (280–320 nm)<sup>(2,12,13,35,43,44)</sup>. We found that the background UVB radiation level was significantly associated with serum 25(OH)D levels in line with previous studies, which have shown this association in infants and school-aged children<sup>(3,17,18,20,30,39,40,45–48)</sup>. Seasonal differences in serum 25(OH)D levels were large, with the prevalence of vitamin D insufficiency being two times higher in winter than in summer, underscoring the large effect of UVB exposure on serum 25(OH)D levels even at relatively low latitudes<sup>(35,43)</sup>. While a previous study estimated that, in adults, 15–42% of vitamin D was derived from food whereas the rest was synthesised<sup>(5,20,21)</sup>, the relative contributions of these two sources have not been reported for children. Our study found that in Japanese

preschool children, the contribution of dietary vitamin D to the variability of serum 25(OH)D level was much smaller and only 1.8% that of UVB exposure. Several reasons may have led to this discrepancy. Our population lived at a much lower altitude (35°N) than the subjects in the previous study (51°N); thus, while the amount of vitamin D consumed was similar, the relative contribution of UVB may have been higher. Our study used a FFQ with a greater possibility of measurement errors in estimating vitamin D intake compared with the previous study which used 4- and 7-d food diaries. Further studies evaluating the contribution of dietary vitamin D on vitamin D status in preschool children are required.


While we found seasonality to have a consistent and strong influence on vitamin D status, we failed to find a consistent significant association between dietary vitamin D and the risk of

vitamin D insufficiency and failed to find any significant association between dietary vitamin D and the average serum 25 (OH) level. Furthermore, the contribution of vitamin D intake on the risk of vitamin D insufficiency was moderate, with risk among children whose intake was in the lowest tertile estimated to be 1.3 times higher than in children whose intake was in the highest tertile, a much smaller value compared that of the effect of UVB exposure. Even when we repeated the analysis after restricting the population to those with low UVB exposure following previous reports suggesting the influence of dietary vitamin D intake was higher when UVB exposure was low<sup>(35,43)</sup>, the correlation between dietary vitamin D intake and serum 25(OH)D level remained weak and insignificant. Taken together, our study suggested that increasing the dietary intake of vitamin D may not substantially alter serum 25(OH)D levels in Japanese children.

This lack of association between dietary intake and serum 25 (OH)D levels did not agree with the findings of previous studies, most of which showed a positive association in children. This discrepancy may be due to the very limited availability of vitamin D-fortified foods or (inactivated) vitamin

D supplements in Japan<sup>(42)</sup>. While the majority of the vitamin D intake in previous cohort studies was derived from fortified milk or supplements, and the amounts deriving from naturally vitamin D-rich foods (such as fish and eggs) contributed to only 9–12% of total intake<sup>(18,40)</sup>, in the participants in our study, 85% of the vitamin D intake came from fish (not fortified). Milk, which is not fortified with vitamin D in Japan and only contains 0.3 µg (12 IU)/100 g<sup>(49)</sup> of the nutrient, contributed a mere 10% to the total vitamin D intake. Our study suggested that increasing the consumption of natural (not fortified) vitamin D-rich foods may not be very effective in increasing serum 25 (OH)D levels. Rather, fortification or supplementation may be the more realistic way of preventing vitamin D deficiency when sufficient UVB exposure is not available<sup>(17,50)</sup>.

Our study also failed to find a significant association between outdoor playtime and the serum 25(OH)D level unlike previous studies of breastfed infants and school-aged children<sup>(20,47,51)</sup>. This may have been due to the crudity of our measurement of sunlight exposure, as we measured only outdoor playtime but did not consider the area of skin exposed, sunscreen use or time of day when the exposure occurred, since very little if any vitamin D is produced before 10.00 hours and after 15.00 hours<sup>(13)</sup>.



**Fig. 2.** Scatter plot between vitamin D intake from food and serum 25-hydroxyvitamin D (25(OH)D) levels with fitted linear regression line. ○, Subjects; —, fitted values.


**Table 5.** Relative contribution of vitamin D intake from food, socio-economic factors and sun exposure on serum 25-hydroxyvitamin D levels

	Relative contribution (%)†
Vitamin D intake from food (µg/d)	1.22
Socio-economic factors‡	22.86
Monthly average of daily UVB radiation*** (kJ/m <sup>2</sup> )	62.76
Daily outdoor play time (min/d)	2.65
Total	100

\*\*\*  $P \leq 0.001$ .

† Percentage of relative contribution was derived by dividing the partial sum of squares for each variable by the partial sum of squares of the full model.

‡ Socio-economic factors include household income, mother's education and father's education.



**Fig. 3.** Vitamin D intake between those with and without vitamin D insufficiency among children with low UVB exposure, using different definitions ((a) UVB strength  $< 15 \text{ kJ/m}^2$ , (b) UVB strength  $< 10 \text{ kJ/m}^2$ ). The box plots show the median and interquartile ranges (IQR); the whiskers represent 2.5 times the IQR above and below the median for each variable. Data not included between the whiskers are shown as dots.  $P$  value calculated for Mann–Whitney  $U$  test is 0.270 for (a) and 0.380 for (b).



Several recent studies showed that vitamin D levels could also vary according to social and environmental factors such as skin colour and socio-economic status<sup>(45,47,52–54)</sup>. In our study, we found the risk of vitamin D insufficiency to vary according to parental characteristics (children with the highest risk, that is, children whose mothers reported their own education status but omitted the father's, were likely to be single mothers). In addition, outdoor playtime was significantly correlated to vitamin D intake from food (data not shown). These findings suggest that health-consciousness, which is likely related to socio-economic status, may be an important contributing factor to both vitamin D intake and outdoor playtime and may explain why we found socio-economic factors to be the second largest contributor to variability in serum 25(OH)D levels following UVB exposure when the relative contribution of each factor was analysed. Future studies on how and why socio-economic status influences serum 25(OH)D are required (Table 5).

These findings suggested several policy implications. Due to the increasing knowledge that subclinical vitamin D deficiency, also called as vitamin D insufficiency, may be deleterious to children's health<sup>(3,4)</sup>, it is recommended that a serum 25(OH)D concentration of 20 ng/ml (equivalent to 50 nmol/l) should be maintained in infants and children<sup>(3,15)</sup>. As Asians consume relatively higher amounts of fish, a rich source of vitamin D, the risk of vitamin D insufficiency was assumed to be low in this population. However, we found that in Japanese children, the prevalence of vitamin D insufficiency was quite high even though their fish consumption was relatively high, outdoor playtime relatively long and latitude not high.

A possible explanation of the lack of an association between vitamin D intake and vitamin D insufficiency may be the recent, popular use of sunscreen and long-sleeved shirts to avoid UVB exposure<sup>(13)</sup>. We showed, in our study, that the contribution of vitamin D intake to serum 25(OH)D levels was small compared with that of vitamin D synthesised through UVB exposure. Thus, small increases in vitamin D intake by increasing the consumption of vitamin D-rich foods may not be sufficient to reduce the risk of vitamin D insufficiency. If increasing UVB exposure by decreasing the use of sunscreen and long sleeves is not justifiable (a valid question, given that Asian populations including the Japanese have a relatively lower risk of skin cancer than Caucasians<sup>(55)</sup>), increasing vitamin D intake through supplementation or fortification at least during seasons, when the background UVB radiation level is low, may be a realistic approach for reducing the prevalence of vitamin D insufficiency in preschool children.

The strengths of our study include the unified age range of the preschool participants (36 months). Our population was also ethnically homogeneous (all Japanese) and the latitude was also the same (Tokyo, 35°N), which maximises the study's power to observe the influence of other factors on serum 25(OH)D concentrations. Another strength is that we used recorded UVB radiation data (reported by the Japan Meteorological Agency) rather than the crude categories of season and were thus able to measure the contribution of UVB on serum 25(OH)D levels.

There were several limitations to our study. First, measurements of vitamin D intake from food may not be precise

since previous validation disclosed that the Pearson and Spearman correlation coefficient for vitamin D was 0.35 and 0.30 for the BDHQ3y and the dietary records, respectively<sup>(23)</sup>. However, the BDHQ3y has a detailed structure for estimating fish intake and is well suited to the dietary culture of Japan, and for nutrients such as EPA and DHA, which are also abundant in oily fish, the correlation between the BDHQ3y and dietary records (0.36–0.40, Asakura *et al.*<sup>(23)</sup>) was similar to that between BDHQ3y estimates and serum values of EPA and DHA (E Ando, N Morisaki, K Asakura, K Ogawa, S Sasaki, R Horikawa and T Fujiwara, unpublished results). We, therefore, concluded that the BDHQ3y was able to estimate vitamin D intake from fish at a level acceptable enough to show a correlation between vitamin D intake and serum values if the influence of UVB were small. In addition, we did not have information on vitamin D supplementation; however, vitamin D supplementation is not commonly available in Japan<sup>(42)</sup>, and 96% of our subjects responded on the BDHQ that they did not consume any vitamin D-fortified foods or vitamin D supplements in the previous month (data not shown). Thus, we believe that measurement errors due to not measuring supplement intake are insignificant. Second, we used CPBA to measure serum 25(OH)D, but liquid chromatography–tandem MS-based methods, which are more accurate and reliable, are now becoming more readily available. Therefore, we may have over- or underestimated 25(OH)D concentrations in our study. Further studies using newer laboratory methods may be required. Third, our study omitted several factors influencing sun exposure, such as clothing, the use of sunscreen and the time of day on which the exposure occurred<sup>(13)</sup>, which likely led to underestimating the effect of UVB exposure time on serum 25(OH)D concentration. Fourth, the sample size of our study, while comparable to those of previous studies, was relatively small ( $n = 574$ ). We may have failed to detect associations due to type 2 error, and further studies with larger samples are warranted to confirm our results. Finally, our study was not entirely representative of the general population as the study participants were not population-based but were volunteers from an affluent subpopulation of Tokyo (35°N). Thus, the families may have been more health conscious and as a result provided their children with vitamin D-rich food and taken their children outdoors more frequently but used more sunscreen than the general population. The smaller variability in vitamin D intake (with few children having low intake) may have led to a reduction in the power to detect its association with serum 25(OH)D levels.

In conclusion, in Japanese preschool children, vitamin D intake from food was not significantly associated with serum 25(OH)D levels regardless of UVB exposure and socio-economic factors. Our results suggest that increasing the intake of vitamin D from food may not substantially affect serum 25(OH)D levels in Japanese preschool children. Other methods of increasing serum 25(OH)D levels apart from promoting increased the consumption of vitamin D-rich food, such as increasing the availability of vitamin D supplementation, fortifying food and re-evaluating how much UVB exposure should be allowed for Asian children, may help reduce vitamin D insufficiency in Japan.

## Conclusions

Nearly 30% of Japanese children 3 years of age had vitamin D insufficiency despite high consumption of fish (naturally rich in vitamin D) and lived at relatively low latitude. In our study, we failed to observe a significant association between vitamin D intake and serum 25(OH)D level as well as the risk of vitamin D insufficiency, which may be due to the extremely limited access to vitamin D-fortified food and supplements for children in Japan. Promoting increases in vitamin D-rich food may not be sufficient to reduce risk of vitamin D insufficiency for children in Japan.

## Acknowledgements

The authors are deeply indebted to all participants in the Seikku Boshi Cohort Study.

This work was supported by the Research Development Grant for Child Health and Development (grant no. H25-4) and the Japan Agency for Medical Research and Development (grant no. AMED-6013).

The authors' responsibilities were as follows: N. M., E. A., K. A., T. F. and R. H. designed the study; N. M. and R. H. conducted research; K. A. and S. S. provided essential materials; E. A. and N. M. analysed the data; E. A. and N. M. wrote the manuscript to which K. A. provided substantial comments; N. M. had primary responsibility for final content; and all authors read and approved the final manuscript.

The authors declare that there are no conflicts of interest.

## Supplementary material

For supplementary material/s referred to in this article, please visit <https://doi.org/10.1017/S0007114518002258>

## References

- Holick MF (1994) McCollum Award Lecture, 1994: vitamin D—new horizons for the 21st century. *Am J Clin Nutr* **60**, 619–630.
- Holick MF (2007) Vitamin D deficiency. *N Engl J Med* **357**, 266–281.
- Casey CF, Slawson DC & Neal LR (2010) Vitamin D supplementation in infants, children, and adolescents. *Am Fam Physician* **81**, 745–748.
- Wagner CL & Greer FR, American Academy of Pediatrics Section on Breastfeeding, *et al.* (2008) Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics* **122**, 1142–1152.
- Bozzetto S, Carraro S, Giordano G, *et al.* (2012) Asthma, allergy and respiratory infections: the vitamin D hypothesis. *Allergy* **67**, 10–17.
- Roth DE, Shah R, Black RE, *et al.* (2010) Vitamin D status and acute lower respiratory infection in early childhood in Sylhet, Bangladesh. *Acta Paediatr* **99**, 389–393.
- Wayse V, Yousafzai A, Mogale K, *et al.* (2004) Association of subclinical vitamin D deficiency with severe acute lower respiratory infection in Indian children under 5 y. *Eur J Clin Nutr* **58**, 563–567.
- Karatekin G, Kaya A, Salihoglu O, *et al.* (2009) Association of subclinical vitamin D deficiency in newborns with acute lower respiratory infection and their mothers. *Eur J Clin Nutr* **63**, 473–477.
- McNally JD, Leis K, Matheson LA, *et al.* (2009) Vitamin D deficiency in young children with severe acute lower respiratory infection. *Pediatr Pulmonol* **44**, 981–988.
- Hypponen E, Laara E, Reunanen A, *et al.* (2001) Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet* **358**, 1500–1503.
- Lee DY, Kwon AR, Ahn JM, *et al.* (2015) Relationship between serum 25-hydroxyvitamin D concentration and risks of metabolic syndrome in children and adolescents from Korean National Health and Nutrition Examination survey 2008–2010. *Ann Pediatr Endocrinol Metab* **20**, 46–52.
- Hilger J, Friedel A, Herr R, *et al.* (2014) A systematic review of vitamin D status in populations worldwide. *Br J Nutr* **111**, 23–45.
- Wacker M & Holick MF (2013) Sunlight and vitamin D: a global perspective for health. *Dermatoendocrinology* **5**, 51–108.
- Golden NH & Abrams SA, Committee on Nutrition (2014) Optimizing bone health in children and adolescents. *Pediatrics* **134**, e1229–e1243.
- Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium (2011) Summary Tables. In *Dietary Reference Intakes for Calcium and Vitamin D* [AC Ross, CL Taylor, AL Yaktine and HB Del Valle, editors]. Washington, DC: National Academies Press.
- Balk SJ, Binns HJ & Brumberg HL (2011) Ultraviolet radiation: a hazard to children and adolescents. *Pediatrics* **127**, 588–597.
- Munasinghe LL, Yuan Y, Willows ND, *et al.* (2017) Vitamin D deficiency and sufficiency among Canadian children residing at high latitude following the revision of the RDA of vitamin D intake in 2010 – Corrigendum. *Br J Nutr* **117**, 1052–1054.
- Ni Chaoimh C, McCarthy EK, Hourihane JO, *et al.* (2018) Low vitamin D deficiency in Irish toddlers despite northerly latitude and a high prevalence of inadequate intakes. *Eur J Nutr* **57**, 783–794.
- Akkermans MD, van der Horst-Graat JM, Eussen SR, *et al.* (2016) Iron and vitamin D deficiency in healthy young children in western Europe despite current nutritional recommendations. *J Pediatr Gastroenterol Nutr* **62**, 635–642.
- Dawodu A, Davidson B, Woo JG, *et al.* (2015) Sun exposure and vitamin D supplementation in relation to vitamin D status of breastfeeding mothers and infants in the global exploration of human milk study. *Nutrients* **7**, 1081–1093.
- Macdonald HM, Mavroei A, Fraser WD, *et al.* (2011) Sunlight and dietary contributions to the seasonal vitamin D status of cohorts of healthy postmenopausal women living at northerly latitudes: a major cause for concern? *Osteoporos Int* **22**, 2461–2472.
- Ogawa K, Morisaki N, Kobayashi M, *et al.* (2018) Maternal vegetable intake in early pregnancy and wheeze in offspring at the age of 2 years. *Eur J Clin Nutr* **72**, 761–771.
- Asakura K, Haga M & Sasaki S (2015) Relative validity and reproducibility of a brief-type self-administered diet history questionnaire for Japanese children aged 3–6 years: application of a questionnaire established for adults in preschool children. *J Epidemiol* **25**, 341–350.
- Kobayashi S, Murakami K, Sasaki S, *et al.* (2011) Comparison of relative validity of food group intakes estimated by comprehensive and brief-type self-administered diet history questionnaires against 16 d dietary records in Japanese adults. *Public Health Nutr* **14**, 1200–1211.

25. Okubo H, Sasaki S, Rafamantanantsoa HH, *et al.* (2008) Validation of self-reported energy intake by a self-administered diet history questionnaire using the doubly labeled water method in 140 Japanese adults. *Eur J Clin Nutr* **62**, 1343–1350.
26. Kobayashi S, Honda S, Murakami K, *et al.* (2012) Both comprehensive and brief self-administered diet history questionnaires satisfactorily rank nutrient intakes in Japanese adults. *J Epidemiol* **22**, 151–159.
27. Asakura K, Haga M, Adachi M, *et al.* (2014) Estimation of food portion sizes frequently consumed by children 3–6 years old in Japan. *J Nutr Sci Vitaminol (Tokyo)* **60**, 387–396.
28. The Council for Science and Technology (2010) *Standard Tables of Food Composition in Japan*, 5th rev. and enlarged ed., pp. 5–229 [Ministry of Education, Culture, Sports, Science and Technology of Japan, editor]. Tokyo: Kagawa Education Institute of Nutrition.
29. Lachin JM (1981) Introduction to sample size determination and power analysis for clinical trials. *Control Clin Trials* **2**, 93–113.
30. Yu A, Kim J, Kwon O, *et al.* (2013) The association between serum 25-hydroxyvitamin D concentration and consumption frequencies of vitamin D food sources in Korean adolescents. *Clin Nutr Res* **2**, 107–114.
31. Lee MS, Li HL, Hung TH, *et al.* (2008) Vitamin D intake and its food sources in Taiwanese. *Asia Pac J Clin Nutr* **17**, 397–407.
32. Okuda M, Sasaki S, Bando N, *et al.* (2009) Carotenoid, tocopherol, and fatty acid biomarkers and dietary intake estimated by using a brief self-administered diet history questionnaire for older Japanese children and adolescents. *J Nutr Sci Vitaminol (Tokyo)* **55**, 231–241.
33. Yamori Y, Nara Y, Iritani N, *et al.* (1985) Comparison of serum phospholipid fatty acids among fishing and farming Japanese populations and American inlanders. *J Nutr Sci Vitaminol (Tokyo)* **31**, 417–422.
34. Nakamura T, Takebe K, Tando Y, *et al.* (1995) Serum fatty acid composition in normal Japanese and its relationship with dietary fish and vegetable oil contents and blood lipid levels. *Ann Nutr Metab* **39**, 261–270.
35. Holick MF (2004) Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart disease, and osteoporosis. *Am J Clin Nutr* **79**, 362–371.
36. Gordon CM, Feldman HA, Sinclair L, *et al.* (2008) Prevalence of vitamin D deficiency among healthy infants and toddlers. *Arch Pediatr Adolesc Med* **162**, 505–512.
37. Gordon CM, DePeter KC, Feldman HA, *et al.* (2004) Prevalence of vitamin D deficiency among healthy adolescents. *Arch Pediatr Adolesc Med* **158**, 531–537.
38. Marrero NM, He FJ, Whincup P, *et al.* (2014) Salt intake of children and adolescents in South London: consumption levels and dietary sources. *Hypertension* **63**, 1026–1032.
39. Petersen RA, Damsgaard CT, Dalskov SM, *et al.* (2016) Vitamin D status and its determinants during autumn in children at northern latitudes: a cross-sectional analysis from the optimal well-being, development and health for Danish children through a healthy New Nordic Diet (OPUS) School Meal Study. *Br J Nutr* **115**, 239–250.
40. Soininen S, Eloranta AM, Lindi V, *et al.* (2016) Determinants of serum 25-hydroxyvitamin D concentration in Finnish children: the Physical Activity and Nutrition in Children (PANIC) study. *Br J Nutr* **115**, 1080–1091.
41. Ministry of Health, Labour, and Welfare of Japan (2016) *The National Health and Nutrition Survey in Japan*. Tokyo: Ministry of Health and Welfare.
42. Itoh M, Tomio J, Toyokawa S, *et al.* (2017) Vitamin D-deficient rickets in Japan. *Glob Pediatr Health* **4**, 2333794X17711342.
43. Holick MF (2006) Resurrection of vitamin D deficiency and rickets. *J Clin Invest* **116**, 2062–2072.
44. MacLaughlin JA, Anderson RR & Holick MF (1982) Spectral character of sunlight modulates photosynthesis of previtamin D<sub>3</sub> and its photoisomers in human skin. *Science* **216**, 1001–1003.
45. Thorisdottir B, Gunnarsdottir I, Steingrimsdottir L, *et al.* (2016) Vitamin D intake and status in 6-year-old Icelandic children followed up from infancy. *Nutrients* **8**, 75.
46. Zhao X, Xiao J, Liao X, *et al.* (2015) Vitamin D status among young children aged 1–3 years: a cross-sectional study in Wuxi, China. *PLOS ONE* **10**, e0141595.
47. Voortman T, van den Hooven EH, Heijboer AC, *et al.* (2015) Vitamin D deficiency in school-age children is associated with sociodemographic and lifestyle factors. *J Nutr* **145**, 791–798.
48. Han SW, Kang HR, Kim HG, *et al.* (2013) Subclinical vitamin D insufficiency in Korean school-aged children. *Pediatr Gastroenterol Hepatol Nutr* **16**, 254–260.
49. Ministry of Education, Culture, Sports, Science and Technology of Japan (2015) Standard tables of food composition in Japan 2015 (seventh revised version). [http://www.mext.go.jp/en/policy/science\\_technology/policy/title01/detail01/1374030.htm](http://www.mext.go.jp/en/policy/science_technology/policy/title01/detail01/1374030.htm) (accessed May 2018).
50. Akkermans MD, Eussen SR, van der Horst-Graat JM, *et al.* (2017) A micronutrient-fortified young-child formula improves the iron and vitamin D status of healthy young European children: a randomized, double-blind controlled trial. *Am J Clin Nutr* **105**, 391–399.
51. Dawodu A, Zalla L, Woo JG, *et al.* (2014) Heightened attention to supplementation is needed to improve the vitamin D status of breastfeeding mothers and infants when sunshine exposure is restricted. *Matern Child Nutr* **10**, 383–397.
52. Madsen KH, Rasmussen LB, Mejborn H, *et al.* (2014) Vitamin D status and its determinants in children and adults among families in late summer in Denmark. *Br J Nutr* **112**, 776–784.
53. Hintzpetter B, Scheidt-Nave C, Muller MJ, *et al.* (2008) Higher prevalence of vitamin D deficiency is associated with immigrant background among children and adolescents in Germany. *J Nutr* **138**, 1482–1490.
54. Carpenter TO, Herreros F, Zhang JH, *et al.* (2012) Demographic, dietary, and biochemical determinants of vitamin D status in inner-city children. *Am J Clin Nutr* **95**, 137–146.
55. Division of Cancer Prevention and Control Centers for Disease Control and Prevention (2017) Skin cancer rates by race and ethnicity. <https://www.cdc.gov/cancer/skin/statistics/race.htm> (accessed February 2018).