

Validation of a Case Definition for Pediatric Brain Injury Using Administrative Data

Jane McChesney-Corbeil, Karen Barlow, Hude Quan, Guanmin Chen,
Samuel Wiebe, Nathalie Jette

ABSTRACT: Background: Health administrative data are a common population-based data source for traumatic brain injury (TBI) surveillance and research; however, before using these data for surveillance, it is important to develop a validated case definition. The objective of this study was to identify the optimal International Classification of Disease, edition 10 (ICD-10), case definition to ascertain children with TBI in emergency room (ER) or hospital administrative data. We tested multiple case definitions. **Methods:** Children who visited the ER were identified from the Regional Emergency Department Information System at Alberta Children's Hospital. Secondary data were collected for children with trauma, musculoskeletal, or central nervous system complaints who visited the ER between October 5, 2005, and June 6, 2007. TBI status was determined based on chart review. Sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) were calculated for each case definition. **Results:** Of 6639 patients, 1343 had a TBI. The best case definition was, "1 hospital or 1 ER encounter coded with an ICD-10 code for TBI in 1 year" (sensitivity 69.8% [95% confidence interval (CI), 67.3-72.2], specificity 96.7% [95% CI, 96.2-97.2], PPV 84.2% [95% CI 82.0-86.3], NPV 92.7% [95% CI, 92.0-93.3]). The nonspecific code S09.9 identified >80% of TBI cases in our study. **Conclusions:** The optimal ICD-10-based case definition for pediatric TBI in this study is valid and should be considered for future pediatric TBI surveillance studies. However, external validation is recommended before use in other jurisdictions, particularly because it is plausible that a larger proportion of patients in our cohort had milder injuries.

RÉSUMÉ: Validation d'une définition de cas de la lésion cérébrale chez l'enfant au moyen de données administratives. Contexte : Les données administratives sur la santé sont une source courante d'informations sanitaires populationnelles pour la surveillance et la recherche sur le traumatisme crânien (TC). Cependant, avant d'utiliser ces données à des fins de surveillance, il est important d'établir une définition de cas validée. L'objectif de cette étude était d'identifier la définition de cas optimale de la 10^e édition de la Classification internationale des maladies (CIM-10), pour identifier les enfants atteints d'un TC à la salle d'urgence ou dans les données administratives hospitalières. Nous avons évalué plusieurs définitions de cas. **Méthodologie :** Les enfants qui se sont présentés à la salle d'urgence ont été identifiés dans le *Regional Emergency Department Information System* de l'Hôpital pour enfants de l'Alberta. Nous avons recueilli les données secondaires des enfants qui se sont présentés à la salle d'urgence pour des problèmes d'origine traumatique, musculosquelettiques ou du système nerveux central entre le 5 octobre 2005 et le 6 juin 2007. Nous avons déterminé s'ils présentaient un TC au moyen d'une revue de dossiers. Nous avons calculé la sensibilité, la spécificité, la valeur prédictive positive (VPP) et la valeur prédictive négative (VPN) de chaque définition de cas. **Résultats :** Parmi les 6 639 patients, 1 343 avaient subi un TC. La meilleure définition de cas était « une visite à l'hôpital ou une visite à la salle d'urgence portant le code de la CIM-10 pour le TC au cours d'une période d'un an » (sensibilité de 69,8% et IC à 95% : 67,3 à 72,2 ; spécificité de 96,7% et IC à 95% de 96,2 à 97,2 ; VPP de 84,2% et IC à 95% de 82,0 à 86,3 ; VPN de 92,7% et IC à 95% de 92,0 à 93,3). Le code non spécifique S09,9 identifiait plus de 80% des cas de TC dans notre étude. **Conclusions :** Dans cette étude, la définition de cas optimale basée sur la CIM-10 du TC chez l'enfant est valide et devrait être prise en compte dans les études de surveillance du TC chez l'enfant à l'avenir. Cependant, nous recommandons de procéder à une validation externe avant son utilisation dans d'autres juridictions, parce qu'il est plausible qu'une grande proportion de patients dans notre cohorte étaient atteints de blessures plus légères.

Keywords: Health Services Research, Brain Injury – Pediatric Health Services

doi:10.1017/cjn.2016.419

Can J Neurol Sci. 2017; 44: 161-169

Traumatic brain injury (TBI) is the leading cause of death and disability in children and youth in the United States, with a combined estimate of 511,257 emergency room (ER) visits, hospitalizations,

and deaths per year, and more than \$1 billion in hospital charges annually.^{1,2} One Canadian report showed that children (<18 years of age) with head injury represented almost 45% of those visiting ERs

From the Department of Community Health Sciences and Institute for Public Health (JM-C, HQ, GC, SW, NJ); Department of Clinical Neurosciences and Hotchkiss Brain Institute (JM-C, SW, NJ); Department of Pediatrics (KB), University of Calgary, Calgary, Alberta.

RECEIVED OCTOBER 27, 2015. FINAL REVISIONS SUBMITTED OCTOBER 5, 2016. DATE OF ACCEPTANCE OCTOBER 6, 2016.

Correspondence to: Nathalie Jette, Foothills Medical Center, Department of Clinical Neurosciences, 1403 29th Street NW, Calgary, Alberta T2N 4N1, Canada.
Email: Nathalie.jette@albertahealthservices.ca

and urgent care centers, and almost 25% of those were hospitalized in acute care.³ In addition, a recent systematic review and meta-analysis of the incidence of TBI from international studies identified that the pediatric (<15 years of age) incidence rate of traumatic brain injury was 134/100,000 person-years.⁴ Two Canadian pediatric studies were included in this review and reported incidence rates for mild TBI of 110/100,000⁵ and 0.81/100,⁶ respectively. TBI thus contributes substantially to the health care burden. Pediatric TBI is of particular concern because it can lead to significant and lifelong deficits,⁷ with a substantial burden on the patient, their caregivers, and society. Unfortunately, ongoing surveillance for pediatric TBI is lacking in Canada, resulting in gaps in the knowledge needed to adequately plan for the resources required for those affected by pediatric TBI.

Administrative data are valuable sources for disease surveillance because they are collected routinely, are cost-effective, cover wide geographic areas, and have a relatively complete capture of all patient encounters with the health care system.⁸ This is especially relevant to the Canadian system because there are comprehensive population-based administrative databases at regional, provincial, and national levels.⁹ In the majority of administrative databases, conditions are coded using the World Health Organization's International Classification of Disease (ICD) system. ICD-coded databases are used internationally to capture causes of morbidity and mortality, and for epidemiological, health management, and reimbursement purposes.¹⁰

To effectively capture cases of pediatric TBI, it is important to develop a validated case definition. A systematic review looking at validated case definitions for TBI identified three studies, all from the United States that met eligibility criteria (Table 1).¹¹ Only one of these studies reported more than one test of diagnostic accuracy, comparing coding of an inpatient database to real-time assessment in the ER.¹² They found low sensitivity (45.9%), but high specificity (97.8%) and negative predictive value (NPV) (99.2%).¹² In addition, only one study validated ICD-10 coding for TBI in administrative data,¹³ whereas the others validated ICD, 9th revision (ICD-9); Table 1.^{12,14} A validated case definition to study TBI outside of the United States has not, to our knowledge, been published. It is important to validate ICD coding specifically in other non-US databases for several reasons. Although the underlying coding framework is the same internationally, different countries may use slightly modified versions of ICD.¹⁵ For example, the United States still uses the ICD-9 Clinical Modification edition,¹⁶ which contains fewer codes than the ICD-10 edition currently being used in Canada. In addition, Canada's health care system differs from that in the United States, where coding standards may vary.¹⁵ Finally, Canadian coders undergo national standardized training, unlike coders in other countries, potentially leading to variations in coding standards between countries.^{15,17}

The goal of the present study was to validate ICD-10 coding for TBI in two administrative databases in a large health region: the Discharge Abstract Database (DAD), which captures all hospitalizations, and the Ambulatory Care Classification System (ACCS), which captures all ER/urgent care visits and hospital clinic visits in the province of Alberta. Because Canada has a universal health care system, these two population-based databases capture all patients who access these services. The two objectives were to: (1) assess the validity of ICD-10 coding for TBI and (2) develop a validated case definition for pediatric TBI. It was our hypothesis that it would be possible to develop an

accurate sensitive case definition to identify pediatric TBI using ICD-10 coded administrative health data.

METHODS

Study Site

Alberta Children's Hospital (ACH) is the primary pediatric hospital for the city of Calgary and surrounding semiurban and rural areas. ACH has a total of 133 inpatient beds, and its ER has 60,000 visits per year for those aged 0 to 18 years. The estimated total population of Calgary was 1,079,310 per the 2006 Canadian census, with 272,275 aged 0 to 19 years.¹⁸

Defining Study Population

The study population includes 7101 children aged 0 to 18 years, with primarily trauma, musculoskeletal, or central nervous system complaints captured between October 4, 2005, and June 6, 2007, at ACH through the Regional Emergency Department Information System (REDIS). These conditions were selected to increase the likelihood of capturing children who may have experienced a TBI.

REDIS is an electronic information system used to collect and track information on all children who visit the ACH ER. The "free text" diagnosis and clinical history fields of the REDIS charts were first reviewed by a trained research assistant to identify cases by looking for key words in the chart that would suggest a neurological problem, such as altered mental state, dizziness, confusion, disorientation, headache, balance problems, nausea and/or vomiting, blurred vision, seizures, concussion, head injury, and/or TBI. These charts were then reviewed again by a pediatric neurologist (KB) with expertise in brain injury to confirm diagnosis. Currently, there is not a single agreed-upon case definition for TBI. For this study, TBI was defined as a force to the body or head that results in disruption of normal function of the brain.¹⁹ TBI was confirmed if the injury was associated with any of the following neurological symptoms: loss of consciousness, altered mental state, posttraumatic amnesia, headache, dizziness, vomiting, confusion, and/or behavioral change. The definition of mild TBI is somewhat controversial in younger children (<5 years of age), in whom it can be difficult to assess an altered mental state and period of amnesia. Therefore, any young child presenting to the ER with a mild head injury without clear evidence contradicting the these criteria was included in the study. Children with simple scalp lacerations, facial injuries/fractures without symptoms of head injury or concussion described previously, or superficial injuries were excluded. Finally, the date the child was first seen in REDIS became the index date for all children.

The quality of the REDIS chart review was further verified among a subset of patients (n = 715) through a follow-up telephone call to confirm TBI diagnosis (n = 670) or who visited the ACH TBI clinic (n = 45). Of the 670 children that were successfully contacted by telephone for follow-up, only 27 (4%) were determined to not be TBI cases upon further questioning. All of the 45 children seen in the TBI clinic were true cases of TBI as originally determined by the REDIS chart review (100%).

Defining TBI in ICD-10 Administrative Data

In this study, TBI coding was validated in two administrative databases: ACCS and the inpatient DAD from April 1, 2002, to

Table 1: Summary of sensitivity, specificity, PPV, and NPV reported by validation studies for TBI

Author, year, location	Data year	Database validated	Reference standard	Study size	ICD version	ICD codes	Sn	Sp	PPV	NPV
Bazarian, 2006, United States ¹²	2003	Single hospital; inpatient, ER visits	ER real-time assessment	516	ICD-9-CM	800.0, 800.5, 801.0, 801.5, 803.0, 803.5, 804.0, 804.5, 850.0, 850.5, 850.9, 854.0, 959.01	45.9	97.8	23.7	99.2
Rodriguez, 2006, United States ¹³	2002	Population-based; death certificates	Medical examiner report or medical chart	980	ICD-9-CM ICD-10	800.0-801.9, 803.0-804.9, 850.0-854.1, 950.1-950.3, 959.01, 995.55, S01.0-S01.9, S02.0, S02.1, S02.3, S02.7-S02.9, S04.0, S06.0-S06.9, S07.0, S07.1, S07.8, S07.9, T01.0, T02.0, T04.0, T06.0, T90.1, T90.2, T90.4, T90.5, T90.8, T90.9	78	-	98	-
Shore, 2005, United States ¹⁴	1999	Population-based; inpatient	Medical chart	1002	ICD-9-CM	800-801.9, 803-804.9	-	-	72.1	-
		Population-based; inpatient	Medical chart	1002	ICD-9-CM	850-854.1, 959.01	-	-	89.5	-
		Population-based; inpatient	Medical chart	698	ICD-9-CM	800-801.9, 803-804.9	-	-	71.2	-
		Population-based; inpatient	Medical chart	698	ICD-9-CM	850-854.1, 959.01	-	-	92.0	-
		Population-based; registry	Medical chart	698	ICD-9-CM	800-801.9, 803-804.9	-	-	63.8	-
		Population-based; registry	Medical chart	698	ICD-9-CM	850-854.1, 959.01	-	-	88.1	-

CM = clinical modification; ER = emergency room; ICD = International Classification of Disease; NPV = negative predictive value; PPV = positive predictive value; Sn = sensitivity; Sp = specificity; TBI = traumatic brain injury.

March 30, 2011. Both of these databases are coded by trained health technologists who undergo national standardized training. The ACCS captures all emergency visits and hospital-based clinic visits in the province of Alberta, and contains up to 15 coding fields for diagnoses and five for procedures using ICD-10 codes. The DAD captures information including diagnoses and in-hospital outcomes on all patients discharged from every hospital in Alberta. Since April 1, 2002, the DAD has contained up to 50 coding fields for diagnoses and 20 for procedures. Both databases contain a primary diagnosis coding field that contains the diagnosis (ICD-10 code) responsible for most resource use during the visit or inpatient admission.¹⁵ The rest of the diagnosis codes are classified as secondary diagnoses. Administrative databases were linked with the REDIS cohort data described previously using the first name, last name, date of birth, and medical record number because ethics restricted us from using the more universal personal health number.

ICD-10 codes to be tested for TBI were selected based on the following: (1) an international systematic review of validated ICD codes to study neurological conditions¹¹; (2) a review of the grey literature and relevant websites (including TBI codes used by the US Centers for Disease Control and Prevention for their TBI surveillance program¹ and codes used by the Canadian National Trauma Registry²⁰; and (3) expert consensus (neurologists, epidemiologists, health services researchers, population health experts) from the Canadian Chronic Disease Surveillance System Neurological Conditions Working Group (unpublished) based on the literature reviews provided previously and expert opinion. Because we were only interested in acute presentations of TBI, some codes used to identify TBI were excluded, such as those used to identify certain TBI-related sequelae (e.g. postconcussive syndrome).

A case of TBI was considered to be present if one or more of the specified TBI-related ICD-10 codes was listed in any of the diagnostic coding fields in either of the linked administrative databases. Table 2 shows the ICD-10 codes that were used to identify TBI in the administrative databases. Nine unique case definitions were subsequently tested to see which produced the best case definition for children with TBI using the REDIS cohort data as the reference standard against the administrative data (DAD and ACCS) (Figure 1). Because TBI is an acute presentation, we only tested different case definitions over a period of 1 or 2 years.

STATISTICAL ANALYSIS

Descriptive statistics were used to describe the breakdown of the study population.

Accepting the REDIS chart data as the reference standard, diagnostic accuracy estimates (sensitivity, specificity, positive predictive value [PPV], and NPV) were calculated for each of the nine case definitions. Sensitivity refers to the proportion of TBI cases identified from REDIS that were coded as TBI cases in the administrative data. Specificity refers to the proportion of non-TBI cases identified from REDIS that were coded as non-TBI cases in the administrative data. PPV refers to the proportion of TBI cases identified in the administrative databases that were deemed “true” TBI cases on the basis of REDIS chart review data. NPV refers to the proportion of non-TBI cases identified in the administrative databases that were deemed “true” non-TBI cases on the basis of REDIS chart review data.

Table 2: ICD-10 codes used to identify TBI in administrative data

ICD-10 code	Definition
S02.0	Fracture of vault of skull
S02.1	Fracture of base of skull
S02.3	Fracture of orbital floor
S02.7	Multiple fractures involving skull and facial bone
S02.8	Fractures of other skull and facial bones
S02.9	Fracture of skull and facial bones, part unspecified
S06	Intracranial injury
S06.0	Concussion
S06.1	Traumatic cerebral edema
S06.2	Diffuse brain injury
S06.3	Focal brain injury
S06.4	Epidural hemorrhage
S06.5	Traumatic subdural hemorrhage
S06.6	Traumatic subarachnoid hemorrhage
S06.8	Other intracranial injuries
S06.9	Intracranial injury, unspecified
S07	Crushing injury of the head
S07.0	Crushing injury of the face
S07.1	Crushing injury of the skull
S07.8	Crushing injury of other parts of the head
S07.9	Crushing injury of the head, part unspecified
S09.9	Unspecified injury of head, including injury of ear NOS, face NOS, and nose NOS
T06.0	Injuries of brain and central nerves with injuries of nerves and spinal cord at neck level

NOS = not otherwise specified.

Ethics approval was obtained from the Conjoint Health Research Ethics Board at the University of Calgary.

RESULTS

Study Population

Of the 7101 children identified through REDIS, 1422 (20%) were determined to have TBI and 5679 served as non-TBI cases (Figure 2). After administrative data linkage, 94% (6639/7101) of the original study population was available for analysis, leaving 1343 with TBI and 5296 without TBI (Figure 2). Because of ethical restrictions, we were not able to link our data using the more universal personal health number, which may have contributed to the 6% of patients that we were unable to link. Based on the available linked data, the study population was 56% male (3372/6639), with a mean age of 9.2 years (range, 2.02 days–18.3 years) for males and 8.5 years (range, 2.61 days–18.6 years) for females.

Validation of ICD-10 Coding

The most frequently used code for TBI cases in our population was S09.9—unspecified injury of head, including injury of ear not




Figure 1: Timeline

otherwise specified (NOS), face NOS, and nose NOS (50.5%)—followed by S06.000—concussion without loss of consciousness without open intracranial wound (22.7%) (Table 3). Similarly, 88.9% of the time, the codes used for TBI were found in the primary diagnosis field of the administrative data.

ICD-10 Codes Used in False-Negative TBI Cases

We investigated which ICD-10 codes were used to identify the false-negative cases (n = 406) in our best case definition, to see if we

were missing important codes used to capture children with TBI and to determine if our case definition required further enhancement. The majority of false-negative cases were classified with nonspecific codes identifying superficial injuries, or general trauma, which with other information in the REDIS chart may have appeared to be a possible TBI but was not coded as such in the administrative data.

Identification of Best Case Definition to Identify TBI

Of the nine case definitions assessed (Table 4), the one with the best diagnostic accuracy to capture cases of pediatric TBI for future surveillance work was “1 ER or 1 hospital in 1 year” with a sensitivity of 69.8% (95% confidence interval [CI], 67.3-72.2), specificity of 96.7% (95% CI, 96.2-97.2), PPV of 84.2% (95% CI 82.0-86.3), and NPV of 92.7% (95% CI, 92.0-93.3).

DISCUSSION

This study investigated whether two Alberta-specific administrative databases (hospital and ER databases) could be used to develop a validated case definition for pediatric TBI. The best case definition to accurately identify the largest proportion of children with TBI in these ICD-10-coded data was “1 hospital or 1 ER visit in 1 year.” This means that at least one TBI-related ICD-10 code was captured in the hospitalization or ER database during a 1-year period. Currently, there is not a single agreed-upon case definition for TBI. Some experts recommend excluding ICD-10 code S09.9 from TBI case definitions because it has been shown to inflate incidence and prevalence figures. We chose to include this code in our study because it was the most frequently used (50.5%) to identify TBI in our pediatric study population. Had we excluded this code, we would have missed 1145 cases (80.5%). There is concern that minor brain injury cases may be missed if codes such as S09.9 are excluded, given their ambiguous nature.²¹ It has also been reported that up to 20% of S09.9 diagnoses may be cases of mild TBI.²¹ Therefore, because a large proportion of patients in our study were identified with code S09.9, it is highly plausible that our sample is characterized largely by patients with milder forms of TBI. It is also important to recognize that we did not evaluate false-positive cases occurring




Figure 2: Study population

Table 3: Frequency of ICD-10 codes used in administrative data to identify TBI

Diagnostic code	Description	Frequency	Percent
S02.000	Fracture of vault of skull, closed	19	0.84
S02.001	Fracture of vault of skull, open	3	0.13
S02.100	Fracture of base of skull, closed	14	0.62
S02.101	Fracture of base of skull, open	10	0.44
S02.300	Fracture of orbital floor, closed	7	0.31
S02.800	Fracture of alveolus, closed Includes maxillary and mandibular alveolar ridge	1	0.04
S02.890	Fractures of other unspecified skull and facial bones, closed	6	0.26
S02.900	Fracture of skull and facial bones, part unspecified, closed	18	0.79
S02.901	Fracture of skull and facial bones, part unspecified, open	1	0.04
S06.0	Concussion	85	3.75
S06.000	Concussion without loss of consciousness without open intracranial wound	514	22.65
S06.010	Concussion with brief loss of consciousness without open intracranial wound	126	5.55
S06.020	Concussion with moderate loss of consciousness without open intracranial wound	1	0.04
S06.090	Concussion with loss of consciousness of unspecified duration without open intracranial wound	77	3.39
S06.190	Traumatic cerebral edema with loss of consciousness of unspecified duration without open intracranial wound	1	0.04
S06.200	Diffuse brain injury without loss of consciousness without open intracranial wound	3	0.13
S06.210	Diffuse brain injury with brief loss of consciousness without open intracranial wound	2	0.09
S06.230	Diffuse brain injury with prolonged loss of consciousness with return to pre-existing level of consciousness without open intracranial wound	1	0.04
S06.290	Diffuse brain injury with loss of consciousness of unspecified duration, without open intracranial wound	5	0.22
S06.340	Focal brain injury with prolonged loss of consciousness without return to pre-existing level of consciousness without open intracranial wound	1	0.04
S06.35	Focal brain injury, without open intracranial wound	1	0.04
S06.390	Focal brain injury with loss of consciousness of unspecified duration, without open intracranial wound	2	0.09
S06.4	Epidural hemorrhage	2	0.09
S06.400	Epidural hemorrhage without loss of consciousness without open intracranial wound	15	0.66
S06.401	Epidural hemorrhage without loss of consciousness with open intracranial wound	1	0.04
S06.420	Epidural hemorrhage with moderate loss of consciousness without open intracranial wound	1	0.04
S06.5	Traumatic subdural hemorrhage	8	0.35
S06.500	Traumatic subdural hemorrhage without loss of consciousness without open intracranial wound	13	0.57
S06.510	Traumatic subdural hemorrhage with brief loss of consciousness without open intracranial wound	5	0.22
S06.530	Traumatic subdural hemorrhage with prolonged loss of consciousness with return to pre-existing level of consciousness without open intracranial wound	1	0.04
S06.590	Traumatic subdural hemorrhage with loss of consciousness of unspecified duration without open intracranial wound	2	0.09
S06.591	Traumatic subdural hemorrhage with loss of consciousness of unspecified duration with open intracranial wound	1	0.04
S06.6	Traumatic subarachnoid hemorrhage	2	0.09
S06.600	Traumatic subarachnoid hemorrhage without loss of consciousness without open intracranial wound	3	0.13
S06.610	Traumatic subarachnoid hemorrhage with brief loss of consciousness without intracranial wound	1	0.04
S06.630	Traumatic subarachnoid hemorrhage with prolonged loss of consciousness with return to pre-existing level of consciousness without open intracranial wound	1	0.04
S06.640	Traumatic subarachnoid hemorrhage with prolonged loss of consciousness without return to pre-existing level of consciousness without open intracranial wound	1	0.04
S06.690	Traumatic subarachnoid hemorrhage with loss of consciousness of unspecified duration without open intracranial wound	2	0.09
S06.691	Traumatic subarachnoid hemorrhage with loss of consciousness of unspecified duration with open intracranial wound	1	0.04
S06.800	Other intracranial injuries without loss of consciousness without open intracranial wound	10	0.44
S06.810	Other intracranial injuries with brief loss of consciousness without open intracranial wound	7	0.31
S06.890	Other intracranial injuries with loss of consciousness of unspecified duration without open intracranial wound	3	0.13
S06.9	Intracranial injury, unspecified, includes brain injury NOS, excludes head injury NOS	16	0.71

Table 3: (Continued)

Diagnostic code	Description	Frequency	Percent
S06.900	Intracranial injury, unspecified without loss of consciousness without open intracranial wound	87	3.83
S06.910	Intracranial injury, unspecified with brief loss of consciousness without open intracranial wound	19	0.84
S06.990	Intracranial injury, unspecified with loss of consciousness of unspecified duration without open intracranial wound	23	1.01
S06.991	Intracranial injury, unspecified without loss of consciousness with open intracranial wound	1	0.04
S09.9	Unspecified injury of the head, including injury of ear NOS, face NOS, and nose NOS	1145	50.46
Total frequency		2269	100

NOS = not otherwise specified.

*Children with TBI in our cohort (true-positive cases) were identified to have the codes listed here.

as a result of the use of this code in our sample. As such, it is important to recognize that the use of this nonspecific code may have resulted in the identification of more false-positive cases, which is a limitation.

Our recommended case definition, “1 hospital or 1 ER visit over 1 year” was selected because it is important to prioritize sensitivity over other measures when the goal is to identify all persons with a certain characteristic (such as TBI) in a population.²² This is relevant to our study because the goal is to use the best case definition for disease surveillance. Although the sensitivity is not perfect, the specificity is very good and, as such, the definition is still valuable for allowing the identification of cohorts for future studies examining a variety of health care issues such as the trajectory of cases through the health care system. This ICD-10 coding validation is important because most countries are now using ICD-10 data to code for morbidity and mortality.⁹

The other tested case definitions revealed varying results. The case definition looking at the DAD alone missed the majority of TBI cases (sensitivity, 3.7-4.2%), but was still associated with excellent specificity (up to 99.8%) and PPV (up to 84.7%). Although this group of hospitalized patients may be good for accurately defining cases for hospital-based outcomes or follow-up studies, this is a poor choice for surveillance. Using the hospitalization data alone may also introduce selection bias, because only the more severe cases tend to be admitted. The case

definition identifying those from the ER alone may also be good for surveillance, but is likely missing the more severe cases; thus, it is important to use a case definition that captures children from both databases. Incorporating both of these databases for national surveillance of pediatric TBI is feasible in Canada because both hospitalization and ER visits are collected prospectively by provincial governments. It would be time-consuming and prohibitively expensive for a clinical network to develop its own pediatric TBI surveillance program. Administrative data are more feasible and are readily available. In the future, electronic medical records or electronic health records may become another source for TBI surveillance, but alternative case definitions incorporating both free text and ICD codes will then need to be developed. Numerous electronic medical record and electronic health record vendors exist, rendering this option more challenging.

Although there are no cutoffs or standard criteria that specify an optimal sensitivity, specificity, PPV, or NPV for case definitions, the validity of our case definition for pediatric TBI is comparable to other conditions. A systematic review looking at validated case definitions for neurological conditions found low sensitivity for Alzheimer’s disease and dementia, Parkinson’s disease, and spinal cord injury.¹¹ Similarly, two separate studies validating ICD-9 codes for depression found low sensitivity, ranging from 18.1% to 34.0%.^{23,24} Considering the results of validation studies for different neurological and mental health

Table 4: TBI validation results for case definitions including hospital or ER visits

TBI case definition	True positives	False negatives	False positives	True negatives	Sensitivity % (95% CI)	Specificity % (95% CI)	PPV % (95% CI)	NPV % (95% CI)
1 hospital, 1 ER, 1 year*	937	406	176	5120	69.8 (67.3-72.2)	96.7 (96.2-97.2)	84.2 (82.0-86.3)	92.7 (92.0-93.3)
1 hospital, 1 ER, 2 years	952	391	266	5030	70.9 (68.5-73.3)	95.0 (94.4-95.6)	78.2 (75.8-80.5)	92.8 (92.1-93.5)
1 hospital, 1 year	50	1293	9	5287	3.7 (2.7-4.7)	99.8 (99.7-99.9)	84.7 (75.6-93.9)	80.3 (79.4-81.3)
1 hospital, 2 years	57	1286	32	5264	4.2 (3.2-5.3)	99.4 (99.2-99.6)	64.0 (54.1-74.0)	80.4 (79.4-81.3)
1 ER, 1 year	932	411	174	5122	69.4 (66.9-71.9)	96.7 (96.2-97.2)	84.3 (82.1-86.4)	92.6 (91.9-93.3)
1 ER, 2 years	946	397	263	5033	70.4 (68.0-72.9)	95.0 (94.4-95.6)	78.2 (75.9-80.6)	92.7 (92.0-93.4)
1 hospital, 2 ERs, 1 year	145	1198	16	5280	10.8 (9.1-12.5)	99.7 (99.6-99.8)	90.1 (85.4-94.7)	81.5 (80.6-82.5)
1 hospital, 2 ERs, 2 years	165	1178	36	5260	12.3 (10.5-14.0)	99.3 (99.1-99.5)	82.1 (76.8-87.4)	81.7 (80.8-82.6)
Any TBI diagnosis	979	364	637	4659	72.9 (70.5-75.3)	88.0 (87.1-88.8)	60.6 (58.2-63.0)	92.8 (92.0-93.5)

CI = confidence interval; ER = emergency room; NPV = negative predictive value; PPV = positive predictive value; TBI = traumatic brain injury.

*This case definition means that at least one TBI-related International Classification of Diseases, 10th revision, code was captured in the hospitalization or ER database during a 1-year period.

conditions, ours has high validity and should be considered for use in pediatric TBI surveillance.

There are several limitations to this study. First, our reference standard was based on a pediatric ER electronic chart review only. It is a concern that misdiagnosis or incomplete documentation of clinical data in a more limited electronic record may affect results. However, when children are admitted to the emergency room, they often receive a more comprehensive workup acutely than if solely seen in community-based locations. In addition, these emergency department electronic charts were reviewed by both a trained research assistant and a pediatric neurologist with TBI expertise; therefore, we postulate that this reference standard is most likely reflecting the “true” diagnosis. We also further confirmed the diagnosis by examining the percentage of children who were not deemed to have TBI upon telephone follow-up (4%) and who were not deemed to have TBI upon a clinic visit (0%). Misclassification bias is a common concern with administrative data; however, one of the primary purposes of a validation study is to examine the validity of these codes and to determine how to best identify children with TBI, so that we are capturing the most accurate picture.

To increase the likelihood of capturing children who may have experienced a TBI (in view of the low incidence of TBI based on epidemiological studies), our enriched population included children with primarily trauma, musculoskeletal, or central nervous system issues. As such, the incidence of TBI in our cohort was higher than expected, and as a result, could have influenced our predictive values (e.g. higher prevalence = higher PPV).

Second, we were not able to assess the severity of TBI cases in our population using a gold standard measure. Further research evaluating TBI severity using the Glasgow Coma Scale, the Abbreviated Injury Scale, or other measures would increase our understanding of which types of TBI are seen in different settings and how case definition accuracy may vary by TBI severity subtype. Despite our inability to determine severity, we initially hypothesized that the majority of cases seen in the ER or hospitalized would be the moderate to severe cases. However, considering the high proportion of children identified in our study with the nonspecific code S09.9 (>80%), it is highly likely that many of the children in our sample had milder injuries. The frequent use of this nonspecific code could also relate to the challenge of accurately identifying symptoms of TBI in younger children.

In addition, we were not able to obtain physician claims data that capture all visits, whether hospital or community-based; therefore, it is possible that we are missing a proportion of children with mild TBI who were seen in the community by their family physician or a walk-in-clinic. However, ACCS captures both emergency visits and hospital-based clinic visits; therefore, we are likely to have captured at least the more symptomatic mild TBI cases.

Third, this study population was selected from one hospital in a large city; therefore, the results may not be generalizable to other regions, including rural areas. In fact, it has been stressed in the literature that case definitions validated in one specific area or age group cannot necessarily be applied to other cohorts if you want to ensure accuracy.²⁵ However, in Canada, professional ICD coders receive standardized national training (17). Therefore, results across different provincial administrative databases should be similar. Generalizing results internationally would likely be difficult because it has been shown there is coding variation between countries.²⁶

In conclusion, we identified that, for surveillance purposes, the case definition “1 hospital or 1 ER visit over 1 year” is best at capturing children with TBI in the two administrative databases using ICD-10 coding. However, there is a need to validate TBI coding in physician claims data to enhance our proposed case definition and to get a more comprehensive picture of all TBI cases whether mild, moderate, or severe. We would also recommend that more efforts be made to educate coders on appropriate use of TBI-related ICD codes to better identify children with the various TBI subtypes, and that physicians receive more training in medical school and residency on the importance of careful chart documentation.

ACKNOWLEDGMENTS AND FUNDING

None.

DISCLOSURES

HQ has received a grant from AIHS and salary support from Alberta Innovates-Health Solutions. NJ is a Principal Investigator for Hotchkiss Brain Institute, the Canada Research Chair Program, and Alberta Innovates Health Solutions; has received a research grant from Pfizer (a shared grant from the University and Pfizer); and has received salary support from a government agency and from principal health research funding sources paid to her university. The remaining authors have nothing to disclose.

REFERENCES

1. Faul M, Xu LWald M, Coronado V. Traumatic brain injury in the United States: emergency department visits, hospitalizations and deaths 2002-2006. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2010.
2. Schneier AJ, Shields BJ, Hostetler SG, Xiang H, Smith GA. Incidence of pediatric traumatic brain injury and associated hospital resource utilization in the United States. *Pediatrics*. 2006;118:483-92.
3. Canadian Institute for Health Information. The burden of neurological diseases, disorders and injuries in Canada. Ottawa: CIHI; 2007.
4. Nguyen R, Fiest K, McChesney J, et al. The international incidence of traumatic brain injury: a systematic review and meta-analysis. *Can J Neurol Sci*. 2016;43:774-85.
5. Gordon KE, Dooley JM, Wood EP. Descriptive epidemiology of concussion. *Pediatr Neurol*. 2006;34:376-8.
6. Willer B, Dumas J, Hutson A, Leddy J. A population based investigation of head injuries and symptoms of concussion of children and adolescents in schools. *Inj Prevent*. 2004;10:144-8.
7. Toga AW, Thompson PM, Sowell ER. Mapping brain maturation. *Trends Neurosci*. 2006;29:148-59.
8. Quan H, Khan N, Hemmelgarn BR, et al. Validation of a case definition to define hypertension using administrative data. *Hypertension*. 2009;54:1423-8.
9. Jette N, Reid AY, Quan H, Hill MD, Wiebe S. How accurate is ICD coding for epilepsy? *Epilepsia*. 2010;51:62-9.
10. World Health Organization. International Classification of Diseases. Available at: <http://www.who.int/classifications/icd/en/>.
11. St Germaine-Smith C, Metcalfe A, Pringsheim T, et al. Recommendations for optimal ICD codes to study neurologic conditions: a systematic review. *Neurology*. 2012;79:1049-55.
12. Bazarian JJ, Veazie P, Mookerjee S, Lerner EB. Accuracy of mild traumatic brain injury case ascertainment using ICD-9 codes. *Acad Emerg Med*. 2006;13:31-8.
13. Rodriguez SR, Mallonee S, Archer P, Gofton J. Evaluation of death certificate-based surveillance for traumatic brain injury—Oklahoma 2002. *Public Health Rep*. 2006;121:282-9.
14. Shore AD, McCarthy ML, Serpi T, Gertner M. Validity of administrative data for characterizing traumatic brain injury-related hospitalizations. *Brain Inj*. 2005;19:613-21.

15. Jette N, Quan H, Hemmelgarn B, et al. The development, evolution, and modifications of ICD-10: challenges to the international comparability of morbidity data. *Med Care*. 2010;48:1105-10.
16. US Centers for Disease Control and Prevention. International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM). Available at: <http://www.cdc.gov/nchs/icd/icd10cm.htm>.
17. Reid AY, St Germaine-Smith C, Liu M, et al. Development and validation of a case definition for epilepsy for use with administrative health data. *Epilepsy Res*. 2012;102:173-9.
18. Statistics Canada. 2006 Community Profiles. Available at: <http://www.census2006.com/census-recensement/2006/dp-pd/prof/92-591/index.cfm?Lang=E>.
19. Menon DK, Schwab K, Wright DW, Maas AI. Position statement: definition of traumatic brain injury. *Arch Phys Med Rehabil*. 2010;91:1637-40.
20. Canadian Institute for Health Information. National Trauma Registry. Available at: http://www.cihi.ca/CIHI-ext-portal/internet/en/document/typesofcare/specializedservices/traumaandinjuries/services_ntr.
21. Chen AY, Colantonio A. Defining neurotrauma in administrative data using the International Classification of Diseases Tenth Revision. *Emerg Themes Epidemiol*. 2011;8:4.
22. Chubak J, Pocobelli G, Weiss NS. Tradeoffs between accuracy measures for electronic health care data algorithms. *J Clin Epidemiol*. 2012;65:343-9.
23. Noyes K, Liu H, Lyness JM, Friedman B. Medicare beneficiaries with depression: comparing diagnoses in claims data with the results of screening. *Psychiatr Serv*. 2011;62:1159-66.
24. Singh JA. Accuracy of Veterans Affairs databases for diagnoses of chronic diseases. *Prev Chronic Dis*. 2009;6:A126.
25. Benchimol EI, Manuel DG, To T, et al. Development and use of reporting guidelines for assessing the quality of validation studies of health administrative data. *J Clin Epidemiol*. 2011;64:821-829.
26. Drosler SE, Romano PS, Tancredi DJ, Klazinga NS. International comparability of patient safety indicators in 15 OECD member countries: a methodological approach of adjustment by secondary diagnoses. *Health Serv Res*. 2012;47:275-92.