



COMPOSITIO MATHEMATICA

Freeness and multirestriction of hyperplane arrangements

Mathias Schulze

Compositio Math. **148** (2012), 799–806.

[doi:10.1112/S0010437X12000164](https://doi.org/10.1112/S0010437X12000164)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY



Freeness and multirestriction of hyperplane arrangements

Mathias Schulze

ABSTRACT

Generalizing a result of Yoshinaga in dimension three, we show that a central hyperplane arrangement in 4-space is free exactly if its restriction with multiplicities to a fixed hyperplane of the arrangement is free and its reduced characteristic polynomial equals the characteristic polynomial of this restriction. We show that the same statement holds true in any dimension when imposing certain tameness hypotheses.

1. Introduction

One of the main themes in the study of hyperplane arrangements is the relation of combinatorial and algebraic or topological data associated with arrangements. In the present article we are concerned with the relation of characteristic polynomials of restrictions on the combinatorial (and topological) side and freeness (of the module of logarithmic differentials/derivations) on the algebraic side. One of the main open problems in this field, Terao’s conjecture, states that freeness is a combinatorial property for simple arrangements.

Let \mathcal{A} be a simple central arrangement of m hyperplanes in an ℓ -dimensional vector space V over a field \mathbb{K} of characteristic 0. Its intersection lattice $L_{\mathcal{A}}$, ordered by reverse inclusion, is a geometric lattice with rank function equal to the codimension in V . Note that V is the unique minimal element in $L_{\mathcal{A}}$. The Möbius function of $L_{\mathcal{A}}$ is the map

$$\mu : L_{\mathcal{A}} \rightarrow \mathbb{Z}$$

defined recursively by the equalities

$$\begin{aligned} \mu(V) &= 1, \\ \forall X \in L_{\mathcal{A}} \setminus \{V\} : \sum_{X \geq Y \in L_{\mathcal{A}}} \mu(Y) &= 0. \end{aligned} \tag{1}$$

It determines the characteristic polynomial of \mathcal{A} , defined as

$$\chi(\mathcal{A}, t) = \sum_{X \in L_{\mathcal{A}}} \mu(X) t^{\dim X}.$$

By (1), $\chi(\mathcal{A}, 1) = 0$ and the quotient

$$\chi_0(\mathcal{A}, t) = \chi(\mathcal{A}, t)/(t - 1) \tag{2}$$

Received 29 April 2011, accepted in final form 18 October 2011, published online 22 March 2012.

2010 Mathematics Subject Classification 14N20 (primary), 16W25, 13D40 (secondary).

Keywords: hyperplane arrangement, free divisor.

The author gratefully acknowledges support by the ‘SQuaREs’ program of American Institute of Mathematics. He would like to thank Graham Denham, Hal Schenck, Max Wakefield, and Uli Walther for helpful discussions. This journal is © [Foundation Compositio Mathematica](http://www.compositio-mathematica.org/) 2012.

is called the reduced characteristic polynomial. For complex arrangements, it is equivalent to the Poincaré polynomial $\pi(\mathcal{A}, t) = t^\ell \chi(\mathcal{A}, -t^{-1})$ of the complement of the arrangement by [OS80].

The complex $\Omega^\bullet(\mathcal{A})$ of logarithmic differential forms along a multiarrangement \mathcal{A} with multiplicities $\mathbf{k} = (k_H)_{H \in \mathcal{A}}$ consists of the S -modules

$$\Omega^p(\mathcal{A}) = \left\{ \omega \in \frac{1}{Q} \Omega_V^p \mid \forall H \in \mathcal{A} : \frac{d\alpha_H}{\alpha_H^{k_H}} \wedge \omega \in \frac{1}{Q} \Omega_V^{p+1} \right\},$$

and it is closed under exterior product. The module $\Omega^1(\mathcal{A})$ and the S -module of logarithmic derivations

$$D(\mathcal{A}) = D_1(\mathcal{A}) = \{ \delta \in D_V \mid \forall H \in \mathcal{A} : \delta(\alpha_H) \in \alpha_H^{k_H} S \},$$

where $D_V = \text{Der}_{\mathbb{K}}(S, S)$ denotes the S -module of polynomial vector fields on V , are mutually S -dual. In particular, $\Omega^1(\mathcal{A})$ and $D(\mathcal{A})$ are reflexive S -modules. If one of them is a free S -module then \mathcal{A} is called a free arrangement.

Consider now a fixed hyperplane $H \in \mathcal{A}$ and the multiarrangement restriction \mathcal{A}^H of \mathcal{A} to H . If \mathcal{A} is defined by $Q = \prod_{H' \in \mathcal{A}} \alpha_{H'} \in S = \text{Sym } V^*$, and H by $\alpha_H \in V^*$, then \mathcal{A}^H is defined by

$$Q_H = \frac{Q}{\alpha_H} \Big|_H \in S' = \text{Sym } H^*.$$

For any $H' \in \mathcal{A} \setminus \{H\}$ the natural multiplicity of $H \cap H' \in \mathcal{A}^H$ is

$$k_{H \cap H'} = \#\{H'' \in \mathcal{A} \setminus \{H\} \mid H \cap H' = H \cap H''\}.$$

In order to formulate our main result, we extend the well-known notion of tameness for arrangements.

DEFINITION 1. We call \mathcal{A} tame if

$$\forall p = 1, \dots, \ell - 1 : \text{pdim } \Omega^p(\mathcal{A}) \leq p. \tag{3}$$

If (3) holds for $\Omega^p(\mathcal{A})$ replaced by $D^p(\mathcal{A})$, we call \mathcal{A} dually tame. We speak of weak tameness, in both cases, if (3) holds for $p = 1$.

Note that all 3-arrangements are both tame and dually tame. For locally free arrangements, such as, for example, generic arrangements, weak tameness coincides with ordinary tameness by [MS01, Corollary 5.4].

Yoshinaga [Yos05, Corollary 3.3] proved our following main result for $\ell = 3$. For the proof we shall use his result [Yos04, Theorem 2.2] for $\ell > 3$ and extend his approach given in [Yos05].

THEOREM 2. Assume that $\ell \leq 4$ or that \mathcal{A} is weakly (dually) tame. Then \mathcal{A} is free if and only if \mathcal{A}^H is free with exponents d'_2, \dots, d'_ℓ and

$$\chi_0(\mathcal{A}, t) = \prod_{k=2}^{\ell} (t - d'_k). \tag{4}$$

The ‘only if’ part holds true for all $\ell \geq 3$ by [Zie89b, Theorem 11] and Terao’s factorization theorem [Ter81].

Assuming freeness of \mathcal{A}^H , condition (4) has the following geometrical interpretation. Let \mathcal{A}_s be the restriction of \mathcal{A} to the affine space H_s defined by $\alpha_H = s$. By the natural \mathbb{K}^* -action, the arrangements $\mathcal{A}_s, s \neq 0$, form a trivial family of non-central simple arrangements and \mathcal{A} can be

identified with the cone of \mathcal{A}_s , for any $s \neq 0$. Then [OT92, Proposition 2.51] shows that

$$\forall s \neq 0 : \chi(\mathcal{A}_s, t) = \chi_0(\mathcal{A}, t).$$

On the other hand, the right-hand side of (4) equals the characteristic polynomial $\chi(\mathcal{A}^H)$ of the multiarrangement $\mathcal{A}^H = \mathcal{A}_0$ as defined in [ATW07]. So condition (4) can be interpreted as the family \mathcal{A}_s being trivial on the level of characteristic polynomials. However, we do not know a concept of characteristic polynomial that covers both the central multiarrangement case and the non-central simple arrangement case.

Whether our result has any implications on Terao’s conjecture is, *a priori*, unclear. For instance, Ziegler [Zie89b, Proposition 10] showed that exponents of multiarrangements are not combinatorial in general.

2. Multirestriction of logarithmic forms

Our starting point is a construction from [Yos05]. Let M^\bullet denote the image of the restriction map

$$\text{res}_H^\bullet : \Omega^\bullet(\mathcal{A}) \rightarrow \Omega^\bullet(\mathcal{A}^H), \quad \frac{d\alpha_H}{\alpha_H} \wedge \eta + \sigma \mapsto \sigma|_H, \tag{5}$$

and let C^\bullet be its cokernel. Then res_H^\bullet factors through $d\alpha_H/\alpha_H \wedge - : \Omega^\bullet \rightarrow d\alpha_H/\alpha_H \wedge \Omega^\bullet$ into the residue map

$$\frac{d\alpha_H}{\alpha_H} \wedge \Omega^\bullet(\mathcal{A}) \rightarrow \Omega^\bullet(\mathcal{A}^H). \tag{6}$$

As the complex $(\Omega^\bullet(\mathcal{A}), d\alpha_H/\alpha_H \wedge -)$ is exact by [OT92, Proposition 4.86],

$$\frac{1}{Q} \Omega^\ell = \Omega^\ell(\mathcal{A}) = \frac{d\alpha_H}{\alpha_H} \wedge \Omega^{\ell-1}(\mathcal{A}) = \frac{d\alpha_H}{\alpha_H} \wedge \frac{\alpha_H}{Q} \Omega^{\ell-1}. \tag{7}$$

By definition,

$$\Omega^{\ell-1}(\mathcal{A}^H) = \frac{1}{Q_H} \Omega_H^{\ell-1}, \tag{8}$$

and it follows using (7) that $C^{\ell-1} = 0$. Thus, $C^p \neq 0$ only for $1 \leq p \leq \ell - 2$.

PROPOSITION 3. *If \mathcal{A}^H is free and $C^1 = 0$ then $C^\bullet = 0$. In particular, freeness of \mathcal{A} implies $C^\bullet = 0$.*

Proof. If \mathcal{A}^H is free then, by Saito’s criterion [Zie89b, Theorem 8] and (8),

$$\Omega^{\ell-1}(\mathcal{A}^H) = \bigwedge^{\ell-1} \Omega^1(\mathcal{A}^H).$$

So [OT92, the proof of Proposition 4.81] shows that

$$\forall p = 0, \dots, \ell - 1 : \Omega^p(\mathcal{A}^H) = \bigwedge^p \Omega^1(\mathcal{A}^H).$$

As restriction commutes with exterior product, these equalities and the surjectivity of res_H^1 implies the surjectivity of res_H^\bullet , and hence $C^\bullet = 0$ as claimed.

Now the second statement follows from [Yos05, Theorem 2.5] □

By [Yos05, Theorem 2.5, Corollary 2.6] and Terao’s factorization theorem, the ‘if’ part of the statement of Theorem 2 is equivalent to

$$(4) \text{ implies } C^1 = 0 \text{ for free } \mathcal{A}^H.$$

The following result is a first step in this direction.

PROPOSITION 4. *If \mathcal{A}^H is free and (4) holds then $\dim(C^p) \leq \ell - 4$ for all $p \in \mathbb{Z}$.*

Proof. By [Yos05, Theorem 2.8], the Poincaré series

$$P(M^\bullet, x, y) = \sum_{p=0}^{\ell-1} P(M^p, x)y^p \tag{9}$$

of M^\bullet and $\Phi(\mathcal{A}, x, y)$ of $\Omega^\bullet(\mathcal{A})$ are related by the equality

$$\Phi(\mathcal{A}, x, y) = \frac{x + y}{x(1 - x)} P(M^\bullet, x, y). \tag{10}$$

Assuming freeness of \mathcal{A}^H , Ziegler’s dual version [Zie89a, Definition-Theorem 3.21] of [ATW07] and [ST87] shows that

$$\chi(\mathcal{A}^H, t) = \lim_{x \rightarrow 1} P(\Omega^\bullet(\mathcal{A}^H), x, t(1 - x) - 1), \tag{11}$$

$$\chi(\mathcal{A}, t) = \lim_{x \rightarrow 1} \Phi(\mathcal{A}, x, t(1 - x) - 1). \tag{12}$$

As in [Yos05, (9)], combining (2), (9), (10), and (12) yields

$$\begin{aligned} \chi_0(\mathcal{A}, t) &= \frac{1}{t - 1} \lim_{x \rightarrow 1} \frac{x + t(1 - x) - 1}{x(1 - x)} \sum_{p=0}^{\ell-1} P(M^p, x)(t(1 - x) - 1)^p \\ &= \lim_{x \rightarrow 1} P(M^\bullet, x, t(1 - x) - 1), \end{aligned} \tag{13}$$

and hence, applying (11) and (13) to the exact sequence

$$0 \rightarrow M^\bullet \rightarrow \Omega^\bullet(\mathcal{A}^H) \rightarrow C^\bullet \rightarrow 0, \tag{14}$$

we obtain

$$\chi(\mathcal{A}^H, t) - \chi_0(\mathcal{A}, t) = \lim_{x \rightarrow 1} P(C^\bullet, x, t(1 - x) - 1). \tag{15}$$

Recall that $C^0 = 0 = C^{\ell-1}$ and set $P^p(x) = (-1)^p P(C^p, x)$ for $p = 0, \dots, \ell - 1$. Assuming (4), the left-hand side of (15) vanishes, and we compute

$$\begin{aligned} 0 &= \lim_{x \rightarrow 1} P(C^\bullet, x, (-t)(1 - x) - 1) \\ &= \lim_{x \rightarrow 1} \sum_{p=0}^{\ell-2} (t(1 - x) + 1)^p P^p(x) \\ &= \lim_{x \rightarrow 1} \sum_{p=0}^{\ell-2} \sum_{r=0}^p \binom{p}{r} t^r (1 - x)^r P^p(x) \\ &= \sum_{r=0}^{\ell-2} t^r \lim_{x \rightarrow 1} \sum_{p=r}^{\ell-2} \binom{p}{r} (1 - x)^r P^p(x). \end{aligned} \tag{16}$$

Considering (16) as an equality of polynomials in $(\mathbb{K}[x])[t]$ shows that

$$\lim_{x \rightarrow 1} \sum_{p=r}^{\ell-2} \binom{p}{r} (1-x)^r P^p(x) = 0, \quad r = 0, \dots, \ell - 2.$$

In particular, for each $k = 0, \dots, \ell - 2$,

$$\lim_{x \rightarrow 1} \sum_{p=r}^{\ell-2} \binom{p}{r} (1-x)^k P^p(x) = 0, \quad r = 0, \dots, k. \tag{17}$$

For $k = \ell - 2$, the matrix of the system of linear equations (17) has full rank. By induction on the pole order of $P^\bullet(x)$, it follows that

$$\lim_{x \rightarrow 1} (1-x)^m P^p(x) = 0, \quad p = 1, \dots, \ell - 2, \tag{18}$$

holds for $m = \ell - 2$. Thus, (17) for $k = \ell - 3$ can be considered as a system of linear equations satisfied by $(\lim_{x \rightarrow 1} (x - 1)^{\ell-3} P^p(x))_{p=1, \dots, \ell-2}$ with matrix

$$\begin{pmatrix} \binom{1}{0} & \binom{2}{0} & \binom{3}{0} & \cdots & \binom{\ell-3}{0} & \binom{\ell-2}{0} \\ \binom{1}{1} & \binom{2}{1} & \binom{3}{1} & \cdots & \binom{\ell-3}{1} & \binom{\ell-2}{1} \\ 0 & \binom{2}{2} & \binom{3}{2} & \cdots & \binom{\ell-3}{2} & \binom{\ell-2}{2} \\ 0 & 0 & \binom{3}{3} & \cdots & \binom{\ell-3}{3} & \binom{\ell-2}{3} \\ \vdots & & \ddots & \ddots & \vdots & \vdots \\ 0 & \dots & \dots & 0 & \binom{\ell-3}{\ell-3} & \binom{\ell-2}{\ell-3} \end{pmatrix}. \tag{19}$$

Using the relation $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$, one can eliminate the lower subdiagonal in (19) by appropriate row operations, keeping all diagonal elements in \mathbb{K}^* . Thus, (19) is invertible and (18) also holds for $m = \ell - 3$. This means that $\ell - 4$ is an upper bound for the pole order of $P(C^p, x)$ at $x = 1$ which equals $\dim C^p$, for all $p = 1, \dots, \ell - 2$. The claim follows. \square

The following statement for $\ell = 3$ also follows from [Yos05, Theorem 3.2].

COROLLARY 5. *Assume that \mathcal{A}^H is free and (4) holds. For $\ell = 3$, $\dim_{\mathbb{K}} C^1 = 0$ and, for $\ell = 4$, $\dim_{\mathbb{K}} C^1 = \dim_{\mathbb{K}} C^2 < \infty$.*

3. Duality of multirestriction for forms and derivations

To understand $C^{\ell-2}$ and deal with the case $\ell = 4$, the following trivial lemma will be useful. For simple arrangements, it is well known. We fix a coordinate system $z_1, \dots, z_\ell \in V^*$ and set $dz = dz_1 \wedge \cdots \wedge z_\ell$.

LEMMA 6. Let \mathcal{A} be a multiarrangement in V with multiplicities $\mathbf{k} = (k_H)_{H \in \mathcal{A}}$ and defining equation $Q \in S$.

- (a) The inner product induces a S -bilinear pairing $D(\mathcal{A}) \times \Omega^p(\mathcal{A}) \rightarrow \Omega^{p-1}(\mathcal{A})$.
- (b) $\langle -, dz \rangle$ induces an isomorphism of graded S -modules $D(\mathcal{A}) \cong Q\Omega^{\ell-1}(\mathcal{A})$.

Proof. (a) Let $\omega/Q \in \Omega^p(\mathcal{A})$ and $\xi \in D(\mathcal{A})$. Then $d\alpha_H \wedge \omega \in \alpha_H^{k_H} \Omega_V^{p+1}$ and $\xi(\alpha_H) \in \alpha_H^{k_H} S$, which implies that

$$d\alpha_H \wedge \langle \xi, \omega \rangle = \langle \xi, d\alpha_H \wedge \omega \rangle + \xi(\alpha_H)\omega \in \alpha_H^{k_H} \Omega_V^p. \tag{20}$$

Thus, $\langle \xi, \omega/Q \rangle = \langle \xi, \omega \rangle/Q \in \Omega^{p-1}(\mathcal{A})$.

- (b) By part (a), each $\xi \in D(\mathcal{A})$ defines an S -linear map

$$\xi = \langle \xi, - \rangle : S \frac{dz}{Q} = \Omega^\ell(\mathcal{A}) \rightarrow \Omega^{\ell-1}(\mathcal{A})$$

which associates to ξ an element $\langle \xi, dz \rangle/Q = \langle \xi, dz/Q \rangle \in \Omega^{\ell-1}(\mathcal{A})$. One computes that

$$\frac{d\alpha_H}{\alpha_H^{k_H}} \wedge \left\langle \xi, \frac{dz}{Q} \right\rangle = \frac{\xi(\alpha_H)}{\alpha_H^{k_H}} dz, \tag{21}$$

which shows that $\xi \mapsto \langle \xi, dz/Q \rangle$ is an isomorphism. □

For the simple arrangement \mathcal{A} , consider the module of derivations that are logarithmic both along \mathcal{A} and along the level sets of α_H ,

$$D_H(\mathcal{A}) = D(\mathcal{A}) \cap D(\alpha_H) = \{ \delta \in D(\mathcal{A}) \mid \delta(\alpha_H) = 0 \}.$$

Then, by [Zie89b, Theorem 11], there is a restriction map

$$\text{res}_H : D_H(\mathcal{A}) \rightarrow D(\mathcal{A}^H),$$

and we denote by M its image and by C its cokernel.

PROPOSITION 7. There is an isomorphism $C[m-1] \cong C^{\ell-2}$ of graded S' -modules.

Proof. By Lemma 6(b), there is an isomorphism of graded S -modules

$$\langle -, dz \rangle : D(\mathcal{A}) \rightarrow Q\Omega^{\ell-1}(\mathcal{A}). \tag{22}$$

Using the Euler derivation $\chi \in D(\mathcal{A})$, one can decompose

$$D(\mathcal{A}) = S\chi \oplus D_H(\mathcal{A}). \tag{23}$$

Then (21) shows that $\langle D_H(\mathcal{A}), dz/Q \rangle \subset \Omega^{\ell-1}(\mathcal{A})$ is the kernel of $d\alpha_H/\alpha_H \wedge -$. Using that the complex $(\Omega^\bullet(\mathcal{A}), d\alpha_H/\alpha_H \wedge -)$ is exact by [OT92, Proposition 4.86], we conclude that

$$\left\langle D_H(\mathcal{A}), \frac{dz}{Q} \right\rangle = \frac{d\alpha_H}{\alpha_H} \wedge \Omega^{\ell-2}(\mathcal{A}).$$

As $M^{\ell-2}$ is the residue of the latter module along H , (22) induces an isomorphism

$$M = D_H(\mathcal{A})|_H \cong Q_H M^{\ell-2}. \tag{24}$$

Again by Lemma 6(b), there is an isomorphism of S' -modules

$$D(\mathcal{A}^H) \cong Q_H \Omega^{\ell-2}(\mathcal{A}^H). \tag{25}$$

The claim follows by combining (24) and (25). □

Assume that $\ell = 4$, that \mathcal{A}^H is free, and that (4) holds. By Corollary 5, Proposition 7, and homogeneity, the cokernel C of res_H is supported at $0 \in H$ only. Therefore, \mathcal{A} is locally free along H by [Yos04, Theorem 2.1]. Then [Yos04, Theorem 2.2] yields the following result.

THEOREM 8. *The statement of Theorem 2 holds true for $\ell = 4$.*

4. Tameness hypotheses on forms and derivations

In order to apply the (weak) tameness hypothesis to (5) and (6), we need the following trivial result.

LEMMA 9. *There is a direct sum decomposition*

$$\Omega^1(\mathcal{A}) \cong S \frac{d\alpha_H}{\alpha_H} \oplus \frac{d\alpha_H}{\alpha_H} \wedge \Omega^1(\mathcal{A}).$$

Proof. Equality (20) with $\xi = \chi$ shows that

$$0 \rightarrow S \frac{d\alpha_H}{\alpha_H} \rightarrow \Omega^1(\mathcal{A}) \rightarrow \frac{d\alpha_H}{\alpha_H} \wedge \Omega^1(\mathcal{A}) \rightarrow 0$$

is split exact, cf. [OT92, the proof of Proposition 4.86]. □

To see the relation with the direct sum decomposition (23), consider the split exact sequence

$$0 \longrightarrow D_H(\mathcal{A}) \longrightarrow D(\mathcal{A}) \xrightarrow{\varphi} S \longrightarrow 0 \tag{26}$$

where

$$\varphi(\delta) = \frac{\delta(\alpha_H)}{\alpha_H} = \left\langle \frac{d\alpha_H}{\alpha_H}, \delta \right\rangle. \tag{27}$$

Applying $-\vee = \text{Hom}_S(-, S)$ to (26) yields a split exact sequence

$$0 \longleftarrow D_H(\mathcal{A})^\vee \longleftarrow \Omega^1(\mathcal{A}) \xleftarrow{\varphi^\vee} S \longleftarrow 0. \tag{28}$$

By (27) and (28), we can identify $\varphi^\vee = d\alpha_H/\alpha_H$, and hence

$$D_H(\mathcal{A})^\vee \cong \Omega^1(\mathcal{A})/S \frac{d\alpha_H}{\alpha_H} \cong \frac{d\alpha_H}{\alpha_H} \wedge \Omega^1(\mathcal{A}).$$

THEOREM 10. *The statement of Theorem 2 holds true if \mathcal{A} is weakly (dually) tame.*

Proof. By [Yos05, Corollary 3.3] and Theorem 8, we may assume that $\ell \geq 5$. We consider the weakly tame case only; the weakly dually tame case can be treated along the same lines using the direct sum decomposition (23) instead of Lemma 9, and proving surjectivity of res_H instead of res_H^1 .

By Lemma 9 and weak tameness, $\text{pdim}_S(d\alpha_H/\alpha_H \wedge \Omega^1(\mathcal{A})) \leq 1$. Then also $\text{pdim}_{S'} M^1 \leq 1$ due to the exact sequence

$$0 \longrightarrow \frac{d\alpha_H}{\alpha_H} \wedge \Omega^1(\mathcal{A}) \xrightarrow{\alpha_H} \frac{d\alpha_H}{\alpha_H} \wedge \Omega^1(\mathcal{A}) \longrightarrow M^1 \longrightarrow 0.$$

Indeed, if $F_\bullet : F_1 \hookrightarrow F_0$ is an S -free resolution of $d\alpha_H/\alpha_H \wedge \Omega^1(\mathcal{A})$, then $F_\bullet/\alpha_H F_\bullet$ is an S' -free resolution of M^1 . This follows immediately from comparing the two spectral sequences associated to the double complex defined by multiplication by α_H on F_\bullet .

By the Auslander–Buchsbaum theorem, this shows that

$$\forall x \in H : \dim S'_x \geq 3 \Rightarrow \text{depth } M_x^1 \geq 2, \quad (29)$$

where x is considered as a scheme-theoretic point. From (14) we get an exact sequence of S' -modules

$$0 \rightarrow M^1 \rightarrow \Omega^1(\mathcal{A}^H) \rightarrow C^1 \rightarrow 0. \quad (30)$$

Assuming that \mathcal{A}^H is free, (29) applied to the localization of (30) at x shows that $\text{depth } C_x^1 \geq 1$ for any $x \in H$ of codimension at least 3. However, by Proposition 4 and our hypothesis $\ell \geq 5$, C^1 has codimension at least 3 in H . Thus, $\text{Ass } C^1 = \emptyset$, $C^1 = 0$, res_H^1 is surjective, and \mathcal{A} is free by [Yos05, Theorem 2.5]. \square

REFERENCES

- ATW07 T. Abe, H. Terao and M. Wakefield, *The characteristic polynomial of a multiarrangement*, Adv. Math. **215** (2007), 825–838; [MR 2355609](#).
- MS01 M. Mustața and H. K. Schenck, *The module of logarithmic p -forms of a locally free arrangement*, J. Algebra **241** (2001), 699–719; [MR 1843320\(2002c:32047\)](#).
- OS80 P. Orlik and L. Solomon, *Combinatorics and topology of complements of hyperplanes*, Invent. Math. **56** (1980), 167–189; [MR 558866\(81e:32015\)](#).
- OT92 P. Orlik and H. Terao, *Arrangements of hyperplanes*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 300 (Springer, Berlin, 1992); [MR 1217488\(94e:52014\)](#).
- ST87 L. Solomon and H. Terao, *A formula for the characteristic polynomial of an arrangement*, Adv. Math. **64** (1987), 159–179.
- Ter81 H. Terao, *Generalized exponents of a free arrangement of hyperplanes and Shepherd-Todd-Brieskorn formula*, Invent. Math. **63** (1981), 159–179; [MR 608532\(82e:32018b\)](#).
- Yos04 M. Yoshinaga, *Characterization of a free arrangement and conjecture of Edelman and Reiner*, Invent. Math. **157** (2004), 449–454; [MR 2077250\(2005d:52044\)](#).
- Yos05 M. Yoshinaga, *On the freeness of 3-arrangements*, Bull. Lond. Math. Soc. **37** (2005), 126–134; [MR 2105827\(2005i:52030\)](#).
- Zie89a G. M. Ziegler, *Combinatorial construction of logarithmic differential forms*, Adv. Math. **76** (1989), 116–154; [MR 1004488\(90j:32016\)](#).
- Zie89b G. M. Ziegler, *Multiarrangements of hyperplanes and their freeness*, in *Singularities (Iowa City, IA, 1986)*, Contemporary Mathematics, vol. 90 (American Mathematical Society, Providence, RI, 1989), 345–359; [MR 1000610\(90e:32015\)](#).

Mathias Schulze mschulze@math.okstate.edu

Department of Mathematics, Oklahoma State University, Stillwater,
OK 74078, USA