60 Poster Presentations

**Methods.** HTW consulted with external organizations to identify the first steps toward effective PPI. Public partners were recruited as a priority before working together on a PPI strategy. Building smart capabilities is key to establishing effective PPI and future-proofing. HTW established a PPI Standing Group to inform HTW throughout its work, including the development of processes and procedures.

Results. Knowledge and resources have been shared and future collaborations identified, including events to encourage new topics from patients and the public. The HTW PPI lead has become a member of key PPI groups, locally and internationally. HTW has recruited public partners who are actively contributing as full members of the Assessment Group and the Appraisal Panel; two members on each Committee. The PPI Standing Group has been established. They have provided advice and co-produced PPI tools for piloting.

Conclusions. The PPI Standing Group concluded that PPI methods and approaches should be tailored for each project based on best practice, and should be piloted to allow them to evolve based on impact evaluation. A PPI strategy or framework would be more useful at a later stage. HTW is committed to identifying and following best practice. Future-proofing and building smart capability will be key to ensuring that HTW develops effective PPI that can be dynamic and responsive to the evolving PPI and HTA landscapes.

## PP125 PhotoVoice: Promoting Knowledge Exchange About Patients' Experiences

Yolanda Triñanes Pego (avalia-t3@sergas.es), Lucía Varela, José Mazaira-Castro and Maria José Faraldo Vallés

**Introduction.** In the past decades the community-based participatory research method known as PhotoVoice has gained relevance, but there are few published studies on its application in the field of health technology assessment (HTA) and clinical practice guidelines (CPGs). The aim of this presentation was to describe a PhotoVoice project linked to a CPG on major depression in children and adolescents.

**Methods.** The design of the study was adapted to the main objective, which was to enhance the understanding of major depression and improve clinical practice with the contributions of clinicians, methodologists, and patients. Seven adolescents and ten of their family members participated in the study through PhotoVoice sessions and focus groups. The audio recordings of all sessions were transcribed verbatim and coded, and a thematic analysis was undertaken.

Results. Six themes emerged: (i) a lack of understanding and information about depression in childhood and adolescence; (ii) the importance of support groups; (iii) the need to favor early care and access to services; (iv) the adaptation of therapeutic strategies tailored to individual needs; (v) the sensitivity of professionals; and (vi) fostering interaction between the health and education systems. Photographic exhibitions were planned to share the main results. These exhibitions were promoted to increase public awareness and reduce stigmatization,

and to reach clinicians and policy makers. From a methodological point of view, the use of PhotoVoice in this study helped to effectively incorporate the lived experiences, concerns, and preferences of patients and their relatives into the CPG. The study also confirmed the value of photographs and participatory methods. The main limitations and strengths of the study, as well as suggestions for future research, are also outlined.

**Conclusions.** PhotoVoice is a flexible, effective, and innovative method of obtaining information about patients' perspectives and experiences, and it offers the added value of being able to reach the main stakeholders, including policymakers and the public.

## PP126 Analyses Of User Requirements In The Evaluation Of Medical Equipment

Ricardo Bertoglio Cardoso (ricardobcardoso@gmail. com), Priscila G. Brust-Renck, Flavio Sanson Fogliatto and Helena Barreto Dos Santos

Introduction. Human-centered approaches to eliciting requirements for medical equipment selection are recognized as improving healthcare outcomes, safety, and end-user satisfaction. Nevertheless, there are many challenges to conducting rigorous investigations to identify requirements that satisfy different hospital services and types of end users (e.g., patients, healthcare professionals, and clinical engineers). By establishing a systematic method for selecting medical recliners, this study provides detailed technical characteristics and user requirements associated with several hospital areas, as well as a comparison between two end users (health professionals and patients) and their different perceptions of usability.

Methods. First, clinical engineers and senior nurses from seven hospital services identified and rated the technical characteristics of medical recliners. Ratings were then used to stratify all services in well-defined similar groups using hierarchical and non-hierarchical clustering algorithms. Next, users of hospital recliners (60 patients and 56 healthcare providers) from each group were interviewed to identify their requirements for an ideal medical recliner. Finally, analyses of variance were performed to identify consensus decisions from users across the different hospital contexts as to which technical characteristics were the most relevant.

**Results.** The contribution of senior nurses and clinical engineers led to the identification of 41 technical characteristics. The analysis of 116 participant interviews identified 95 different requirements, extracted from 1,052 user suggestions. Correspondence analysis of the most important requirements, combined for each of the three stratified service groups, indicated that two-thirds of all user requirements (14 out of 20) were fulfilled by five out of 32 quantitative technical characteristics, regardless of context.

**Conclusions.** Human-centered methods can identify similarities between health technology characteristics and decrease the complexity associated with selecting technologies, while simultaneously fulfilling the requirements of multiple users and hospital departments.