THE BRITISH JOURNAL OF NUTRITION DIRECTIONS TO CONTRIBUTORS

(Revised September 1980)

The British Journal of Nutrition publishes reports in English of original work in all branches of nutrition from any country. It does not print reviews of the literature or polemical articles, but the Editorial Board is willing to consider original articles critically re-examining published information and the conclusions drawn from it. The aim of all work presented should be to develop nutritional concepts.

Papers submitted for publication should be as concise as possible. Economy of space should not, however, be achieved by suppressing useful results.

Papers should be accompanied by a signed statement to the effect that the author accepts the conditions laid down in Directions to Contributors. Special attention is directed to the sections below about the preparation of the typescript; care in this matter will hasten publication. The Editors will return any typescript that does not conform to these conditions. Contributors of accepted articles will be asked to assign their copyright, on certain conditions, to The Nutrition Society to help protect their material.

Communications. Papers submitted for publication should be sent to Dr G. A. J. Pitt (*British Journal of Nutrition*), Department of Biochemistry, University of Liverpool, PO Box 147, Liverpool L69 3BX.

General. Submission of a paper to the Editorial Board will be held to imply that it represents the results of original research or of an original interpretation of existing knowledge not previously published; that it is not under consideration for publication elsewhere; and that if accepted for the *British Journal of Nutrition* it will not be published elsewhere in the same form, in English or any other language, without the consent of the Editorial Board.

Authors' names should be given without titles or degrees and one forename may be given in full. The name and address of the laboratory or institution where the work was performed should be given. Any necessary descriptive material about the author, e.g. Beit Memorial Fellow, should appear in parentheses after the author's name or at the end of the paper and not in the form of a footnote.

Typescripts should bear the name and address of the person to whom the proof of the paper is to be sent and should also give a shortened version of the paper's title, not exceeding forty-five letters and spaces in length, suitable for a running title in the published pages of the work.

If a typescript that has been returned to the author for revision is not resubmitted within three months, it may, on resubmission, be deemed a new paper and the date of receipt altered accordingly.

Form of Papers Submitted for Publication. The onus of preparing a paper in a form suitable for sending to press lies in the first place with the author. Authors should consult a current issue in order to make themselves familiar with the practice of the British Journal of Nutrition as to typographical and other conventions, use of cross-headings, layout of tables and so on. Attention to these and other details (mentioned below) in the preparation of the typescript before it is sent to the Editors will shorten the time required for publication: the need for undue amounts of editorial revision caused by badly prepared typescript will lead to delay in publication. Papers on specialized aspects of the subject should be so presented as to make them intelligible, without undue difficulty, to the ordinary reader of the Journal. Sufficient information should be given to permit repetition of the published work by any competent reader of the Journal.

Papers should be in double-spaced typing on one side of sheets of paper (thin paper should not be used) of uniform size with wide margins. At the ends of lines of typing, words

12

should not be hyphenated unless hyphens are to be printed. A space of 50 mm should be left at the top of the first sheet. The use of line-numbered paper is encouraged. The top copy and one photocopy should be submitted, packed flat. The paper should be written in English, the spelling being that of the Concise Oxford Dictionary, 6th ed. Oxford: Clarendon Press, 1976, and should, in general, be divided into the following parts. (a) Synopsis: each paper must open with a synopsis not more than 5% of the length of the following text. This synopsis should aim at giving a picture in miniature of the entire article. The past tense should be used in referring to the author's experimental work. The present tense may be used where reference to existing knowledge is necessary, or where the author is stating what is shown or concluded. The change of tense should clearly differentiate the author's contribution from what is already known. The sequence in the synopsis should be the same as that in the paper. It is desirable to divide the synopsis into a series of numbered paragraphs giving, where relevant, the following information: a succinct account of the experimental work with essential facts about apparatus, chemicals, methods and animals; the results, singling out new information; the conclusions from the results. (b) Introductory paragraph: it is not now customary to introduce a paper with a full account of the relevant literature, but the introductory paragraph should help the reader by indicating briefly the nature of the question asked and the reasons for asking it. The answer obtained should be indicated if it is possible to do so shortly. (c) Experimental methods adopted: with chemical papers, the Experimental part will normally appear towards the end, but with other types of publication Methods should appear after the introduction. (d) Results: these should be given as concisely as possible, with the help of figures or tables. (e)Discussion: it is desirable that the presentation of the results and the discussion of their significance should be considered separately. (f) References: these should be given in the text thus: Osborne & Mendel (1916), (Culbertson & Thomas, 1934); where a paper to be cited has more than two authors, citations should appear thus (Osborne et al. 1919). Where more than one paper by the same authors has appeared in one year the reference should be given as follows: Osborne & Mendel (1914a); Osborne & Mendel (1914b); or Osborne & Mendel (1914a, b); (Osborne & Mendel, 1914b, 1916: Culbertson & Thomas, 1934). In the text, references should be given in chronological order. Where the lists of authors of two papers appearing in the same year begin with the same name but differ subsequently, references should include sufficient names to differentiate between the two papers, e.g. Thomas, Smith et al. (1960), Thomas, Wilson et al. (1960). At the end of the paper, on a page(s) separate from the text, references should be listed in alphabetical order according to the name of the first author of the publication quoted, names with prefixes being entered under the prefix, and should include the author's initials; the title of the paper (except that of a thesis) should not be included. Names and initials of authors of unpublished work should be given in the text and not included in the References. Titles of journals should be abbreviated as in the *World List of Scientific Periodicals*, 4th ed. London: Butterworths, 1963–5. References to books and monographs should include the town of publication and the number of the edition to which reference is made. Thus:

- Ahrens, E. H. Jr, Hirsch, J., Insull, W. Jr & Peterson, M. L. (1958). In Chemistry of Lipids as Related to Atherosclerosis, p. 222 [I. H. Page, editor]. Springfield, Ill.: C. C. Thomas.
- Culbertson, C. C. & Thomas, B. H. (1934). Rep. Iowa agric. Exp. Stn 1933-4, p. 51.
- Fairey, N. H. (1938). Nature, Lond. 142, 1156.
- King, H. (1941). J. chem. Soc. p. 338.
- Osborne, T. B. & Mendel, L. B. (1914a). J. biol. Chem. 17, 325.
- Osborne, T. B. & Mendel, L. B. (1914b). J. biol. Chem. 18, 1.
- Osborne, T. B. & Mendel, L. B. (1916). Biochem. J. 10, 534.
- Osborne, T. B., Mendel, L. B. & Ferry, E. L. (1919). J. biol. Chem. 37, 233.
- Starling, E. H. (1915). Principles of Human Physiology, 2nd ed. London: J. & A. Churchill.
- Summerson, W. H. (1938). J. biol. Chem. 123, cxix.
- Wilson, J. (1965). Leber's disease. PhD Thesis, University of London.

Units. Results should be presented in metric units according to the International System of Units (see *Quantities, Units, and Symbols,* London: The Royal Society, 1971, and Metric Units, Conversion Factors and Nomenclature in Nutritional and Food Sciences. London: The Royal Society, 1972~reproduced in Proc. Nutr. Soc. (1972), **31**, 239).

Energy measurements should be expressed in joules.

For substances of known molecular weight, e.g. glucose, urea, Ca, Na, Fe, K, P, values should be expressed as mol/1; for substances of indeterminate molecular weights, e.g. phospholipids, proteins, and for trace elements, e.g. Cu, Zn, g/l should be used.

Time. The 24h clock should be used, e.g. 15.00 hours.

Statistical Treatment of Results. In general, it is not necessary to publish all the individual results of replicated tests. A statement of the number, their mean value and some appropriate measure of their variability is usually sufficient.

The methods of analysis followed should be indicated, but statistical details, such as tables of analysis of variance, should be given only if they are relevant to the discussion. A statement that the difference between the means for two groups of values is statistically significant should include an indication of the level of significance attained.

Where means are quoted, normally the measure of variation most appropriate is the standard error of the mean; occasionally, as for instance where there is specific interest in the distribution of the individual values in the sample on which the mean is based, the standard deviation may be more appropriate. The measure adopted must be clearly stated in the text and in each table. Where the measure of variation is presented as a separate column, \pm should not be repeated before each value.

Figures. These include graphs, histograms, complex formulas, metabolic pathways. Originals and one photocopy should be submitted, each on a separate sheet, not larger over all than the sheets on which the paper itself is typed, and packed flat. Mounting on heavy cardboard is undesirable. Photographs of line drawings are acceptable if printed on matt paper. In curves presenting experimental results, the determined points should be clearly shown, the symbols used being, in order of preference, $\bigcirc, \spadesuit, \triangle, \clubsuit, \square, \blacksquare, \times +$. Curves and symbols should be drawn with a mechanical aid and not free-hand, and should not extend beyond the experimental points. Scale-marks on the axes should be on the inner side of each axis and should extend beyond the last experimental point.

Numbers and letters should be written not on the figure but in the correct position on a flyleaf of tracing paper firmly attached. On the flyleaf should be typed or stencilled: (a) the title of the paper and names of the authors; (b) the figure number. Legends for all figures should be typed on one separate sheet (two or more, if necessary) and numbered corresponding to the relevant figures. Each figure, with its legend, should be comprehensible without reference to the text. The approximate position of each should be indicated in the margin of the text thus: 'Fig. 1 near here'.

Plates. Glossy photographs are required, and should be accompanied by a legend prepared as above. The size of photomicrographs may have to be altered in printing. To avoid mistakes, the magnification will be shown by a scale on the photograph itself, e.g. thus: $[1 \mu m]$. The scale with the appropriate unit should be drawn by the author on the fiyleaf together with any lettering and will be inserted by the Press. Do not write details on the back of prints, bend, use paper-clips or mark in any way. The plate number, title of the paper and authors' names should be typed on a label and pasted on to the back of the print.

Tables. Tables should carry headings describing their content and should be comprehensible without reference to the text. The dimensions of the values, e.g. mg/kg, should be given at the top of each column and not repeated on each line of the table. Tables should not normally be included in the body of the text, but should be typed on separate sheets. Tables should not be subdivided by ruled lines. Their approximate position should be indicated in the margin of the text thus: 'Table 1 near here'.

Diagrams. Diagrams to appear as tables (e.g. flow diagrams) should be prepared as for Tables using Letraset or stencils. No flyleaf is required.

Key Words. Authors should supply two to three key words or phrases (each containing up to three words) on the title page of typescripts. These will be used to compile subject indexes of published papers.

Chemical Formulas. These should be written as far as possible on a single horizontal line. With inorganic substances, formulas may be used, particularly in the Experimental part, at the discretion of the Editors. With salts, it must be stated whether or not the anhydrous material is used, e.g. anhydrous $CuSO_4$, or which of the different crystalline forms is meant, e.g. $CuSO_4$. $5H_2O$, $CuSO_4$. H_2O .

Descriptions of Solutions, Compositions and Concentrations. Solutions of common acids, bases and salts should be defined in terms of molarity (M), e.g. 0.1 M-NaH₂PO₄. Compositions expressed as mass per unit mass (w/w) should have values expressed as ng, µg, mg or g per kg; similarly for concentrations expressed as mass per unit volume (w/v), the denominator being the litre. Concentrations or compositions should not be expressed on a percentage basis. The common measurements used in nutritional studies, e.g. digestibility, biological value and net protein utilization, should be expressed as decimals rather than as percentages, so that amounts of available nutrients can be obtained from analytical results by direct multiplication. See Metric Units, Conversion Factors and Nomenclature in Nutritional and Food Sciences. London: The Royal Society, 1972 (para 8).

Nomenclature of Vitamins. Most of the names for vitamins and related compounds that are accepted by the Editors are those recommended by the IUNS Committee on Nomenclature. See Nutr. Abstr. Rev. A (1978), 48, 831.

Vitamin A_1 RetinolRetinene or retinalRetinaldehydeVitamin A acidRetinoic acidVitamin A_2 or 3-dehydroretinolDehydroretinolRetinene2 or 3-dehydroretinalDehydroretinolRetinene2 or 3-dehydroretinalDehydroretinolRetinene2 or 3-dehydroretinalDehydroretinolRetinene2 or 3-dehydroretinalErgocalciferolVitamin D2 or calciferolErgocalciferolVitamin D3CholecalciferolVitamin K1 orPhylloquinoneVitamin K2 series*MenaquinonesVitamin B1, aneurin(e)Thiaminor thiamineThiaminVitamin B2 or riboflavineNicotinic acidNicotinic acid or niacinNicotinic acidNicotinic acid or folacin(e)Pteroylmonoglutamic acidVitamin B120 orPyridoxalPyridoxalPyridoxalPyridoxaminePyridoxalVitamin B120 orNitritocobalaminNitritocobalaminHydroxocobalaminVitamin B120 orNitritocobalaminNitritocobalaminPyridoxalPyridoxaminePuridoxalPyridoxalPuridoxalPyridoxaninePuridoxalVitamin B120 orNitritocobalaminHydroxocobalmideNitritocobalaminVitamin C or L-ascorbicAscorbic acidacidL-dehydroascorbic acidDehydroascorbic acidDehydroascorbic acid	Previous name	Recommended name
Vitamin A acidRetinoic acidVitamin A2 or 3-dehydroretinolDehydroretinolRetinene2 or 3-dehydroretinalDehydroretinolRetinene2 or 3-dehydroretinalDehydroretinolVitamin D2 or calciferolErgocalciferolVitamin D3CholecalciferolVitamin K1 orPhylloquinoneVitamin K2 series*MenaquinonesVitamin K3, menadioneMenadioneor menaphthoneThiaminVitamin B1, aneurin(e)Thiaminor thiamineNicotinic acid or niacinNiacinamide or nicotinicNicotinic acidNiacinamide or nicotinicNicotinic acidVitamin B1, adermin orPyridoxalPyridoxalPyridoxalPyridoxalPyridoxalPyridoxalPyridoxalPyridoxalidionNitritocobalaminVitamin B12aNitritocobalaminVitamin B12aNitritocobalaminVitamin C or L-ascorbicAscorbic acidAscorbic acidAscorbic acid	Vitamin A ₁	Retinol
Vitamin A_2 or 3-dehydro- retinolDehydroretinolRetinene2 or 3-dehydro- retinalDehydroretinolRetinene2 or 3-dehydro- retinalDehydroretinolVitamin D_2 or calciferolDehydroretinaldehydeVitamin D_2 or calciferolCholecalciferolVitamin D_3 CholecalciferolVitamins ESee Generic descriptorsVitamin K_1 or phylloquinonePhylloquinoneVitamin K_2 series*MenaquinonesVitamin B_1, aneurin(e) or thiamineThiaminVitamin B_2 or riboflavineRiboflavinNicotinic acid or niacin nicotinic acid or niacin acid amideNicotinic acid Nicotinic acid Nicotinic acid Nicotinic acid NicotinieFolic acid or folacin(e)Pteroylmonoglutamic acid PyridoxalPyridoxal Pyridoxal Nitamin B_{12} or cobalamin Vitamin B_{12a}, B_{12b} or hydroxocobamideNitritocobalamin Y-Inositol Pantothenic acid BiotinNitritocobalamin Choline p-Aminobenzoic acid acidNitritocobalamin Ascorbic acid Ascorbic acid	Retinene or retinal	Retinaldehyde
retinolDehydroretinaldehydeRetinene, or 3-dehydro- retinalDehydroretinaldehydeRetinene, or 3-dehydro- retinalDehydroretinaldehydeVitamin D_2 or calciferolErgocalciferolVitamin D_3 CholecalciferolVitamins ESee Generic descriptorsVitamin K_1 or phylloquinonePhylloquinoneVitamin K_2 series*MenaquinonesVitamin K_3, menadione or menaphthoneMenadioneVitamin B_1, aneurin(e) or thiamineThiaminVitamin B_2 or riboflavine Nicotinic acid or niacin niacinamide or nicotinic acid amideRiboflavin Nicotinic acid Nicotinic acid NicotinamideFolic acid or folacin(e) Vyridoxal Pyridoxal Nitamin B_{12} or cobalamin Vitamin B_{12a}, B_{12b} or hydroxocobamidePyridoxal Pyridoxal Pyridoxal Pyridoxal PyridoxanineVitamin B_12a, B_{12b} or hydroxocobamideNitritocobalamin Puntothenic acid BiotinNitritocobalamin Vitamin C or t-ascorbic acidAscorbic acid Ascorbic acid	Vitamin A acid	Retinoic acid
retinolDehydroretinaldehydeRetinene, or 3-dehydro- retinalDehydroretinaldehydeRetinene, or 3-dehydro- retinalDehydroretinaldehydeVitamin D_2 or calciferolErgocalciferolVitamin D_3 CholecalciferolVitamins ESee Generic descriptorsVitamin K_1 or phylloquinonePhylloquinoneVitamin K_2 series*MenaquinonesVitamin K_3, menadione or menaphthoneMenadioneVitamin B_1, aneurin(e) or thiamineThiaminVitamin B_2 or riboflavine Nicotinic acid or niacin niacinamide or nicotinic acid amideRiboflavin Nicotinic acid Nicotinic acid NicotinamideFolic acid or folacin(e) Vyridoxal Pyridoxal Nitamin B_{12} or cobalamin Vitamin B_{12a}, B_{12b} or hydroxocobamidePyridoxal Pyridoxal Pyridoxal Pyridoxal PyridoxanineVitamin B_12a, B_{12b} or hydroxocobamideNitritocobalamin Puntothenic acid BiotinNitritocobalamin Vitamin C or t-ascorbic acidAscorbic acid Ascorbic acid	Vitamin A, or 3-dehydro-	Dehydroretinol
retinal Vitamin D_2 or calciferol Vitamin D_3 Vitamin D_3 Vitamin D_3 Vitamin D_3 Vitamin D_3 Vitamin D_3 Vitamin D_3 Vitamin E Vitamin E Vitamin K_1 or phylloquinone Vitamin K_2 series Vitamin K_3 , menadione or menaphthone Vitamin B_1 , aneurin(e) or thiamine Vitamin B_2 or riboflavine Nicotinic acid or niacin Nicotinic acid or folacin(e) Vitamin B_1 or cobalamin Vitamin B_{12} or cobalamin Vitamin B_{12} Inositol or meso-inositol Pantothenic acid Biotin Choline p-Aminobenzoic acid Ascorbic acid	retinol	
Vitamin D_3 CholecalciferolVitamins ESee Generic descriptorsVitamins K1 orPhylloquinonephylloquinoneVitamin K2 seriesVitamin K3, menadioneMenaquinonesor menaphthoneThiaminVitamin B1, aneurin(e)Thiaminor thiamineRiboflavinNicotinic acid or niacinNicotinic acidNiacinamide or nicotinicNicotinic acidacid amidePteroylmonoglutamic acidFolic acid or folacin(e)Pteroylmonoglutamic acidVitamin B6, adermin orpyridoxalPyridoxalPyridoxalPyridoxalPyridoxalVitamin B12aOr cobalaminVitamin B12aNitritocobalaminVitamin B12aNitritocobalaminVitamin B12aPantothenic acidBiotinBiotinCholinePantothenic acidP-Aminobenzoic acidP-Aminobenzoic acidAscorbic acidAscorbic acid		Dehydroretinaldehyde
Vitamin D_3 CholecalciferolVitamins ESee Generic descriptorsVitamins K1 orPhylloquinonephylloquinoneVitamin K2 seriesVitamin K3, menadioneMenaquinonesor menaphthoneThiaminVitamin B1, aneurin(e)Thiaminor thiamineRiboflavinNicotinic acid or niacinNicotinic acidNiacinamide or nicotinicNicotinic acidacid amidePteroylmonoglutamic acidFolic acid or folacin(e)Pteroylmonoglutamic acidVitamin B6, adermin orpyridoxalPyridoxalPyridoxalPyridoxalPyridoxalVitamin B12aOr cobalaminVitamin B12aNitritocobalaminVitamin B12aNitritocobalaminVitamin B12aPantothenic acidBiotinBiotinCholinePantothenic acidP-Aminobenzoic acidP-Aminobenzoic acidAscorbic acidAscorbic acid	Vitamin D ₂ or calciferol	Ergocalciferol
Vitamins \vec{E} See Generic descriptorsVitamin K_1 or phylloquinonePhylloquinoneVitamin K_2 series*MenaquinonesVitamin K_3 , menadione or menaphthoneMenadioneVitamin B_1 , aneurin(e) or thiamineThiaminVitamin B_2 or riboflavineRiboflavinNicotinic acid or niacin Nicotinic acid or niacinNicotinic acidNiacinamide or nicotinic acid amidePteroylmonoglutamic acidFolic acid or folacin(e)Pteroylmonoglutamic acidPyridoxal PyridoxalPyridoxalPyridoxaninePyridoxalVitamin B_{12a} , B_{12b} or hydroxocobamideNitritocobalaminVitamin B_{12c} NitritocobalaminVitamin B_{12c} , or hydroxocobamideNitritocobalaminVitamin B_{12c} , or hydroxocobamidePyridoxalPyridoxali PyridoxalicPyridoxalPyridoxic Pantothenic acid BiotinPantothenic acid BiotinPantothenic acid Vitamin C or t-ascorbic acidAscorbic acid Ascorbic acid		
phylloquinoneVitamin K_2 series*MenaquinonesVitamin K_3 , menadioneMenadioneor menaphthoneMenadioneVitamin B_1 , aneurin(e)Thiaminor thiamineRiboflavinNicotinic acid or niacinNicotinic acidNicotinic acid or niacinNicotinic acidNiacinamide or nicotinicNicotinic acidacid amidePteroylmonoglutamic acidFolic acid or folacin(e)Pteroylmonoglutamic acidPyridoxalPyridoxalPyridoxalPyridoxalPyridoxanineCyanocobalaminVitamin B_{12a} , B_{12b} orHydroxocobalaminNitritocobalaminWitritocobalaminVitamin B_{12a} , B_{12b} orNitritocobalaminPantothenic acidBiotinBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or t-ascorbicAscorbic acid		See Generic descriptors
phylloquinoneVitamin K_2 series*MenaquinonesVitamin K_3 , menadioneMenadioneor menaphthoneMenadioneVitamin B_1 , aneurin(e)Thiaminor thiamineRiboflavinNicotinic acid or niacinNicotinic acidNicotinic acid or niacinNicotinic acidNiacinamide or nicotinicNicotinic acidacid amidePteroylmonoglutamic acidFolic acid or folacin(e)Pteroylmonoglutamic acidPyridoxalPyridoxalPyridoxalPyridoxalPyridoxanineCyanocobalaminVitamin B_{12a} , B_{12b} orHydroxocobalaminNitritocobalaminWitritocobalaminVitamin B_{12a} , B_{12b} orNitritocobalaminPantothenic acidBiotinBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or t-ascorbicAscorbic acid	Vitamin K, or	
Vitamin K2 series*MenaquinonesVitamin K3, menadione or menaphthoneMenadioneVitamin B1, aneurin(e) or thiamineThiaminVitamin B2 or riboflavine Nicotinic acid or niacin Nicotinic acid or niacin Nicotinic acid or niacin acid amideRiboflavinNicotinic acid or niacin Nicotinic acid or niacin acid amideRiboflavin Nicotinic acid Nicotinic acid Nicotinic acid Nicotinic acid Nicotinic acid Nicotinic acid Pyridoxal Pyridoxal Pyridoxal Pyridoxal Pyridoxal Nitritocobalamin Vitamin B122 or cobalamin Vitamin B122 or meso-inositol Pantothenic acid BiotinPyridoxal 	phylloquinone	
Vitamin K_3 , menadione or menaphthoneMenadioneVitamin B_1 , aneurin(e) or thiamineThiaminVitamin B_2 or riboflavineRiboflavinNicotinic acid or niacin Nicotinic acid or niacinNicotinic acidNicotinic acid or niacin nicotinic acid or niacinNicotinic acidNicotinic acid or niacin acid amideNicotinic acidFolic acid or folacin(e) Vitamin B_6 , adermin or pyridoxalPteroylmonoglutamic acidPyridoxal Vitamin B_{12} or cobalamin Vitamin B_{12} or cobalamin Vitamin B_{120} or hydroxocobamidePyridoxal Pyridoxal Pyridoxacobalamin HydroxocobalideVitamin B_{12c} Nitritocobalamin Pantothenic acid BiotinNitritocobalamin Pantothenic acid Biotin Choline p -Aminobenzoic acid Ascorbic acid	Vitamin K, series	*Menaquinones
or menaphthone Vitamin B_1 , aneurin(e) or thiamine Vitamin B_2 or riboflavine Nicotinic acid or niacin Nicotinic acid or niacin Nicotinic acid or niacin Nicotinic acid or niacin Nicotinic acid Nicotinic acid Pteroylmonoglutamic acid Pyridoxal Pyrido		
Vitamin B_1 , aneurin(e) or thiamineThiaminVitamin B_2 or riboflavineRiboflavinNicotinic acid or niacinNicotinic acidNicotinic acid or niacinNicotinic acidNiacinamide or nicotinic acid amideNicotinic acidFolic acid or folacin(e)Pteroylmonoglutamic acidVitamin B_6 , adermin or pyridoxalPyridoxalPyridoxalPyridoxalPyridoxaminePyridoxalVitamin B_{12a} , B_{12b} or hydroxocobamideNitritocobalaminVitamin B_{12c} NitritocobalaminVitamin B_{12c} NitritocobalaminPantothenic acidPantothenic acidBiotinBiotinCholine p-Aminobenzoic acidCholine p-Aminobenzoic acid		
or thianineRiboflavinVitamin B_2 or riboflavineRiboflavinNicotinic acid or niacinNicotinic acidNiacinamide or nicotinicNicotinic acidNiacinamide or nicotinicNicotinic acidAitaria amideNicotinamideFolic acid or folacin(e)Pteroylmonoglutamic acidVitamin B_6 , adermin orPyridoxalPyridoxalPyridoxalPyridoxalPyridoxalVitamin B_{12a} , B_{12b} orHydroxocobalaminVitamin B_{12a} , B_{12b} orHydroxocobalaminNitritocobalaminVitamin B_{12c} NitritocobalaminY-InositolPantothenic acidBiotinBiotinBiotinCholineCholinep-Aminobenzoic acidAscorbic acid		Thiamin
Nicotinic acid or niacinNicotinic acidNiacinamide or nicotinicNicotinic acidNicotinamideNicotinamideFolic acid or folacin(e)Pteroylmonoglutamic acidVitamin B ₆ , adermin orPyridoxinepyridoxolPyridoxalPyridoxalPyridoxalPyridoxaminePyridoxalminVitamin B ₁₂ or cobalaminCyanocobalaminVitamin B _{12e} , orHydroxocobalaminVitamin B _{12e} NitritocobalaminVitamin B _{12e} NitritocobalaminInositol or meso-inositolPantothenic acidBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid		
Nicotinic acid or niacinNicotinic acidNiacinamide or nicotinicNicotinic acidNicotinamideNicotinamideFolic acid or folacin(e)Pteroylmonoglutamic acidVitamin B ₆ , adermin orPyridoxinepyridoxolPyridoxalPyridoxalPyridoxalPyridoxaminePyridoxalminVitamin B ₁₂ or cobalaminCyanocobalaminVitamin B _{12e} , orHydroxocobalaminVitamin B _{12e} NitritocobalaminVitamin B _{12e} NitritocobalaminInositol or meso-inositolPantothenic acidBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	Vitamin B, or riboflavine	Riboflavin
acid amideFolic acid or folacin(e)Pteroylmonoglutamic acidFolic acid or folacin(e)Pteroylmonoglutamic acidVitamin B_{e} , adermin or pyridoxalPyridoxinePyridoxalPyridoxalPyridoxaminePyridoxalVitamin B_{12} or cobalaminCyanocobalaminVitamin B_{12a} , B_{12b} or hydroxocobamideHydroxocobalaminVitamin B_{12c} NitritocobalaminVitamin B_{12c} NitritocobalaminPantothenic acidPantothenic acidBiotinBiotinCholineCholinep-Aminobenzoic acidAscorbic acid	Nicotinic acid or niacin	Nicotinic acid
Folic acid or folacin(e) Vitamin B ₆ , adermin or pyridoxolPteroylmonoglutamic acid PyridoxinePyridoxal Pyridoxal Vitamin B ₁₂ or cobalamin Vitamin B _{12a} , B _{12b} or hydroxocobalmidePyridoxal Pyridoxal Pyridoxamine Cyanocobalamin Hydroxocobalamin Hydroxocobalamin PyridoxanineVitamin B _{12a} Nitritocobalamin Postiol Pantothenic acid Biotin P-Aminobenzoic acid AcidNitritocobalamin Pyridoxal Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalamin Pyridoxacobalami	Niacinamide or nicotinic	Nicotinamide
Vitamin B_{e} , adermin or pyridoxolPyridoxinePyridoxalPyridoxalPyridoxaminePyridoxamineVitamin B_{12} or cobalaminCyanocobalaminVitamin B_{12a} , B_{12b} or hydroxocobalmideHydroxocobalaminVitamin B_{12e} NitritocobalaminInositol or meso-inositol Ψ -InositolPantothenic acidBiotinBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	acid amide	
pyridoxolPyridoxalPyridoxalPyridoxaminePyridoxamineVitamin B_{12} or cobalaminCyanocobalaminVitamin B_{12a} , B_{12b} orHydroxocobalaminhydroxocobamideVitamin B_{12c} Vitamin B_{12c} NitritocobalaminInositol or meso-inositol Ψ -InositolPantothenic acidPantothenic acidBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	Folic acid or folacin(e)	Pteroylmonoglutamic acid
pyridoxolPyridoxalPyridoxalPyridoxaminePyridoxamineVitamin B_{12} or cobalaminCyanocobalaminVitamin B_{12a} , B_{12b} orHydroxocobalaminhydroxocobamideVitamin B_{12c} Vitamin B_{12c} NitritocobalaminInositol or meso-inositol Ψ -InositolPantothenic acidPantothenic acidBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	Vitamin B_6 , adermin or	Pyridoxine
PyridoxaminePyridoxamineVitamin B_{12} or cobalaminCyanocobalaminVitamin B_{12a} , B_{12b} or hydroxocobamideHydroxocobalaminVitamin B_{12a} NitritocobalaminVitamin B_{12c} NitritocobalaminInositol or meso-inositolPantothenic acidPantothenic acidBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	pyridoxol	-
Vítamin B_{12} or cobalaminCyanocobalaminVítamin B_{12a} , B_{12b} or hydroxocobamideHydroxocobalaminVitamin B_{12a} NitritocobalaminVitamin B_{12c} NitritocobalaminInositol or meso-inositolP-InositolPantothenic acidBiotinBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or t-ascorbic acidAscorbic acid	Pyridoxal	Pyridoxal
Vitamin B_{12a} , B_{12b} or hydroxocobamideHydroxocobalaminVitamin B_{12c} NitritocobalaminInositol or meso-inositol Ψ -InositolPantothenic acidPantothenic acidBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	Pyridoxamine	Pyridoxamine
hydroxocobamide Vitamin B_{12c} Nitritocobalamin Inositol or meso-inositol Ψ -Inositol Pantothenic acid Pantothenic acid Biotin Biotin Choline Choline p-Aminobenzoic acid p-Aminobenzoic acid Ascorbic acid	Vitamin B_{12} or cobalamin	Cyanocobalamin
$\begin{array}{llllllllllllllllllllllllllllllllllll$	Vitamin B_{12a} , B_{12b} or	Hydroxocobalamin
Inositol or meso-inositol Pantothenic acidΨ-Inositol Pantothenic acidBiotin Choline p-Aminobenzoic acidBiotin Biotin Choline p-Aminobenzoic acidVitamin C or L-ascorbic acidAscorbic acid		
Pantothenic acidPantothenic acidBiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	Vitamin B _{12c}	Nitritocobalamin
BiotinBiotinCholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acid	Inositol or meso-inositol	Ψ-Inositol
CholineCholinep-Aminobenzoic acidp-Aminobenzoic acidVitamin C or L-ascorbicAscorbic acidacidAscorbic acid	Pantothenic acid	
<i>p</i> -Aminobenzoic acid Vitamin C or L-ascorbic acid		
Vitamin C or L-ascorbic Ascorbic acid		
acid		
L-dehydroascorbic acid Dehydroascorbic acid		Ascorbic acid
	L-dehydroascorbic acid	Dehydroascorbic acid

* Details of the nomenclature for these and other naturally occurring quinones should follow the Tentative Rules of the IUPAC-IUB Commission on Biochemical Nomenclature (see *Biochem. J.* (1967), **102**, 15).

Generic descriptors. The terms vitamin A, vitamin C and vitamin D may still be used where appropriate for example in phrases such as 'vitamin A deficiency', 'vitamin D activity'.

Vitamin E. The term vitamin E should be used as the descriptor for all tocol and tocotrienol derivatives exhibiting qualitatively the biological activity of α -tocopherol. The term **tocopherols** should be used as the generic descriptor for all methyl tocols. Thus, the term **tocopherol** is not synonymous with the term **vitamin** E.

Vitamin K. The term vitamin K should be used as the generic descriptor for 2-methyl-1,4-naphthoquinone (menaphthone) and all derivatives exhibiting qualitatively the biological activity of phylloquinone (phytylmenaquinone).

Niacin. The term **niacin** should be used as the generic descriptor for pyridine 3-carboxylic acid and derivatives

exhibiting qualitatively the biological activity of nicotinamide.

Folic acids. The term **folic acid** may be used to designate the naturally occurring pteroylglutamine acids.

Vitamin \hat{B}_6 . The term vitamin \hat{B}_6 should be used as the generic descriptor for all 2-methylpyridine derivatives exhibiting qualitatively the biological activity of pyridoxine.

Vitamin B_{12} . The term vitamin B_{12} , should be used as the generic descriptor for all corrinoids exhibiting qualitatively the biological activity of cyanocobalamin. The term corrinoids should be used as the generic descriptor for all compounds containing the corrin nucleus and thus chemically related to cyanocobalamin. The term corrinoid is not synonymous with the term vitamin B_{12} .

Amounts of vitamins and summation. All amounts of vitamins should be expressed in terms of their mass rather than in terms of i.u. See Metric Units, Conversion Factors and Nomenclature in Nutritional and Food Sciences. London: The Royal Society, 1972 (paras. 8 and 14-20).

Nomenclature of Fatty Acids. In the description of results obtained for the analysis of fatty acids by conventional gas-liquid chromatography, the shorthand designation proposed by Farquhar, J. W., Insull, W., Rosen, P., Stoffel, W. & Ahrens, E. H. (Nutr. Rev. (1959), 17, Suppl.) for individual fatty acids should be used in the text, tables and figures. Thus 18:1 should be used to represent a fatty acid with eighteen carbon atoms and one double bond; if the position and configuration of the double bond is unknown, this fatty acid should not be referred to as oleic acid. The shorthand designation should also be used in the synopsis but sentences should be constructed so that it is clear to the non-specialist reader that 18:1 refers to a fatty acid; for example, '... resulted in an increase in the concentration of the fatty acid 18:1 in the liver triglycerides...'. If the positions and configurations of the double bonds are known, and these are important to the discussion, then a fatty acid such as linoleic acid may be referred to as $18:2 \Delta 9$ -cis, 12-cis (positions of double bonds related to carbon atom 1). However, when essential and related fatty acids derived from animal tissues are being considered, it is preferable to refer to fatty acids such as linoleic acid as 18:2w6 and arachidonic acid as $20:4\omega6$ (position of double bonds related to the ω -carbon atom); it is assumed that the double bonds are methylene-interrupted and are of the cis-configuration (see Holman, R. T. in Progress in the Chemistry of Fats and Other Lipids, vol. 9, part 1, p. 3. Oxford: Pergamon Press, 1966). Groups of fatty acids that have a common chain length but vary in their double bond content or double bond position should be referred to, for example, as C₂₀ fatty acids or C20 polyunsaturated fatty acids. Impure samples of fatty acids such as those used in the preparation of diets should be referred to, for example as 'linoleic acid'.

Nomenclature of Enzymes. The nomenclature should be that of the Recommendations of the Commission on Enzymes of the International Union of Biochemistry, 1972 (*Enzyme Nomenclature*. Amsterdam: Elsevier Publishing Co., 1973). Relevant *EC* numbers should be given.

Nomenclature of Micro-organisms. The correct name of the organism, conforming with international rules of nomenclature, must be used: if desired, synonyms may be added in brackets when the name is first mentioned. Names of bacteria must conform with the current Bacteriological Code and the opinions issued by the International Committee on Systematic Bacteriology. Names of algae and fungi must conform with the current International Code of Botanical Nomenclature. Names of protozoa must conform with the current International Code of Zoological Nomenclature. The following books may be found useful:

- Bergey's Manual of Determinative Bacteriology, 8th edn (1974), edited by R. E. Buchanan and N. E. Gibbons. Baltimore: The Williams and Wilkins Co.
- The Yeasts, a Taxonomic Study, 2nd edn (1970), edited by J. Lodder. Amsterdam: North Holland Publishing Co.
- Ainsworth and Bisby's Dictionary of the Fungi, 6th edn (1971). Kew: Commonwealth Mycological Institute.

Other Nomenclature, Symbols and Abbreviations. Authors should follow current numbers of the British Journal of Nutrition in this connexion. The IUPAC Rules on chemical nomenclature should be followed, and the Recommendations of the IUPAC-IUB Commission on Biochemical Nomenclature (see Biochem. J. (1978) 169, 11). The symbols and abbreviations, other than units, are essentially those listed in British Standard 1991: 1967 Letter Symbols, Signs and Abbreviations. Part I. General (incorporating amendment no. 1, issued 7 July 1970). Day should be abbreviated to d, for example 7d; except for example, 'each day', '7th day' and 'day 1'.

When an element, e.g. nitrogen, is referred to frequently, the first mention, both in the synopsis and the main text, should be in full, after that by its chemical symbol N. Well-known abbreviations for chemical substances may be used without explanation, thus: RNA for ribonucleic acid and DNA for deoxyribonucleic acid. Other substances that are mentioned frequently may also be abbreviated, the abbreviation being placed in parentheses at the first mention, thus: free fatty acids (FFA) after that, FFA. Well-known terms which are used frequently may be abbreviated in a similar fashion (the abbreviation being in small capitals (underlined twice in the typescript)), thus: basal metabolic rate or basal metabolism (BMR); biological value (BV); dry matter (DM); metabolizable energy (ME). Abbreviations in tables must be defined in footnotes. Signs for footnotes should be used in the sequence: *[†]‡_{||}§¶, then ** etc., (omit * or †, or both, from the sequence if they are used to indicate levels of significance).

Spectrophotometric terms and symbols are those proposed in *IUPAC Manual of Symbols and Terminology* for *Physicochemical Quantities and Units* (1975) London: Butterworths). For mathematical notation and numerals the rules laid down in *Proc. R. Soc.* A (1909), **82**, 14 should be followed. The attention of authors is particularly drawn to the following symbols: m (= milli) =

 10^{-3} , μ (= micro) = 10^{-6} , n (= nano) = 10^{-9} and p (= pico) = 10^{-12} . Note also that ml (millilitre) should be used instead of cc, μ m (micrometre) instead of μ (micron) and μ g (microgram) instead of γ .

Numbers. Figures should be used with units, for example 10 g, 7 d, 4 years (except when beginning a sentence, thus 'Four years ago...'); otherwise, words (except when 100 or more), thus: one man, ten ewes, ninety-nine flasks, three times (but with decimal 2.5, times), 100 patients, 120 cows, 136 samples.

Ethics of Human Experimentation. The notice of contributors is drawn to the guide-lines in the Declaration of Helsinki (1964) (Br. Med. J. (1964), ii 177) and to the Report of ELSE as reprinted in Br. J. Nutr. (1973), 29, 149. A paper describing any experimental work on human subjects should include a statement that the Ethical Committee in the Institution in which the work was performed, where such a Committee exists, has approved it. A paragraph headed *Ethical considerations* in which the experiments are discussed and justified from an ethical standpoint should form the last paragraph of the Experimental section.

Proofs. Proofs are sent to authors in order that they may make sure that the paper has been correctly set up in type and not that they may add new material. Otherwise, increased printing charges are inevitable. Excessive alteration may have to be disallowed or made at the author's expense. Authors will not receive the typescript of their paper with the proof. The symbols used to indicate corrections should be those laid down in *British Standard* 5261 C: 1976. All corrections should be made in ink in the margins; mark made in the text should be only those indicating the place to which the correction refers.

Corrected proofs should be returned without delay to Miss Margot Lee, *British Journal of Nutrition*, Department of Biochemistry, University of Liverpool, PO Box 147, Liverpool L69 3BX.

Reprints. Orders for reprints should be addressed to The University Press, Cambridge. Where at least one author of a paper is a member of The Nutrition Society, twenty-five reprints are supplied free of cost. Additional reprints may be purchased if the Press are notified on the appropriate form immediately the proof is received.