

caused ballismus movements in this case. In patients who present with short duration monoballismus, evaluation for subthalamic nuclei function, seizure disorders and other origins of ballismus are warranted.

FUNDING ACKNOWLEDGEMENTS: Smell & Taste Treatment and Research Foundation

127

Cardiovascular Safety Assessment of Deutetrabenazine in Healthy Volunteers and Implications for Patients With Huntington Disease or Tardive Dyskinesia

Donna S. Cox, PhD¹; Micha Levi, PhD¹; Laura Rabinovich-Guilatt¹; David Truong, PharmD, MS¹; and David Stamler, MD¹

¹ Teva Pharmaceutical Industries, Frazer, Pennsylvania, USA

ABSTRACT: Introduction: Deutetrabenazine is approved for treating Huntington disease (HD) chorea and is being evaluated for tardive dyskinesia (TD).

OBJECTIVE: To assess the effect of deutetrabenazine on cardiac repolarization.

METHODS: A QT interval study was performed to evaluate effects of deutetrabenazine 12 and 24 mg on cardiac repolarization, as assessed by time-matched change from baseline, placebo-adjusted, in Fridericia-corrected QT interval ($\Delta\Delta$ QTcF). Moxifloxacin (400 mg) and tetrabenazine (50 mg) were the positive control and comparator, respectively. An exposure–response analysis was developed from this study to predict maximal effects on QTcF at maximum recommended dosing based on CYP2D6 status, an approach consistent with regulatory guidance at predicting QT interval effects.

RESULTS: Maximal $\Delta\Delta$ QTcF between the least-squares mean (90% two-sided confidence interval) of deutetrabenazine 12 and 24 mg ($n = 45$ in each group) were 2.8 (0.7–4.8) ms and 4.5 (2.4–6.5) ms, respectively. The $\Delta\Delta$ QTcF increase with tetrabenazine ($n = 45$) was 7.6 (5.6–9.5) ms. Assay sensitivity was verified with moxifloxacin ($n = 47$), which produced a maximal effect on $\Delta\Delta$ QTcF of 14.0 (11.9–16.0) ms. A linear model was developed that described a correlation between plasma concentrations from pivotal HD and TD trials ($n = 101$) and QT interval prolongation. Using that model and the individual predicted C_{max} for HD and TD patients, the placebo-adjusted change from baseline in QTcF for deutetrabenazine at maximal recommended daily doses was found to be 5.4 (2.5–9.5) ms.

CONCLUSIONS: Patients receiving the maximal recommended doses of deutetrabenazine are predicted to have a QTcF increase below the level of regulatory concern.

Presented at: Psych Congress; September 16–19, 2017; New Orleans, Louisiana, USA.

FUNDING ACKNOWLEDGEMENTS: This study was funded by Teva Pharmaceutical Industries, Petach Tikva, Israel

128

Effect of DR/ER-MPH on Early Morning and Late Afternoon/Evening Functioning in Children With ADHD: Analysis of PREMB-R Items From a Phase 3 Trial

Steven R. Pliszka, MD¹; Valerie K. Arnold, MD²; Andrea Marraffino, PhD³; Norberto J. DeSousa, MA⁴; Bev Incedon, PhD⁴; F. Randy Sallee, MD, PhD⁴; Timothy E. Wilens, MD⁵; and Jeffrey H. Newcorn, MD⁶

¹ The University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

² University of Tennessee Health Science Center, Memphis, Tennessee, USA

³ Meridien Research, Inc., Maitland, Florida, USA

⁴ Ironshore Pharmaceuticals & Development, Inc., Camana Bay, Grand Cayman, Cayman Islands

⁵ Massachusetts General Hospital, Boston, Massachusetts, USA

⁶ Mount Sinai Medical Center, New York, New York, USA; On behalf of the HLD200-108 Study Group

ABSTRACT: Objective: In a phase 3 trial of children with ADHD, DR/ER-MPH (formerly HLD200), a delayed-release and extended-release methylphenidate, improved ADHD symptoms and reduced at-home early morning and late afternoon/evening functional impairments versus placebo, as measured by the validated Parent Rating of Evening and Morning Behaviors-Revised, Morning (PREMB-R AM) and Evening (PREMB-R PM) subscales. This post hoc analysis evaluated the effect of DR/ER-MPH versus placebo on individual PREMB-R AM/PM item scores.

METHOD: Data were analyzed from a pivotal, randomized, double-blind, multicenter, placebo-controlled, parallel-group, phase 3 trial of DR/ER-MPH in children (6–12 years) with ADHD (NCT02520388). Using the 3-item PREMB-R AM and 8-item PREMB-R PM, both key secondary endpoints, investigators evaluated early morning and late afternoon/evening functional impairment by scoring each item on a severity scale from 0 (none) to 3 (a lot). For post hoc analyses, treatment comparisons between DR/ER-MPH and placebo at endpoint were determined by using least squares mean changes from