

Effects of psychological eating behaviour domains on the association between socio-economic status and BMI

Antje Löffler^{1,2,3,*}, Tobias Luck^{1,2}, Francisca S Then^{1,2}, Claudia Luck-Sikorski^{1,4,5}, Alexander Pabst¹, Peter Kovacs⁴, Yvonne Böttcher^{4,6}, Jana Breitfeld⁴, Anke Tönjes⁷, Annette Horstmann^{4,8}, Markus Löffler⁹, Christoph Engel⁹, Joachim Thiery¹⁰, Arno Villringer^{8,11}, Michael Stumvoll⁴ and Steffi G Riedel-Heller¹

¹Institute of Social Medicine, Occupational Health and Public Health (ISAP), University of Leipzig, Leipzig, Germany: ²LIFE–Leipzig Research Center for Civilization Diseases, University of Leipzig, Leipzig, Germany: ³Brandenburg University of Technology, Department of Public Health, Universitätsplatz 1, 01968 Senftenberg, Germany: ⁴IFB Adiposity Diseases, Leipzig University Medical Center, Leipzig, Germany: ⁵University of Applied Health Sciences, Gera, Germany: ⁶Institute of Clinical Medicine, Section for Clinical Molecular Biology, University of Oslo, Oslo, Norway: ⁷Division of Endocrinology and Nephrology, Medical Department, University of Leipzig, Leipzig, Germany: ⁸Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany: ⁹Institute for Medical Informatics, Statistics and Epidemiology (IMISE), University of Leipzig, Leipzig, Germany: ¹⁰Institute of Laboratory Medicine, Clinical Chemistry and Molecular Diagnostics, University of Leipzig, Leipzig, Germany: ¹¹Day Clinic of Cognitive Neurology, University of Leipzig, Leipzig, Germany

Submitted 16 December 2016: Final revision received 5 June 2017: Accepted 12 June 2017: First published online 24 July 2017

Abstract

Objective: The current study investigates potential pathways from socio-economic status (SES) to BMI in the adult population, considering psychological domains of eating behaviour (restrained eating, uncontrolled eating, emotional eating) as potential mediators stratified for sex.

Design: Data were derived from the population-based cross-sectional LIFE-Adult-Study. Parallel-mediation models were conducted to obtain the total, direct and indirect effects of psychological eating behaviour domains on the association between SES and BMI for men and for women.

Setting: Leipzig, Germany.

Subjects: We studied 5935 participants aged 18 to 79 years.

Results: Uncontrolled eating mediated the association between SES and BMI in men only and restrained eating in both men and women. Emotional eating did not act as mediator in this relationship. The total effect of eating behaviour domains on the association between SES and BMI was estimated as $\beta = -0.03$ (SE 0.02; 95% CI $-0.062, -0.003$) in men and $\beta = -0.18$ (SE 0.02; 95% CI $-0.217, -0.138$) in women.

Conclusions: Our findings do not indicate a strong overall mediation effect of the eating behaviour domains restrained eating, uncontrolled eating and emotional eating on the association between SES and BMI. Further research on other pathways of this association is strongly recommended. Importantly, our findings indicate that, independent from one's social position, focusing on psychological aspects in weight reduction might be a promising approach.

Keywords
Socio-economic status
BMI
Eating behaviour
Restrained eating
Uncontrolled eating
Emotional eating

In Western countries, high standards of living provide good conditions for health, formally irrespective of one's social position. Despite these good theoretical conditions, there is an inverse gradient between socio-economic status (SES) and health. For example, in Germany, the estimated risk for diabetes mellitus type 2, heart disease or liver disease is twice as high for individuals with the lowest SES compared with individuals with the highest SES⁽¹⁾.

Moreover, risk factors for common civilization diseases like physical inactivity and higher smoking prevalence, as well as less use of primary prevention services are more common in population groups with lower and lowest SES. This is also true for the risk factor high BMI^(1,2). For example, the German Health Update 2012 for the German population aged between 18 and 79 years revealed that 31.4% of women with the lowest SES were identified as

*Corresponding author: Email antje.loeffler@b-tu.de

overweight (BMI ≥ 25.0 kg/m²) and another 23.6% as obese (BMI ≥ 30.0 kg/m²). But in women with the highest SES, the estimated overweight and obesity prevalence was much lower: 24.2% of the women were overweight and another 9.4% were obese⁽³⁾. For men, the findings are less clear. Nevertheless, obesity prevalence was also significantly higher in men with the lowest SES than in men with the highest SES (20.0 v. 14.1% in the age group 18 to 79 years)⁽³⁾. Overweight and obesity carry high social and economic burden. For example, in Germany, direct costs of obesity were estimated as €29.39 billion and the indirect costs as €33.65 billion per year⁽⁴⁾. Therefore, it is of utmost importance to identify the underlying links between SES and BMI, which might help to develop suitable prevention strategies against overweight and obesity.

One link between SES and BMI is eating behaviour, which differs between the SES groups. For example, higher consumption of sugar-sweetened beverages is more common in the lower and lowest SES groups^(5,6). Regarding the various interacting factors which lead to one's individual eating behaviour, including genetic, sociocultural and personal ones, there is no simple explanation for the differences in eating behaviour between SES groups. Therefore, we hypothesized that psychological factors of eating behaviour might be underlying factors between the inverse association of SES and BMI. Common and well-established psychological domains of eating behaviour are: (i) a domain which describes the degree of cognitive control in daily food intake (cognitive restrained, restrained eating); (ii) a domain which describes the loss of control over food intake (uncontrolled eating, disinhibition); and (iii) a domain which describes eating in response to emotional triggers like feeling anxious or depressed (emotional eating)⁽⁷⁻¹³⁾. The association between these psychological domains of eating behaviour and BMI are well studied. Especially the domains uncontrolled eating and emotional eating have been found to be associated with higher BMI values⁽¹⁴⁻¹⁷⁾. As eating behaviour has not only been found to be associated with BMI, but also with SES, in the present study we particularly aimed at investigating potential pathways between SES and the psychological eating behaviour domains 'restrained eating', 'uncontrolled eating' and 'emotional eating' and the BMI in the adult population. To the very best of our knowledge, this has not been done before.

Methods

Ethics statement

All participants provided written informed consent prior to their participation in the study. The study complies with the ethical standards of the Declaration of Helsinki and has been approved by the ethics committee of the University of Leipzig, Germany.

Participants

Data were derived from the large population-based LIFE-Adult-Study that was conducted by the LIFE-Leipzig Research Center for Civilization Diseases in Leipzig, Germany (August 2011–November 2014). The LIFE-Adult-Study aims to examine the prevalence, early onset markers, genetic predisposition and the role of lifestyle factors in major civilization diseases such as obesity, diabetes and CVD. The study design of the LIFE-Adult-Study has been described in detail elsewhere⁽¹⁸⁾. An age- and sex-stratified sample of residents of the city of Leipzig, Germany, was randomly selected from lists provided by the Leipzig registry office. The response rate was 33%. Eating behaviour was completely assessed in 6657 participants. We excluded participants whose eating behaviour might have been influenced by symptoms of chronic inflammatory bowel disease, treatment of cancer disease, insulin treatment or intake of antipsychotic drugs (n 722). The present analysis was based on a final sample of 5935 participants aged 18–79 years.

Data collection and assessment procedures

All participants underwent a comprehensive assessment programme including a variety of physical and medical examinations, clinical interviews and standardized questionnaires. The assessments were conducted by trained study personnel at the LIFE research centre located on the premises of the University Hospital of Leipzig. For the present report, data were analysed from the following assessments.

Eating behaviour

Eating behaviour was assessed with the Fragebogen zum Essverhalten (FEV), the German version of the Three-Factor Eating Questionnaire (TFEQ)^(8,12). The present analysis was based on a revised twenty-nine-item version of the TFEQ, which covers three domains of eating behaviour: 'uncontrolled eating', 'restrained eating' and 'emotional eating'. The internal consistency of the scales (Cronbach's α) was stated as 0.802 for the domain of uncontrolled eating (contains fifteen items), 0.840 for the domain of restrained eating (contains eleven items) and 0.780 for the domain of emotional eating (contains three items)⁽¹⁷⁾.

As mentioned above, the domain of uncontrolled eating describes the tendency to lose control over food intake, the domain of restrained eating describes a permanent cognitive control of food intake order to lose or to control body weight, and the domain of emotional eating describes eating in response to emotional triggers like feeling anxious or depressed.

Sociodemographic characteristics

Sociodemographic information was obtained via a standardized interview, which included questions about the educational and occupational qualification of the participant, the net household income and actual occupational status. This information was used to assess the multidimensional

SES-Index-Score in accordance with the approach used in the German Health Interview and Examination Survey for Adults⁽¹⁹⁾. The SES-Index-Score is based on the three domains of 'education', 'work' and 'income'. For each participant of the LIFE-Adult-Study, each of the three domains was rated between 1 point as the lowest possible value and 7 points as the highest possible value. The points of the three domains were then added to a final score (range 3–21 points) and participants were categorized into one of the three SES groups: 'low' (3–8 points), 'medium' (9–14 points) or 'high' (15–21 points).

Anthropometry

Body weight was measured with an electronic scale (SECA 701, Seca GmbH & Co. KG) with a precision of 0.01 kg. Body height was assessed by means of a stadiometer (SECA 240) to the nearest 0.1 cm. BMI was calculated as the weight in kilograms divided by the squared height in metres (kg/m²).

Statistical analysis

Statistical analyses were performed with the statistical software package IBM SPSS Statistics version 20.0, including the SPSS macro 'PROCESS' from Preacher and Hayes⁽²⁰⁾. All analysis employed an α level of 0.05 for statistical significance (two-tailed). Group differences were analysed using Mann–Whitney *U* tests, χ^2 tests and Kruskal–Wallis tests as appropriate.

To examine the pathways between SES, BMI and the eating behaviour domains, we conducted mediation analysis. For our analyses, we hypothesized that the psychological domains of eating behaviour (uncontrolled eating, restrained eating, emotional eating) mediate the association between SES-Index-Score and BMI. Previous analysis required differences in eating behaviour between men and women. Therefore, we conducted sex-stratified analysis⁽²¹⁾. We also used age as covariate, as age was found to be an influencing factor on eating behaviour⁽²¹⁾.

To detect the presence of mediation effects, we first examined the associations between the obtained variables. Therefore, we conducted Pearson's correlation analysis. Then we developed a parallel-mediation model to examine the potential direct, indirect and total effects of eating behaviour on the association between SES-Index-Score and BMI, controlled for covariates. Parallel mediation was used to simultaneously include the three eating behaviour domains in one model. A 'direct effect' describes the direct association between SES-Index-Score and BMI in the presence of the mediator(s). An 'indirect effect' describes the product of the paths between SES-Index-Score and BMI and between SES-Index-Score, the mediator(s) and BMI. The 'total effect' is the sum of the indirect and direct effect. Thus, a full mediation of the association between SES-Index-Score and BMI by eating behaviour would be present if there was an indirect effect via eating behaviour, but no direct effect of SES-Index-Score

on BMI in the model. A partial mediation would be present if both the path between SES-Index-Score and BMI and the path between SES-Index-Score, eating behaviour and BMI was significant. To achieve scale similarity, we conducted a *z*-transformation of all included variables. A *z*-scale has an arithmetic mean of 0 and an sd of 1, and allows the interpretation of regression coefficients more easily. Significance of the direct and indirect effects was tested with both the Sobel *z*-test and a bootstrapping procedure of about 10 000 resamples. The bootstrap procedure providing 95% CI is additionally recommended for significance testing as it respects irregularity of the sampling distribution and provides more power⁽²⁰⁾.

Results

Sample description

Overall, analysis of the association between SES, eating behaviour and BMI was based on a sample of 5935 participants of the LIFE-Adult-Study. Characteristics of the study sample are shown in Table 1.

Table 2 presents a detailed overview on the association between the factors BMI and eating behaviour with SES. Group comparisons using Kruskal–Wallis tests revealed a higher mean BMI in lower SES groups, especially in women (χ^2 for men = 0.848, $P = 0.654$; χ^2 for women = 87.137, $P < 0.001$). Additional information about the proportion of obesity in SES classes is given. The proportion of obesity was higher in lower and lowest SES in both men and women compared with the highest SES group. Values of restrained eating were higher in the higher SES groups in both men and women (χ^2 for men = 71.304, $P < 0.001$; χ^2 for women = 20.241, $P < 0.001$). Values of uncontrolled eating and emotional eating did not differ between lowest and highest SES groups.

Intercorrelation between SES-Index-Score, BMI, eating behaviour and the covariates

As shown in Table 3, we found weak (Pearson's correlation coefficient $r < 0.3$) but significant associations between SES-Index-Score and BMI for both men and women. We also found weak associations between SES-Index-Score and all three eating behaviour domains as well as between BMI and the eating behaviour domains. Finally, we found a moderate ($r > 0.4$ and < 0.7) association between the eating behaviour domains uncontrolled and emotional eating, and a weak association between the domains emotional and restrained eating.

Parallel-mediation model

We tested the pathways between SES-Index-Score, BMI and eating behaviour in separate parallel-mediation models for men and for women (see Figs. 1 and 2). In the models, the psychological eating behaviour domains uncontrolled eating, restrained eating and emotional

Table 1 Main characteristics of and significance differences between the study sample of men and women (*n* 5935), LIFE-Adult-Study, Leipzig, Germany, August 2011–November 2014

	Men (<i>n</i> 2905)		Women (<i>n</i> 3030)		<i>P</i> value†
	Mean or %	SD or <i>n</i>	Mean or %	SD or <i>n</i>	
Sociodemographic characteristics					
Age					
Age (years; range 18–79)	54.8	12.0	53.8	11.1	<0.001
Socio-economic status (SES)					
SES-Index-Score (range 3–21)	13.5	3.3	13.1	3.1	<0.001
Low SES (% , <i>n</i>)	4.8	139	4.3	129	<0.001
Middle SES (% , <i>n</i>)	49.5	1439	57.4	1740	
High SES (% , <i>n</i>)	45.7	1327	38.3	1161	
Anthropometric measures					
BMI (kg/m ²)	27.4	4.08	26.8	5.40	<0.001
Eating behaviour domains‡					
Restrained eating score (range 0–15)	4.6	3.52	6.7	3.90	<0.001
Uncontrolled eating score (range 0–11)	2.7	2.53	3.1	2.80	<0.001
Emotional eating score (range 0–3)	0.2	0.60	0.6	0.99	<0.001

†Group differences in continuous variables were analysed with the Mann–Whitney *U* test, group differences in categorical variables were analysed with the χ^2 test.

‡Higher scores indicate stronger characteristic values in the domains.

Table 2 Differences in BMI and eating behaviour scores according to socio-economic status (SES) for men and women (*n* 5935) aged 18–79 years, LIFE-Adult-Study, Leipzig, Germany, August 2011–November 2014

	Low SES		Medium SES		High SES		χ^2	<i>P</i> value†
	Mean	SD	Mean	SD	Mean	SD		
Men								
BMI	27.65	5.17	27.45	4.14	27.27	3.88	0.848	0.654
Obese‡	28.80		24.30		20.90		7.289	0.026
Restrained eating score§	3.36	3.26	4.22	3.42	5.12	3.56	71.304	<0.001
Uncontrolled eating score§	2.53	2.48	2.64	2.50	2.87	2.56	7.858	0.020
Emotional eating score§	0.18	0.54	0.17	0.52	0.23	0.66	2.719	0.257
Women								
BMI	28.63	6.13	27.37	5.49	25.79	4.99	87.137	<0.001
Obese‡	36.40		26.70		17.10		47.917	<0.001
Restrained eating score§	5.26	3.78	6.76	3.94	6.84	3.82	20.241	<0.001
Uncontrolled eating score§	2.89	2.90	3.08	2.79	3.21	2.80	3.684	0.159
Emotional eating score§	0.62	1.01	0.56	0.96	0.66	1.04	5.948	0.051

†Group differences were analysed with the Kruskal–Wallis *H* test.

‡Proportion of obesity in %; obesity is classified as BMI ≥ 30.0 kg/m²(27).

§Higher scores indicate stronger characteristic values in the domains.

Table 3 Pearson's correlation coefficients between socio-economic status (SES; i.e. SES-Index-Score), BMI, eating behaviour and age for men and women (*n* 5935) aged 18–79 years, LIFE-Adult-Study, Leipzig, Germany, August 2011–November 2014

	BMI	RS	UC	EE	Age
Men					
SES	-0.04*	0.15*	0.07*	0.06*	0.01*
BMI		0.14*	0.32*	0.15*	0.17*
RS			0.00*	0.08*	0.22*
UC				0.37*	-0.14*
EE					-0.07*
Women					
SES	-0.18*	0.06*	0.01*	0.04*	-0.11*
BMI		0.15*	0.25*	0.22*	0.23*
RS			-0.01*	0.09*	0.20*
UC				0.48*	-0.14*
EE					-0.07*
Age					

RS, restrained eating; UC, uncontrolled eating; EE, emotional eating.

**P* < 0.05.

eating were included as potential mediators between SES-Index-Score and BMI. Age was included as a covariate. Results of the mediation models are presented in Table 4.

In both men and women, we found a significant direct path between SES-Index-Score and BMI. The effect was estimated as $\beta = -0.07$ (SE 0.01; *P* < 0.001) for men and $\beta = -0.19$ (SE 0.02; *P* < 0.001) for women. Regarding the paths between SES-Index-Score, eating behaviour variables and BMI in detail, we found differences between the models for men and women. For men, we found direct significant paths between SES-Index-Score and all three eating behaviour domains (restrained eating: $\beta = 0.13$ (SE 0.02), *P* < 0.001; uncontrolled eating: $\beta = 0.06$ (SE 0.02), *P* < 0.05; emotional eating: $\beta = 0.04$ (SE 0.01), *P* < 0.05). The paths between the eating behaviour domains and BMI were also significant (restrained eating: $\beta = 0.09$ (SE 0.02), *P* < 0.001; uncontrolled eating: $\beta = 0.30$ (SE 0.02), *P* < 0.001;




Fig. 1 Parallel-mediation model between socio-economic status (SES; i.e. SES-Index-Score), BMI and psychological domains of eating behaviour for men (*n* 2905) aged 18–79 years, LIFE-Adult-Study, Leipzig, Germany, August 2011–November 2014. *Significant path (z-test) with α level of < 0.05; **significant path (z-test) with α level of < 0.001




Fig. 2 Parallel-mediation model between socio-economic status (SES; i.e. SES-Index-Score), BMI and psychological domains of eating behaviour for women (*n* 3030) aged 18–79 years, LIFE-Adult-Study, Leipzig, Germany, August 2011–November 2014. **Significant path (z-test) with α level of < 0.001

Table 4 Results from the mediation models with direct, indirect and total effects of socio-economic status (SES; i.e. SES-Index-Score) and eating behaviour on BMI for men ($r^2 = 0.032$) and women ($r^2 = 0.075$) aged 18–79 years, LIFE-Adult-Study, Leipzig, Germany, August 2011–November 2014

	β	SE	<i>P</i> value (normal theory)	Bootstrap 95% CI	
Men					
Total effect†	-0.033	0.015	0.032	-0.062	-0.003
Direct effect‡	-0.066	0.014	<0.001	-0.094	-0.038
Indirect effect§					
Restrained eating score	0.012	0.003	<0.001	0.008	0.018
Uncontrolled eating score	0.016	0.005	0.003	0.009	0.030
Emotional eating score	0.002	0.001	0.066	0.000	0.006
Women					
Total effect†	-0.178	0.020	<0.001	-0.217	-0.138
Direct effect‡	-0.191	0.019	<0.001	-0.228	-0.153
Indirect effect§					
Restrained eating score	0.011	0.003	0.002	0.006	0.027
Uncontrolled eating score	-0.004	0.005	0.909	-0.010	0.009
Emotional eating score	0.003	0.003	0.144	-0.001	0.001

†Total effect = direct effect + indirect effects.

‡Direct effect = path between SES-Index-Score and BMI in presence of mediators.

§Indirect effect = paths between SES-Index-Score, eating behaviour domains and BMI.

emotional eating: $\beta = 0.05$ (SE 0.02), $P < 0.05$). For women, there were no significant paths between SES-Index-Score and the eating behaviour domains uncontrolled eating and emotional eating, but between SES-Index-Score and the domain restrained eating ($\beta = 0.09$ (SE 0.02), $P < 0.001$).

Similarly to the results for men, however, all three eating behaviour variables showed significant paths to BMI also in women (restrained eating: $\beta = 0.12$ (SE 0.02), $P < 0.001$; uncontrolled eating: $\beta = 0.25$ (SE 0.02), $P < 0.001$; emotional eating: $\beta = 0.11$ (SE 0.02), $P < 0.001$). The total effect

between SES-Index-Score and BMI regarding eating behaviour variables was estimated as $\beta = -0.03$ (SE 0.02; 95% CI -0.062, -0.003) for men and as $\beta = -0.18$ (SE 0.02; 95% CI -0.217, -0.138) for women.

Overall, the mediation models indicated a partial mediation by psychological eating behaviour domains in both men and women. In men, restrained eating ($\beta = 0.012$ (SE 0.003); 95% CI 0.008, 0.018) and uncontrolled eating ($\beta = 0.016$ (SE 0.005); 95% CI 0.009, 0.030) mediated the association between SES-Index-Score and BMI. In women, a mediation effect of restrained eating between SES-Index-Score and BMI was found ($\beta = 0.011$ (SE 0.003); 95% CI 0.006, 0.027). Emotional eating did not act as a mediator between SES-Index-Score and BMI in both men and women; results rather showed an independent impact of emotional eating on BMI.

Discussion

In the present study, we aimed to investigate potential pathways from SES to BMI in the adult population, considering the psychological eating behaviour domains restrained eating, uncontrolled eating and emotional eating as potential mediators stratified for sex. We hypothesized that psychological domains of eating behaviour mediate the association between SES and BMI.

In detail, the eating behaviour factor restrained eating was found to attenuate the indirect association between SES-Index-Score and BMI. This means that low restrained eating might explain higher BMI in lower SES. Higher SES is associated with higher restrained eating. Importantly, restrained eating might be a strong result of individual health considerations. It has been shown that highly educated individuals have higher health considerations, which might in turn lead to restrained eating to stop weight gain, to maintain weight or to lose weight⁽²²⁾. Thus, this result corroborates health awareness as one important component in weight reduction programmes or in part of health education in general.

We also found a significant mediation effect of uncontrolled eating on the association between SES-Index-Score and BMI. However, as this effect was found only in men, and moreover was only small, further studies may be waited for to confirm this finding. Nevertheless, as the direct association between uncontrolled eating and BMI is well studied, coping with a loss of control in food intake might be part of a weight reduction programme^(14–17).

Emotional eating did not mediate the association between SES-Index-Score and BMI in both men and women. Thus, our analysis suggests that emotional eating is rather directly associated with BMI – independent of SES. The positive association between emotional eating and BMI was also found in several other studies and underlines the importance of this eating behaviour domain for understanding weight gain^(13,15–17,23,24). Therefore, alternative

coping strategies for emotions might be a very useful approach in attempts to lose or to maintain weight independent from one's social position.

However, regarding the low total effect of all the eating behaviour domains on the association between SES-Index-Score and BMI in mediation analyses, our study, in general, does not indicate a strong mediation effect of psychological domains of eating behaviour. As we were able to focus only on three domains of eating behaviour, more research is encouraged to investigate further potentially important aspects (e.g. social norms, habits, food marketing strategies) in mediation analyses of the association between SES and BMI. Nevertheless, independent from one's social position, focusing on psychological aspects in a weight reduction process might be a promising approach. Therefore, the identification of internal or external food stimuli, such as food advertising, might help to develop individual coping strategies in a weight reduction process. For example, a review from Mantzios *et al.* found that 'mindfulness eating' might reduce emotional as well as uncontrolled eating, which are both associated with higher BMI values^(17,25).

Our study is not without limitations. First, there is the probability of selection bias because of a relatively low response rate and some missing information about eating behaviour. Second, individuals with lower SES were under-represented, which might have caused bias, too. Finally, self-reported data of eating behaviour might have also led to some bias, such as over- or under-reporting. Nevertheless, irrespective of these limitations, we found that the association between SES and BMI was partly mediated by eating behaviour.

However, the association between SES and BMI, in general, seems to be complex and there might be more than one pathway involved. To the very best of our knowledge, our study is the first that examined a pathway via the psychological domains of eating behaviour. So far, few studies also examined other potential psychological aspects in mediation analysis. Beydoun *et al.*, for example, found that depressive symptoms may mediate the inverse association between SES and obesity in women⁽²⁶⁾.

Acknowledgements

Acknowledgements: The authors would like to thank all participants and the staff at LIFE–Leipzig Research Center for Civilization Diseases. *Financial support:* This publication is supported by LIFE–Leipzig Research Center for Civilization Diseases, Universität Leipzig. LIFE is funded by the European Union through the European Regional Development Fund (ERDF) and by the Free State of Saxony within the framework of the excellence initiative. F.S.T. has been supported by LIFE–Leipzig Research Center for Civilization Diseases, Universität Leipzig. Her collaboration within LIFE was funded by means of the

European Social Fund and the Free State of Saxony. Y.B. is funded by IFB Adiposity Diseases. IFB Adiposity Diseases is supported by the Federal Ministry of Education and Research (BMBF), Germany (grant number FKZ: 01EO1501). *Conflict of interest:* None of the authors has any conflict of interest to declare. *Authorship:* M.L., J.T., C.E., A.V., M.S. and S.G.R.-H. conceived and designed the study. A.L. performed the statistical analysis, interpreted the data and drafted the manuscript. T.L. and F.S.T. helped to interpret the data and to perform the statistical analysis and were involved in drafting and revising the manuscript critically for important contents. A.P. advised in statistical analysis. S.G.R.-H. supervised the study. She as well as M.L., J.T., C.E., A.V., M.S., C.L.-S., A.P., P.K., Y.B., J.B., A.T. and A.H. revised the manuscript critically for important contents. All authors approved the final version. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the ethics committee of the University of Leipzig, Germany. All subjects provided written informed consent prior to their participation in the study.

References

- Lampert T, Kroll LE, Lippe E von der et al. (2013) Socioeconomic status and health: results of the German Health Interview and Examination Survey for Adults (DEGS1). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* **56**, 814–821.
- Weyers S, Dragano N, Richter M et al. (2010) How does socio economic position link to health behaviour? Sociological pathways and perspectives for health promotion. *Glob Health Promot* **17**, 25–33.
- Robert Koch-Institut (editor) (2012) *Daten und Fakten: Ergebnisse der Studie 'Gesundheit in Deutschland aktuell 2010': Gesundheitsberichterstattung-Hefte*. Berlin: Robert Koch-Institut.
- Effertz T, Engel S, Verheyen F et al. (2015) The costs and consequences of obesity in Germany: a new approach from a prevalence and life-cycle perspective. *Eur J Health Econ* **17**, 1141–1158.
- Malik VS, Schulze MB & Hu FB (2006) Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr* **84**, 274–288.
- Weyers S, Fekete C, Dragano N et al. (2013) Education, selected nutrition attitudes and nutrition behaviour. *Prävention und Gesundheitsförderung* **8**, 289–294.
- Ganley RM (1988) Emotional eating and how it relates to dietary restraint, disinhibition, and perceived hunger. *Int J Eat Disord* **7**, 635–647.
- Stunkard AJ & Messick S (1985) The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *J Psychosom Res* **29**, 71–83.
- Hyland ME, Irvine SH, Thacker C et al. (1989) Psychometric analysis of the Stunkard–Messick Eating Questionnaire (SMEQ) and comparison with the Dutch Eating Behavior Questionnaire (DEBQ). *Curr Psychol* **8**, 228–233.
- Karlsson J, Persson LO, Sjostrom L et al. (2000) Psychometric properties and factor structure of the Three-Factor Eating Questionnaire (TFEQ) in obese men and women. Results from the Swedish Obese Subjects (SOS) study. *Int J Obes Relat Metab Disord* **24**, 1715–1725.
- Mazzeo SE, Aggen SH, Anderson C et al. (2003) Investigating the structure of the eating inventory (three-factor eating questionnaire): a confirmatory approach. *Int J Eat Disord* **34**, 255–264.
- Pudel V & Westenhoefer J (1989) *Fragebogen zum Eßverhalten*. Göttingen: Hogrefe.
- van Strien T, Frijters JER, Bergers GPA et al. (1986) The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *Int J Eat Disord* **5**, 295–315.
- French SA, Epstein LH, Jeffery RW et al. (2012) Eating behavior dimensions. Associations with energy intake and body weight. A review. *Appetite* **59**, 541–549.
- Geliebter A & Aversa A (2003) Emotional eating in overweight, normal weight, and underweight individuals. *Eat Behav* **3**, 341–347.
- Koenders PG & van Strien T (2011) Emotional eating, rather than lifestyle behavior, drives weight gain in a prospective study in 1562 employees. *J Occup Environ Med* **53**, 1287–1293.
- Löffler A, Luck T, Then FS et al. (2015) Eating behaviour in the general population: an analysis of the factor structure of the German version of the Three-Factor-Eating-Questionnaire (TFEQ) and its association with the body mass index. *PLoS One* **10**, e0133977.
- Loeffler M, Engel C, Ahnert P et al. (2015) The LIFE-Adult-Study: objectives and design of a population-based cohort study with 10,000 deeply phenotyped adults in Germany. *BMC Public Health* **15**, 691.
- Lampert T, Kroll L, Müters S et al. (2013) Messung des sozioökonomischen Status in der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* **56**, 631–636.
- Preacher KJ & Hayes AF (2008) Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* **40**, 879–891.
- Löffler A, Luck T, Then FS et al. (2015) Age- and gender-specific norms for the German version of the Three-Factor Eating-Questionnaire (TFEQ). *Appetite* **91**, 241–247.
- Ball K, Crawford D & Mishra G (2006) Socio-economic inequalities in women's fruit and vegetable intakes: a multi-level study of individual, social and environmental mediators. *Public Health Nutr* **9**, 623–630.
- Kontinen H, Mannisto S, Sarlio-Lahteenkorva S et al. (2010) Emotional eating, depressive symptoms and self-reported food consumption. A population-based study. *Appetite* **54**, 473–479.
- Horstmann A, Kovacs P, Kabisch S et al. (2013) Common genetic variation near MC4R has a sex-specific impact on human brain structure and eating behavior. *PLoS One* **8**, e74362.
- Mantzios M & Wilson JC (2015) Mindfulness, eating behaviours, and obesity: a review and reflection on current findings. *Curr Obes Rep* **4**, 141–146.
- Boydoun MA, Kuczumski MTF, Mason MA et al. (2009) Role of depressive symptoms in explaining socioeconomic status disparities in dietary quality and central adiposity among US adults: a structural equation modeling approach. *Am J Clin Nutr* **90**, 1084–1095.
- World Health Organization (2000) *Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation. WHO Technical Report Series no. 894*. Geneva: WHO.