

Positive relationship between consumption of specific fish type and *n*-3 PUFA in milk of Hong Kong lactating mothers

Vincy Wing-Si Wong¹, Yuk-Fan Ng¹, Suk-Mei Chan², Yi-Xiang Su³, Kevin Wing-Hin Kwok¹, Hing-Man Chan¹, Chi-Leung Cheung¹, Hang-Wai Lee¹, Wing-Yiu Pak¹, Shi-Ying Li^{1*} and Man-Sau Wong^{1*}

¹Department of Applied Biology and Chemical Technology, Laboratory for Infant & Child Nutrition, Food Safety and Technology Research Centre, The Hong Kong Polytechnic University, Hong Kong, China

²Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, China

³Department of Nutrition, School of Public Health, Guangdong Provincial Key Laboratory of Food, Nutrition and Health, Sun Yat-Sen University, Guangzhou 510080, People's Republic of China

(Submitted 16 July 2018 – Final revision received 14 March 2019 – Accepted 24 March 2019 – First published online 17 June 2019)

Abstract

Residents of Hong Kong have undergone a dietary transition from a traditional Chinese diet that is high in seafood to a more Western diet. This may have affected the nutritional composition of breast milk of Hong Kong mothers. The present study aims to investigate the relationship between the dietary pattern and the fatty acid profile of the breast milk of lactating women in Hong Kong. Seventy-three volunteering healthy Hong Kong lactating mothers participated in the study. Their dietary intakes were assessed by using a 3-d dietary record and FFQ. The mean *n*-3 fatty acid levels were approximately 0.4% (EPA) and 0.9% (DHA) of total fatty acids in the breast milk of lactating mothers who had exclusively breastfed their infants aged 2–6 months. Maternal dietary intakes of *n*-3 fatty acids were positively associated with their levels in the breast milk. The levels of maternal intakes of freshwater and saltwater fish, especially the consumption of salmon, croaker and mandarin, were significantly correlated with the content of DHA in breast milk. The present study is among the very few in the literature to determine the fatty acid profile of breast milk in Hong Kong populations and verify certain dietary factors that influence this profile. High levels of *n*-3 PUFA, especially DHA, were observed in the breast milk of Hong Kong lactating women. The findings may serve as a dietary reference for lactating mothers to optimise the fatty acid profile of their breast milk.

Key words: DHA: Seafood: Diet: *n*-3 Fatty acids: PUFA: Hong Kong

It is well recognised that breast milk provides infants with the best source of nutrients for healthy growth and development⁽¹⁾. The WHO recommends that infants should be exclusively breastfed for the first 6 months of their lives, and from then on be given appropriate complementary foods along with breastfeeding until the age of 2 years or beyond⁽²⁾. The first 1000 d from conception to age 2 years are a critical period of growth and development during which malnutrition may result in retarded growth and altered metabolism⁽³⁾. However, the rate of exclusive breastfeeding in Hong Kong up to 4 months and up to 6 months was 31 and 1%, respectively, in 2016⁽⁴⁾. These low rates may be caused by a number of factors, including aggressive marketing and promotion of the use of infant formula and inadequate support in workplace and community as well as misconception of the nutritional values of breast milk⁽⁵⁾.

Milk fatty acids, including long-chain PUFA, play an essential role in infant growth, optimal visual and neurological development^(6,7). Humans are unable to synthesise essential fatty acids, linoleic acid and α -linolenic acid. Also, the conversion efficiency of precursor fatty acids to arachidonic acid (AA) and DHA is low⁽⁸⁾. Maintaining an adequate balanced intake of AA and DHA is important for the development and function of the immune system in the early stage of life⁽⁹⁾. Therefore, optimum dietary intakes of both AA and DHA are not only important to breast-fed infants, but are even more important to formula-fed infants who would not receive any long-chain PUFA unless their infant formula is supplemented with them.

The composition of fatty acids in breast milk is regulated by various factors, such as endogenous fatty acids production in the mammary glands and the fatty acids released by maternal adipose tissue⁽¹⁰⁾; it is also significantly influenced by maternal diets

Abbreviations: AA, arachidonic acid; DRI, Dietary Reference Intake.

* **Corresponding authors:** Man-Sau Wong, email man-sau.wong@polyu.edu.hk; Shi-Ying Li, email christine.sy.li@polyu.edu.hk

and nutritional status^(11–13). Seafood, especially oily fish, is a prominent dietary source of *n*-3 fatty acids, particularly DHA. DHA concentration in milk varies from 0.06 to 1.4% among populations⁽¹⁴⁾. Many studies have shown that lactating women living in coastal regions or women who have high fish consumption, like those in Japan⁽¹⁵⁾, Korea⁽¹⁶⁾ and some cities in China^(17,18), have higher *n*-3 fatty acids concentration in their breast milk than those living in inland regions. However, information on the relationship between the milk fatty acid proportions and specific fish consumption is limited. In 2011, the per capita average seafood consumption level in Hong Kong was 3.8 and two times higher than that of the global and China figures, respectively. The city has the second highest per capita seafood consumption in Asia and is the seventh highest in the world⁽¹⁹⁾. Moreover, in recent decades, rapid economic development has led to changes in the lifestyle and dietary habits of the populations⁽²⁰⁾. To our knowledge, although there are some studies on the fatty acid composition of breast milk in Asian and Western countries, the last one conducted on Hong Kong lactating women took place a decade ago⁽²¹⁾. Therefore, the objectives of the present study were (i) to investigate the dietary pattern of Hong Kong lactating women, (ii) to determine the fatty acid profile of their breast milk and (iii) to examine the correlations between the nutritional composition of human milk and dietary characteristics of local mothers.

Methods

Justification of sample size

The optimal sample size was estimated based on a previous published study⁽²²⁾ in which the variability of the level of PUFA in the breast milk of Chinese women (mean 25.5 (SD 5.27) mol% of the total fatty acid content, *n* 82) was considered. The study results were adopted as the subjects of the study are ethnically, geographically and culturally most similar to those of our present study. Cochran's sample size formula was then applied to determine the optimal sample size: $n \geq (ZS/E)^2$ where *n* = sample size, *S* = standard deviation, *Z* = confidence level (95% confidence = 1.96) and *E* = range of possible random error. Assuming that possible random error (*E*) was 4–5% of the mean value, *E* would be equal to 25.5 × 4/5% and the optimal sample size (*N*) was estimated to be $\geq (1.96 \times 5.27/1.02 \sim 1.275)^2 = 65 \sim 102$.

Participants

Subjects were recruited via convenience sampling with the use of posters and email announcements as well as online social networking platforms between August 2014 and June 2015 (Fig. 1). Eligible subjects met the following criteria: they (1) were healthy Chinese lactating mothers aged 18–40 years old; (2) were Hong Kong residents who had resided in Hong Kong for a continuous period of not less than 18 months; (3) had delivered a full term (>37 gestation weeks) singleton who had normal birth weight of >2500 g and no known abnormality; and (4) had infants aged within 2 to 6 months who had been exclusively breastfed or had infants aged over 6 months who had been partially breastfed. Those who were concurrently participating in

any clinical trial or study, taking contraceptive medication after giving birth, suffering from renal, liver or thyroid dysfunction, cognitive impairment or any other indication of a major medical or psychological illness were excluded. A total of 123 volunteers were screened in the present study, among which fifty were excluded. The high exclusion rate (40.7%) of subjects was due to the intake of fatty acid-containing supplements. Seventy-three subjects were recruited for the study. They were divided into three groups according to the lactation stages they were in. Mothers with infants aged 2–6 months who were exclusively breastfed were assigned to Group 1, mothers with infants aged between 7 and 12 months who were partially breastfed were assigned to Group 2 and mothers with infants aged over 12 months who were partially breastfed were assigned to Group 3 (Fig. 1). Researchers were not allowed to influence the diets and extent of breastfeeding of subjects. Subjects were allowed to give their infants complementary foods at their own choice. An informed consent form was signed by each subject after they had received detailed explanation of the study information sheet and protocol. Ethics approval was obtained from the Human Subjects Ethics Sub-Committee and the biological safety and chemical safety of study were approved by the Health, Safety and Environment Office, The Hong Kong Polytechnic University, Hong Kong.

Study design

Data collection

Anthropometric and demographic data collection. Data were collected during interview. Weight and height were measured using standardised methods with the use of a digital electronic balance (REX TE856; Rex Products Ltd) and a measuring tape. They were measured to the nearest 0.1 kg and 0.1 cm, respectively. BMI (kg/m²) was then calculated as body weight (kg) divided by body height (m) squared.

Demographic information such as subject's age, occupation, family income and information on other lifestyle habits such as smoking status and alcohol use was collected using a standardised questionnaire. Information on other maternal characteristics such as the number of pregnancies and births they had had, previous history of lactation, breastfeeding practice and pattern of the current lactation was also collected.

Assessment of dietary intake

Dietary records of three consecutive days prior to the interview were assessed using a 3-d dietary record sheet, which was mailed to each subject after appointment of interview had been made. Prior to the interview, subjects were asked to record the time, types and amounts of both food and beverage, condiments and oils consumed using standard household measures (e.g. bowls and cups). Also, cooking methods were recorded to estimate the amount of cooking oil consumed. Subjects were reminded to consume their usual diets. On the day of the interview, subjects were reminded to bring the records with them to the interview venue. The records were checked by our research staff for any unclear or missing food items. During the interview, a food book/album consisting of photos of local commonly




Fig. 1. Consolidated Standards of Reporting Trials (CONSORT) diagram of the study.

consumed foods and guidelines on estimating food portion sizes and food container volumes was provided to subjects to assist them complete the dietary records. Meanwhile, utensils, including bowls, plates and spoons, of different standardised portion sizes were shown to subjects to help quantify the dietary intake. Moreover, subjects were asked whether they had carried out certain dietary practices such as addition of extra sauces and removal of fat in meats to minimise estimation bias. Subjects were interviewed again over the phone to clarify any incomplete items, if necessary. The records were then analysed using the nutritional analysis software The Food Processor Nutrition Analysis and Fitness software version 10.13.1 (ESHA Research). Traditional Chinese and local foods had been input into the database based on information obtained from China and the Centre for Food Safety, Hong Kong. Maternal intakes of energy, macronutrients and the proportions of several long-chain-PUFA were estimated based on the 3-d dietary records.

To assess their dietary habits during the past 3 months, subjects were asked to complete a FFQ, modified from a previous population survey⁽²³⁾, by addition of PUFA-rich food items in

accordance with the study purpose. The questionnaire comprises eleven food categories with a total of 153 food items, including vegetables and beans, mushrooms and algae, fishes and seafood, fish soups and soup remains, eggs, dairy products and beverages, snacks and soups as well as condiments and oils. The category of fish and seafood includes items on the list of 'Advice for pregnant women, women planning pregnancy and young children on fish consumption', which was issued by the Center for Food Safety, Hong Kong Special Administrative Region, in 2013⁽²⁴⁾. As it is not possible to include all food items in the questionnaire, rows of 'others' were provided to the respondents to record other consumed foods in each part. To minimise assessment error, our subjects were reminded to average the consumption amounts of seasonal foods in the past 3 months. Subjects' consumption frequency, of which nine options ranging from never to everyday in the past 3 months, and the average amounts of specific food items consumed were recorded. The mean daily intake of different kinds of foods was calculated by dividing the total amount by 90 d, whereas that of each food group was calculated by the sum of mean daily intake

of foods in that food group. The mean intake of EPA and DHA from total freshwater fish, saltwater fish and some specific fish species consumed by the lactating women in the present study was then calculated by using the Food Composition Database of the US Department of Agriculture, Agricultural Research Service (2015) and the China Food Composition (Chinese Edition, 2002).

Breast milk collection and analysis

On the day of interview, after completing the dietary record, each subject was asked to supply a specimen of milk that was fully expressed from one breast in a private interview room. Subjects were allowed to use an electric breast pump (mini electric breast pump, Medela Inc.) we provided or their hands to express milk into sterile polystyrene containers without preservatives. The samples were kept in ice buckets and transported to the laboratory within 2 h of collection. All milk samples were divided into aliquots on receipt. One aliquot was stored at -20°C and batched for fatty acid analysis. The remaining aliquots were stored at -80°C until further analysis.

Determination of milk fatty acid profile

The test method for fatty acid determination was modified from AOAC 996.06.

Fat extraction. Fat was extracted from 1 g of milk samples in a Mojonnier flask; 100 mg pyrogallol acid, 2 ml (5 mg/ml C_{11} TAG) internal standard solution and a few boiling granules were then added to the flask; 2 ml ethanol was added and the solution was mixed well; and 4 ml H_2O and 2 ml NH_4OH were added to the mixture. The flask was placed into a basket in a shaking water bath at $70\text{--}80^{\circ}\text{C}$ at a moderate agitation speed for 10 min. The solution was mixed every 5 min. The flask was then cooled down to room temperature. Ethanol was added to fill the bottom reservoir of the flask and mixed gently. Then, 20 ml diethyl ether and 20 ml petroleum ether were added to the flask and then the mixture was shaken gently. The ether (top) layer was decanted into a 150 ml boiling tube. Addition of diethyl ether and petroleum ether was repeated twice. Ether was evaporated with the aid of a nitrogen blow concentrator (Eyla MGS2200).

Methylation. Fat was then dissolved in chloroform and diethyl ether. The mixture was transferred to a test tube and then evaporated to dryness using a nitrogen blow concentrator. Fatty acid methyl esters (FAME) were obtained after trans-methylation of total lipids by using 7% BF_3 in toluene (1 ml) for 45 min at 100°C . FAME were extracted with hexane (1 ml).

GC determination. With added internal standard (FAME C_{11} : 0), samples were injected into a gas chromatograph (GC system, 7890A; Agilent Technologies), separated by a capillary column (Supleco, SP 2560) and analysed by a flame ionised detector. Commercially available FAME standard (Nu-Check- Prep, catalogue no. GLC- Nestle-36 (Nu-Check-Prep GLC-Nestle36, or equivalent grade)) was injected to the gas chromatograph to establish a calibration curve for peak identification on the basis of retention times. Specific fatty acid was then quantified by

calculation from the calibration curve with reference to an internal standard.

Statistical analyses

All analyses were performed using the software SPSS version 23. Normality of data was tested by Shapiro–Wilk test while homogeneity of variances was assessed by Levene's test. Normal distributed data are shown as means and standard deviations, whereas non-normally distributed data are shown as medians and interquartile ranges. Correlation of maternal intake of $n\text{-}3$ fatty acids as well as freshwater and saltwater fish species and fatty acid concentration in breast milk was determined by the Spearman rank correlation test and stepwise multiple linear regressions. $P < 0.05$ (two-tailed) was considered statistically significant for all analyses.

Results

Subjects' characteristics

The anthropometric and socio-economic characteristics of seventy-three subjects, namely their age, body weight, body height, BMI, education level, household income and occupation, are shown in Table 1. The subjects were young women with a

Table 1. Anthropometric and socio-economic characteristics of seventy-three lactating women (Mean values and standard deviations; numbers of participants and percentages)

Variable	<i>n</i>	%
Age (years)		
Mean	32.1	
SD	3.7	
Body weight (kg)		
Mean	56.9	
SD	8.5	
Body height (m)		
Mean	1.60	
SD	0.05	
BMI* (kg/m ²)		
Mean	22.2	
SD	3.1	
Underweight	3	4.1
Normal	46	63.0
Overweight	13	17.8
Obese	11	15.1
Maternal education level†		
Low	17	23.3
High	56	76.7
Monthly household income‡		
Low	23	31.5
High	50	68.5
Occupation		
Housewife	16	21.9
Working full time	57	78.1
Duration of lactation (months)		
<0–6	29	39.7
>6–12	20	27.4
>12–24	24	32.9

* Underweight: BMI < 18.5 kg/m²; normal: BMI $18.5\text{--}23.0$ kg/m²; overweight: BMI $23\text{--}25.0$ kg/m²; obese: BMI ≥ 25 kg/m².

† High: tertiary education or above.

‡ High: monthly household income $>$ HK\$ 30 000.

Table 2. Dietary intake of macronutrients of the seventy-three lactating women* (Mean values and standard deviations; medians and interquartile ranges (IQR))

Energy and nutrients	Dietary intake		Recommendation from Chinese DRI 2013
	Mean	SD	
Energy (kcal)†	2370.2	407.1	2300
Carbohydrates (g)	258.6	49.5	
% Energy	44.0	6.6	50–65 (AMDR)
% subjects < AMDR		80.0	
% subjects within range		20.0	
% subjects > AMDR		0.0	
Protein (g)	112.1	27.2	80 g (RNI)
% Energy	18.9	3.1	
% subjects <AMDR		0.0	
% subjects within range		10.0	
% subjects >AMDR		90.0	
Fat (g)	98.1	24.4	
% Energy	36.9	5.0	20–30 (AMDR)
% subjects <AMDR		0.0	
% subjects within range		10.0	
% subjects >AMDR		90.0	
SFA (g)	24.3	7.5	
% Energy	9.1	2.0	<10 (AMDR)
MUFA (g)	30.9	10.5	
% Energy	11.6	2.9	
PUFA (g)			
Median	17.7		
IQR	9.0		
% Energy	7.1	1.6	
LA (g)	10.8	4.1	4 (RNI)
% Energy	4.1	1.3	
ALA (g)	1.4	1.4	0.60 (AI)
% Energy	0.54	0.59	
AA (g)			
Median	0.26		
IQR	0.19		
EPA			
Median	0.08		
IQR	0.12		
DHA			0.2 (AI)
Median	0.20		
IQR	0.29		
EPA+DHA			0.25 (AI)
Median	0.28		
IQR	0.42		

DRI, Dietary Reference Intake; AMDR, acceptable macronutrient distribution range; RNI, recommended nutrient intake; LA, linoleic acid; AI, adequate intake; ALA, α -linolenic acid; AA, arachidonic acid.

* Data are calculated from the 3-d dietary records.

† To convert energy in kcal to kJ, multiply by 4.184.

mean age of about 32 years and a mean BMI around 22 kg/m². Nearly one-third of the subjects were overweight (BMI \geq 23 kg/m²) or obese (BMI \geq 25 kg/m²)⁽²⁵⁾. The majority were working full-time. Two-thirds of them had both tertiary or above education level and high monthly household income (>HK\$ 30 000)⁽²⁶⁾. Positive correlation was shown between the maternal education level and household income. Subjects with infants aged from 2 to 6 months were exclusively breastfeeding (*n* 29).

Dietary intake of lactating mothers

The mean dietary intakes of macronutrients and selected kinds of PUFA of lactating women, as calculated from the 3-d dietary records, are shown in Table 2. The energy intake of the lactating

Table 3. Food group consumption of the seventy-three lactating women* (Medians and interquartile ranges (IQR); mean values and standard deviations)

Food groups	Intake (g/d)		Servings/d†		Suggested servings/d from Hong Kong ⁽⁴²⁾
	Median	IQR	Median	IQR	
Grains	510.3	238.8	6.8	3.2	4–5
Vegetables	119.3	93.0	1.5	1.2	4–5
Nuts‡	0.0	6.7			
Fruits	68.3	120.6	0.9	1.5	3
Meats					6–7
Mean	268.9		9.0		
SD	109.7		3.7		
Red meats	91.2	99.6			
Poultry	51.7	74.8			
Seafood‡	87.5	92.0			
Eggs	25.8	34.4			
Dairy products	41.7	161.6	0.2	0.7	2

* Data are calculated from the 3-d dietary records.

† One serving: grains: 75 g; vegetables: 80 g; fruits: 80 g; dairy products: 240 ml; meats: 30 g (cooked).

‡ Food items rich in *n*-3 fatty acids.

Table 4. Dietary intake of EPA and DHA from freshwater fish, saltwater fish and some fish species consumed in Hong Kong by seventy-three lactating women* (Mean values and standard deviations)

	Daily intake of EPA and DHA from fish (mg/d)			
	EPA		DHA	
	Mean	SD	Mean	SD
Fresh water fish	22.59	32.79	30.89	43.11
Grass fish	0.67	0.90	2.00	2.70
Mandarin fish	9.22	22.39	11.00	26.73
Largemouth bass	1.33	6.16	1.59	7.35
Mud carp	0.65	2.03	1.68	5.30
Bid head fish	2.97	12.80	3.55	15.28
Saltwater fish (g)	1.23	2.25	2.53	3.95
Salmon	88.82	91.02	139.62	143.08
Pomfret	3.37	7.59	2.07	4.67
Croaker	5.82	11.84	15.17	30.83
Golden thread	3.53	8.24	8.69	20.27
Grouper	0.97	2.23	2.56	5.58

* Data are estimated from the FFQ.

mothers was about 2400 kcal/d (10 040 kJ/d) while the percentage energy intake of mothers from carbohydrates, protein and fat was about 44, 18 and 37 %, respectively. The subjects' mean dietary intake of the macronutrients was out of the acceptable macronutrient distribution range (AMDR) suggested by the Chinese Dietary Reference Intakes (DRI) 2013⁽²⁷⁾. The diets of these lactating women were characterised by high consumption of protein and fat. Maternal mean protein intake was higher than the Chinese recommended nutrient intake (RNI) for lactating women by 40 %. The mean maternal energy intake from fat was higher than the upper limit of the Chinese AMDR by 23 %. On the other hand, maternal intakes of several *n*-3 fatty acids (i.e. α -linolenic acid, EPA and DHA) as well as *n*-6 fatty acids (i.e. linoleic acid) met the levels recommended by the Chinese DRI.

The mean intake of different food groups of the lactating mothers, as calculated from the 3-d dietary records, is shown

Table 5. Milk fatty acid (FA) composition of the seventy-three lactating women at different stages of lactation* (Mean values and standard deviations)

Fatty acids (%wt/wt of all FA)	All subjects (n 73)		Group 1 (n 29)		Group 2 (n 20)		Group 3 (n 24)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
SFA								
10:0	1.18	0.40	1.29	0.41	1.12	0.46	1.10	0.31
12:0	6.02	1.78	5.08	1.80 ^b	6.43	1.65 ^a	6.76	1.48 ^a
14:0	8.03	6.14	8.44	9.48	7.21	1.84	8.05	2.45
15:0	0.19	0.11	0.18	0.08	0.21	0.16	0.20	0.08
16:0	19.0	2.43	18.32	3.04	19.21	1.65	19.51	1.77
17:0	0.25	0.06	0.26	0.07	0.25	0.05	0.25	0.07
18:0	5.89	1.13	5.61	1.12	6.13	0.92	6.03	1.23
20:0	0.05	0.09	0.12 ^a	0.09	0.03 ^b	0.08	0.01 ^b	0.03
21:0	0.40	0.33	0.20	0.22 ^b	0.50	0.30 ^a	0.55	0.36 ^a
22:0	0.29	0.24	0.19	0.19 ^b	0.35	0.19 ^a	0.36	0.28 ^a
24:0	0.03	0.06	0.07	0.09	ND		ND	
Total SFA	41.43	6.07	39.86	7.77	41.50	4.46	42.91	4.44
MUFA								
14:1 <i>n</i> -5	0.05	0.08	0.09	0.08 ^a	0.02	0.07 ^b	0.03	0.06 ^b
15:1	ND		ND		ND		0.01	0.07
16:1 <i>n</i> -7	2.48	0.74	2.67	0.75	2.27	0.87	2.47	0.54
17:1 <i>n</i> -7	0.13	0.10	0.05	0.09 ^b	0.18	0.07 ^a	0.19	0.05 ^a
18:1 <i>n</i> -7	33.52	3.64	33.60	4.09	34.17	3.20	33.12	3.69
20:1 <i>n</i> -9	0.15	0.27	0.38	0.32	ND		ND	
22:1 <i>n</i> -9	0.02	0.05	0.06	0.06	ND		ND	
24:1 <i>n</i> -9	0.07	0.11	0.05	0.08	0.07	0.11	0.07	0.10
Total MUFA	36.42	3.66	36.91	4.35	36.71	3.11	35.88	3.50
<i>n</i>-6 PUFA								
18:2 <i>n</i> -6 (LA)	17.16	3.50	17.89	4.09	16.83	3.63	16.72	2.45
18:3 <i>n</i> -6	1.07	0.77	1.66	0.87 ^a	0.72	0.39 ^b	0.65	0.30 ^b
20:2 <i>n</i> -6	0.08	0.15	0.21	0.17	ND		ND	
20:3 <i>n</i> -6	0.09	0.16	0.23	0.19	ND		ND	
20:4 <i>n</i> -6 (AA)	0.71	0.22	0.74	0.24	0.66	0.18	0.67	0.21
22:2 <i>n</i> -6	0.02	0.07	0.06	0.11	ND		ND	
Total <i>n</i> -6 PUFA	18.65	3.60	19.44	4.29	18.21	3.57	18.05	2.49
<i>n</i>-3 PUFA								
18:3 <i>n</i> -3 (ALA)	1.49	1.12	2.20	1.02	2.01	0.66	1.79	0.35
20:3 <i>n</i> -3	0.01	0.03	0.03	0.04	ND		ND	
20:5 <i>n</i> -3 (EPA)	0.29	0.29	0.36	0.31	0.26	0.28	0.23	0.22
22:5 <i>n</i> -3	0.41	0.56	0.34	0.18	0.40	0.27	0.53	0.93
22:6 <i>n</i> -3 (DHA)	0.80	0.65	0.86	0.66	0.91	0.88	0.64	0.35
Total <i>n</i> -3 PUFA	3.54	1.41	3.79	1.54	3.58	1.38	3.20	1.26
Total PUFA	22.14	4.27	23.23	4.99	21.79	4.01	21.25	3.06
<i>n</i>-6/<i>n</i>-3 PUFA	5.84	1.87	5.68	1.90	5.66	1.98	6.19	1.75
18:2 <i>n</i> -6 / 18:3 <i>n</i> -3	9.12	2.35	8.95	2.73	8.84	2.39	9.55	1.81
20:5 <i>n</i> -3 + 22:6 <i>n</i> -3	1.09	0.83	1.22	0.86	1.17	1.04	0.87	0.52
Total fat (g/l)	41.34	13.67	39.13	14.69 ^{a,b}	36.76	9.76 ^b	47.84	13.22 ^a

ND, not detectable; LA, linoleic acid; AA, arachidonic acid; ALA, α -linolenic acid.

^{a,b} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

in Table 3. Intakes of grains and meats could meet the recommended levels. However, intakes of vegetables, fruits and dairy products were all below the recommended levels.

The mean dietary intake of EPA and DHA from the freshwater fish, saltwater fish and some fish species consumed in Hong Kong of the seventy-three lactating women, as estimated from the FFQ, is shown in Table 4. The maternal dietary intake of EPA and DHA from salmon ranked the highest among all types of fish species.

Breast milk fatty acid composition of lactating mothers

The fatty acid composition in the breast milk of the seventy-three lactating women is shown in Table 5. The three categories of fatty acids, SFA, MUFA, PUFA, represented approximately 42, 36 and 22%, respectively, of total fatty acids. Palmitic acid accounted for the largest proportion of SFA while oleic acid

and linoleic acid accounted for the largest proportion of MUFA and PUFA, respectively.

The fatty acid composition of the breast milk of mothers in the three groups is also shown in Table 5. Significant differences in concentrations of some fatty acids were mainly observed between mothers from Group 1 and Group 2 as well as between mothers from Group 1 and Group 3. Mean DHA level was approximately 0.9% of total fatty acids in the milk from mothers who had exclusively breastfed their infants aged within 6 months.

Correlation between maternal dietary intake and fatty acid composition of breast milk

Correlation tests were conducted between the dietary intake of *n*-3 fatty acids of the seventy-three lactating women and their levels in breast milk (Fig. 2). Results showed that levels of




Fig. 2. Scatter plots of the intakes of *n*-3 fatty acids (FA) of the seventy-three lactating women and their contents in breast milk (Spearman's correlation coefficient (r_s) between each pair of factors is shown in the plot). ALA, α -linolenic acid.

Table 6. Spearman's correlation coefficients between dietary intake of freshwater and saltwater fish species consumed in Hong Kong and contents of DHA and EPA in breast milk

Dietary intake (g/d)	Breast milk fatty acids (%wt/wt of all fatty acids)	
	EPA	DHA
Freshwater fish	0.158	0.285**
EPA from freshwater fish	0.212	0.312**
DHA from freshwater fish	0.185	0.296**
Grass fish	0.093	0.194
Mandarin fish	0.250*	0.268*
Largemouth bass	-0.123	-0.007
Mud carp	0.039	0.105
Bid head fish	0.011	0.023
Saltwater fish	0.284**	0.404**
DHA from saltwater fish	0.024	0.121
EPA from saltwater fish	0.048	0.128
Salmon	0.165	0.214*
Pomfret	0.004	0.167
Croaker	0.177	0.241**
Golden thread	0.098	0.161
Grouper	-0.029	0.093

Significant correlation: * $P < 0.05$, ** $P < 0.01$ (two-tailed).

maternal DHA ($R = 0.290$, $P = 0.008$) and α -linolenic acid ($R = 0.239$, $P = 0.023$) intake and their levels in milk were significantly associated. Moreover, the results revealed significant associations between DHA content in the milk samples and maternal intakes of both freshwater and saltwater fish as well as some frequently consumed types of fish in Hong Kong (Table 6). In stepwise linear regression analysis, we studied the association of milk DHA concentration with maternal intake of freshwater fish, saltwater fish, age, BMI, education level and monthly household income, respectively. Maternal intake of saltwater fish was the only significant predictor of the content of DHA in milk ($R^2 = 0.242$, $P < 0.001$) while the other variables were not.

Discussion

Hong Kong people have a local dietary habit with high seafood consumption. Over the past decades, the dietary patterns of Hong Kong people have had substantial changes, which may have significant impacts on the fatty acid components of the breast milk of Hong Kong mothers⁽²⁸⁾. The last study on the

breast milk fatty acid components in Hong Kong dates back to 1997⁽²¹⁾. Previous studies have reported different breast milk fatty acid profiles among mainland Chinese women living in urban, rural, coastal and inland regions and showed maternal diet impacts on the fatty acids in milk^(18,22,29,30). The present study shows the dietary patterns of Hong Kong lactating mothers and confirms the association between fatty acid contents in their breast milk and their consumption of aquatic foods. More importantly, for the first time, the study demonstrates the associations between the fatty acid contents in milk of mothers and their specific fish consumptions.

The present study shows that the concentrations of DHA (0.86%) and *n*-3 fatty acids (3.79%) in breast milk are significantly higher than those observed in Hong Kong in 1997 (0.56, 2.17%)⁽²¹⁾ and those from coastal areas of China (0.41–0.61%, 1.68–3.34%)^(18,22,29). Moreover, the contents of DHA (0.86%) and AA (0.74%) in breast milk of Hong Kong lactating mothers are higher than their worldwide mean (0.32, 0.47%)⁽³¹⁾.

The high content of DHA (0.86%) in breast milk of Hong Kong mothers is comparable with that from some island countries and regions such as Japan (0.99%)⁽¹⁵⁾, Korea (0.96%) and Taiwan (0.98, 1.13%), regions that account for the high levels of DHA concentrations in the breast milk of Asian mothers⁽³²⁾. DHA is synthesised limitedly in infants from the age of 0 to 6 months, regarded as a conditional essential fatty acid for infants⁽³³⁾. Assuming that the daily milk consumption of milk is 750 ml, among the twenty-nine milk samples from lactating mothers with infants aged from 2 to 6 months old, twenty-four could provide their infants (i.e. more than 80%) with adequate level of DHA (100 mg) as recommended by the Chinese DRI 2013⁽²⁷⁾.

The average fish intake of lactating mothers in our study is ten servings per week, which is higher than the recommended weekly consumption suggested by the US Department of Agriculture⁽³⁴⁾. There is increasing amount of evidence suggesting that maternal diet affects milk lipid profile to a certain extent⁽³⁵⁾. It is well known that both seafood and fish are prominent dietary sources of *n*-3 fatty acids, especially DHA and EPA⁽³⁶⁾. The findings are in line with those of other studies, suggesting a positive association between DHA concentration in milk of lactating mothers and their frequency of fish consumption⁽¹⁴⁾. However, our results only show significant association between milk EPA and intake of saltwater fish, but not freshwater fish. The most frequently consumed type of fish by the subjects in our study is salmon, which is regarded as a type of fatty fish containing high levels of EPA and DHA⁽³⁷⁾. Our subjects' high consumption of *n*-3 fatty acids-rich fish may explain the result that there is correlation only between the intake of saltwater fish and the content of EPA in the milk samples. The DHA content in the breast milk from Hong Kong mothers (0.86%) is higher than that from mainland Chinese mothers (0.38%), while the EPA content in the breast milk from Hong Kong mothers (0.36%) is comparable with that from mainland Chinese mothers (0.41%)⁽³⁰⁾. Our study shows that DHA content in breast milk is more influenced by maternal saltwater fish intake (r 0.404) than EPA content (r 0.284). This may explain the significant differences in DHA contents in breast milk between Hong Kong mothers and mainland Chinese mothers. However, correlation analysis indicates that

the intakes of EPA and DHA from total consumption of saltwater fish are not associated with the milk concentrations of the corresponding fatty acid. Thus, the results are contradictory to the findings on the correlation between intake of saltwater fish and the DHA and EPA content of breast milk. Such discrepancy may be due to the use of both Chinese and US databases as references in the present study to estimate the dietary intake of EPA and DHA from different fish. The level of fatty acids in fish varies with many factors such as growing location and fish feeds, which may result in the inconsistency in findings of the correlation test. As expected, PUFA content in breast milk apparently varies with both the quantities and types of fish consumed. Populations in Hong Kong consume a wide range of fish, among which maternal intakes of salmon, croaker and mandarin fish are significantly correlated with milk DHA content, while maternal intake of mandarin fish is significantly correlated with milk EPA content (Table 6).

The AA concentration of 0.74% of our milk samples is within the recommended range of 0.35–1.00%⁽³⁸⁾ and slightly lower than the DHA concentration (0.86%). The results of the present study reveal that, in line with the results of some previous studies showing that the content of AA in breast milk is relatively stable and not influenced by maternal dietary intake⁽³⁹⁾, the correlation between maternal intake of AA and milk AA is insignificant. The concentration of AA in breast milk is commonly higher than that of DHA; it is less dependent on dietary intake, whereas that of DHA fluctuates more due to changes in dietary intake. Interestingly, AA levels in breast milk also tend to be higher when DHA levels are higher, suggesting the importance of maintaining a certain ratio of AA and DHA. However, we do not know yet the right balance between these two; but recent systematic reviews have pointed toward the beneficial effect of having more DHA in breast milk on some health outcomes such as the cognitive and visual development in preterm infants^(9,40).

The present cross-sectional study examines the dietary patterns and fatty acid profile of breast milk in Hong Kong mothers at different stages of lactation (0–24 months). It is likely that there are factors other than diet that affect the breast milk composition. Milk fatty acid contents may change along the different stages of lactation⁽³⁵⁾. Many women in Hong Kong follow traditional Chinese cultural practices during the first month postpartum, with restrictions on their diet, hygiene and physical activities⁽⁴¹⁾, but the impacts of these practices on breast milk composition are not well studied.

There are some limitations of the present study. The recruitment procedure, including promotion on campus via poster and email or via online social platforms and the voluntary participation of the women, may have resulted in selection bias, as women who use online social platforms are expected to have high educational level and economic status. Also, the generalisability of the present results is limited by the small sample size. The preliminary results of the present study are able to give us more information about the milk fatty acid profile of Hong Kong lactating women, and thus support us to design further studies with larger sample sizes.

In conclusion, the present study determines the fatty acid profile of breast milk in Hong Kong populations and verifies that certain dietary factors could influence the fatty acid profile. High levels of *n*-3 PUFA, especially DHA, have been observed in the breast milk of Hong Kong lactating women. Specifically, maternal intakes of certain fish types are significantly correlated with

milk DHA content. This may serve as a dietary reference for lactating mothers to optimise their breast milk fatty acid profile.

Acknowledgements

The authors wish to acknowledge the support of Laboratory for Infant & Child Nutrition, Food Safety and Technology Research Centre, Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University and the mothers who donated their milk for our study.

The present study was funded by the Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University.

V. W.-S. W., M.-S. W., S.-Y. L., S.-M. C., Y.-X. S., H.-M. C. and K. W.-H. K. designed the study, analysed and interpreted the data and drafted the manuscript. H.-W. L. and W.-Y. P. performed experiments. C.-L. C. and Y.-F. N. designed the study.

The authors declared that they have no conflict of interest.

References

- Agostoni C, Braegger C, Decsi T, *et al.* (2009) Breast-feeding: a commentary by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr* **49**, 112–125.
- Gersovitz M, Madden JP, Smiciklas-Wright H (1978) Validity of the 24-hr. dietary recall and seven-day record for group comparisons. *J Am Diet Assoc* **73**, 48–55.
- Barker DJ, Eriksson JG, Forsen T, *et al.* (2002) Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol* **31**, 1235–1239.
- Moreira AS, Teixeira M, da Silveira Osso F, *et al.* (2009) Left ventricular hypertrophy induced by overnutrition early in life. *Nutr Metab Cardiovasc Dis* **19**, 805–810.
- Tarrant M, Fong DY, Wu KM, *et al.* (2010) Breastfeeding and weaning practices among Hong Kong mothers: a prospective study. *BMC Pregnancy Childbirth* **10**, 27.
- Lauritzen L, Hansen HS, Jorgensen MH, *et al.* (2001) The essentiality of long chain *n*-3 fatty acids in relation to development and function of the brain and retina. *Prog Lipid Res* **40**, 1–94.
- Tau GZ, Peterson BS (2010) Normal development of brain circuits. *Neuropsychopharmacol* **35**, 147–168.
- Gibson RA, Muhlhausler B, Makrides M (2011) Conversion of linoleic acid and α -linolenic acid to long-chain polyunsaturated fatty acids (LCPUFAs), with a focus on pregnancy, lactation and the first 2 years of life. *Matern Child Nutr* **7**, 17–26.
- Lien EL, Richard C, Hoffman DR (2018) DHA and ARA addition to infant formula: current status and future research directions. *Prostaglandins Leukot Essent Fatty Acids* **128**, 26–40.
- Jensen RG (1996) The lipids in human milk. *Prog Lipid Res* **35**, 53–92.
- Makrides M, Simmer K, Neumann M, *et al.* (1995) Changes in the polyunsaturated fatty acids of breast milk from mothers of full-term infants over 30 wk of lactation. *Am J Clin Nutr* **61**, 1231–1233.
- Neville MC, Picciano MF (1997) Regulation of milk lipid secretion and composition. *Annu Rev Nutr* **17**, 159–183.
- Bokor S, Koletzko B, Decsi T (2007) Systematic review of fatty acid composition of human milk from mothers of preterm compared to full-term infants. *Ann Nutr Metab* **51**, 550–556.
- Antonakou A, Skenderi KP, Chiou A, *et al.* (2013) Breast milk fat concentration and fatty acid pattern during the first six months in exclusively breastfeeding Greek women. *Eur J Nutr* **52**, 963–973.
- Yuhans R, Pramuk K, Lien EL (2006) Human milk fatty acid composition from nine countries varies most in DHA. *Lipids* **41**, 851–858.
- Golfetto I, McGready R, Ghebremeskel K, *et al.* (2007) Fatty acid composition of milk of refugee Karen and urban Korean mothers. Is the level of DHA in breast milk of Western women compromised by high intake of saturated fat and linoleic acid? *Nutr Health* **18**, 319–332.
- Gao Y, Zhang J, Wang C, *et al.* (2011) [Fatty acid composition of mature human milk in three regions of China]. *Wei Sheng Yan Jiu (J Hygiene Res)* **40**, 731–734.
- Urwin HJ, Zhang J, Gao Y, *et al.* (2013) Immune factors and fatty acid composition in human milk from river/lake, coastal and inland regions of China. *Br J Nutr* **109**, 1949–1961.
- Food and Agriculture Organization of the United Nations (2014) *The State of World Fisheries and Aquaculture 2014 (SOFIA)*. Rome, Italy: FAO Fisheries and Aquaculture Department.
- Department of Health (2013) Diet, physical activity and health: Hong Kong situation. http://www.change4health.gov.hk/filemanager/common/image/strategic_framework/action_plan/action_plan_2_e.pdf (accessed May 2016).
- Chen ZY, Kwan KY, Tong KK, *et al.* (1997) Breast milk fatty acid composition: a comparative study between Hong Kong and Chongqing Chinese. *Lipids* **32**, 1061–1067.
- Peng Y, Zhou T, Wang Q, *et al.* (2009) Fatty acid composition of diet, cord blood and breast milk in Chinese mothers with different dietary habits. *Prostaglandins Leukot Essent Fatty Acids* **81**, 325–330.
- Woo J, Leung SSF, Ho SC, *et al.* (1997) A food frequency questionnaire for use in the Chinese population in Hong Kong: description and examination of validity. *Nutr Res* **17**, 1633–1641.
- Centre for Food Safety (2003) *Advice for Pregnant Women, Women Planning Pregnancy and Young Children on Fish Consumption*. Hong Kong, China: Department of Food & Environmental Hygiene.
- Arenz S, Ruckerl R, Koletzko B, *et al.* (2004) Breast-feeding and childhood obesity—a systematic review. *Int J Obes Relat Metab Disord* **28**, 1247–1256.
- Chan MF, Mok E, Wong TK, *et al.* (2008) Investigating the health profiles of Hong Kong Chinese: results of a cluster analysis. *J Clin Nurs* **17**, 911–920.
- Chinese Nutrition Society (2013) *Chinese Dietary Reference Intakes*. China: Science Press.
- Zhang J, Dhakal IB, Zhao Z, *et al.* (2012) Trends in mortality from cancers of the breast, colon, prostate, esophagus, and stomach in East Asia: role of nutrition transition. *Eur J Cancer Prev* **21**, 480–489.
- Li J, Fan Y, Zhang Z, *et al.* (2009) Evaluating the trans fatty acid, CLA, PUFA and erucic acid diversity in human milk from five regions in China. *Lipids* **44**, 257–271.
- Jiang J, Wu K, Yu Z, *et al.* (2016) Changes in fatty acid composition of human milk over lactation stages and relationship with dietary intake in Chinese women. *Food Funct* **7**, 3154–3162.
- Brenna JT, Varamini B, Jensen RG, *et al.* (2007) Docosahexaenoic and arachidonic acid concentrations in human breast milk worldwide. *Am J Clin Nutr* **85**, 1457–1464.
- Li SY, Dong XL, Wong WSV, *et al.* (2015) Long-chain polyunsaturated fatty acid concentrations in breast milk from Chinese mothers: comparison with other regions. *Int J Child Health Nutr* **4**, 230–239.
- Food and Agriculture Organization of the United Nations (2010) *Fats and Fatty Acids in Human Nutrition: Report of a Joint FAO/WHO Expert Consultation*. Rome: FAO.
- US Food & Drug Administration (2017) Eating fish: what pregnant women and parents should know. <https://www.fda.gov/Food/ResourcesForYou/Consumers/ucm393070.htm> (accessed April 2018).

35. Koletzko B (2016) Human milk lipids. *Ann Nutr Metab* **69**, 27–40.
36. Grote V, Verduci E, Scaglioni S, *et al.* (2016) Breast milk composition and infant nutrient intakes during the first 12 months of life. *Eur J Clin Nutr* **70**, 250–256.
37. Philibert A, Vanier C, Abdelouahab N, *et al.* (2006) Fish intake and serum fatty acid profiles from freshwater fish. *Am J Clin Nutr* **84**, 1299–1307.
38. Antonakou A, Skenderi KP, Chiou A, *et al.* (2013) Breast milk fat concentration and fatty acid pattern during the first six months in exclusively breastfeeding Greek women. *Eur J Nutr* **52**, 963–973.
39. Demmelmair H, Baumheuer M, Koletzko B, *et al.* (2001) Investigation of long-chain polyunsaturated fatty acid metabolism in lactating women by means of stable isotope techniques. *Adv Exp Med Biol* **501**, 169–177.
40. Koletzko B, Boey CC, Campoy C, *et al.* (2014) Current information and Asian perspectives on long-chain polyunsaturated fatty acids in pregnancy, lactation, and infancy: systematic review and practice recommendations from an early nutrition academy workshop. *Ann Nutr Metab* **65**, 49–80.
41. Liu N, Mao L, Sun X, *et al.* (2006) Postpartum practices of puerperal women and their influencing factors in three regions of Hubei, China. *BMC Public Health* **6**, 274.
42. Family Health Service (2016) Healthy eating during pregnancy and breastfeeding. https://www.fhs.gov.hk/english/health_info/woman/20036.html (accessed June 2016)