

Review: Pork production with maximal nitrogen efficiency

S. Millet^{†,a}, M. Aluwé, A. Van den Broeke, F. Leen, J. De Boever and S. De Campeneere

ILVO (Flanders Research Institute for Agriculture, Fisheries and Food), Scheldeweg 68, 9090 Melle, Belgium

(Received 5 March 2017; Accepted 11 September 2017; First published online 25 October 2017)

During growth, pigs convert plant protein into animal protein. The major part of the ingested protein is excreted via manure, with potential nitrogen (N) losses to the environment. To limit N losses and increase sustainability of pork production, the efficiency of protein conversion should be maximized. The aim of this paper is to critically evaluate diet and management strategies linked with N efficiency. Besides nutrition, we discuss three management strategies observed in science and in practice to be linked with improved N efficiency: genetic selection, castration and slaughter weight. Because diet has a marked effect on eventual N losses, it must also be taken into account when evaluating management strategies. A reductionist approach, such as feeding the same diet across all management treatments, may overestimate the effect of a management strategy and eventually lead to incorrect conclusions. The amount of excreted N depends on the amount of ingested N, the amount of absorbed N, the amino acid (AA) balance in the diet and the animal's N and AA requirements. Daily multiphase feeding adapted to the individual animal's AA needs is likely to be the most N efficient. For animals housed in groups, phase feeding is necessary. When combined with periods of temporary AA restriction, N efficiency can be further improved. Specific AA consumption must be balanced by applying the ideal protein concept. With better knowledge of the requirements of individual animals and the commercial availability of certain AAs, the total dietary CP level can be lowered within limits. Further research is needed on the minimal CP level that allows maximal performance. For this end a useful parameter may be the ratio of standardized ileal digestible (SID) lysine : apparent total tract digestible CP level. By combining optimal nutrition and management, a whole body N efficiency approaching 60% may be achievable in the near future.

Keywords: amino acid, diet, growing-finishing pig, management strategies, nitrogen efficiency

Implications

This study reviews several management strategies that affect the amount of nitrogen (N) used per kg of meat produced. Genetic selection, castration and slaughter weight have an effect. Diet is still the most important factor affecting N efficiency, however. When the diet is adapted to the animal requirements, a whole body N efficiency close to 60% seems attainable for group-fed pigs. This would improve the environmental sustainability of pork production. When evaluating the effect of management strategies on N efficiency, all of these concepts need to be taken into account to avoid false conclusions.

Introduction

In contemporary pig production, ~6.3 kg N is used to raise an 8-kg piglet to a 110-kg finishing pig (own calculations based on current Belgian feeds and breeds, Table 1).

While ~46% of this N is retained in the animal, the other 54% is excreted, mainly through feces and urine. Part of the excreted N can be re-used as a fertilizer but part of it is either lost to the air as ammonia (NH₃), nitrous oxide (N₂O) and N oxide (NO_x) emissions or lost to ground and surface waters via leaching and runoff of nitrate (NO₃) and other N compounds (Leip *et al.*, 2014). Increasing the efficiency of converting plant protein into animal protein would decrease the environmental burden per kg pork and may improve the sustainability of pig production.

Nitrogen efficiency decreases ~3% (from 46% to 43%) when the sow's production of piglets is included in the calculation (Table 1). In piglet production, N is retained in piglets (200 g N per 8 kg piglet, Table 1) and in the sow (BW gain). Most of the N is excreted to the urine and feces, yielding a N efficiency level around 24%. Relative to the fattening phase, the effect of the piglet production on N efficiency is small. For a sow producing 27 piglets per year, the sow and her piglets consume around 28.8 kg N per year, while the 27 piglets consume around 170.1 kg N to grow from 8 to 110 kg (Table 1). This may explain why efforts to

^a The content of this manuscript was presented at the 5th EAAP International Symposium on Energy and Protein Metabolism and Nutrition, Krakow, Poland.

[†] E-mail: sam.millet@ilvo.vlaanderen.be

Table 1 Estimation of nitrogen input and nitrogen efficiency of a hybrid sow and her offspring (27 piglets per year over 2.4 cycles) in a commercial Belgian farm (Hybrid sow × Piétrain boar)

Parameters	Estimate	Reference or calculation
Sow and piglet phase (until weaning)		
Days per phase		
Lactation (a)	67	2.4 cycles × 28 days
Gestation + non-productive days	298	365-a
Daily feed intake (kg)		
Lactation	6.9	CVB (2016)
Gestation	2.7	
Creep feed intake per pig (kg) (b)	0.275	Estimation ¹
Dietary CP content (g/kg)		Estimation ¹
Lactation	160	
Gestation	130	
Creep feed	170	
Nitrogen intake of the sow (kg) (c)	28.6	$= \sum_1^i \frac{(\text{Dietary CP}_i \times \text{number of days}_i \times \text{Daily feed intake}_i)}{6.25}$ with <i>i</i> the phase in the reproductive cycle
Nitrogen intake of the piglets (kg) (d)	0.2	$\frac{27 \text{ piglets} \times b \times \text{CP content creep feed}}{6.25}$
Nitrogen content of 8 kg piglet (kg) (e)	0.2	Warnants <i>et al.</i> (2006)
BW gain of the sow, (kg) (f)	48.0	2.4 cycles × 20 kg (CVB, 2016)
CP content of sows' BW gain (g/kg) (g)	178.0	Van den Broeke <i>et al.</i> (2017) ²
Nitrogen retained in the sow (kg) (h)	1.4	$\frac{(f \times g)}{6.25}$
Nitrogen retained in the piglets (kg) (i)	5.4	27 × e
Nitrogen efficiency (%)	24	$\frac{(h+i)}{(c+d)} \times 100$
Piglets from weaning to slaughter		
Nitrogen content of 110 kg pig with 62% meat (kg) (j)	3.1	Millet <i>et al.</i> (2010)
Dietary CP content (g/kg)		Estimation ¹
8 to 25 kg	174	
25 to 45 kg	159	
45 to 70 kg	146	
70 to 110 kg	138	
Feed conversion ratio (kg/kg)		Estimation ¹
8 to 25 kg	1.7	
25 to 45 kg	2.1	
45 to 70 kg	2.6	
70 to 110 kg	3.3	
Nitrogen input (kg) (k)	6.3	$\sum_1^i \frac{(\text{Dietary CP}_i \times \text{FCR}_i \times \text{Body weight gain}_i)}{6.25}$ with <i>i</i> the feeding phase index
Whole body nitrogen efficiency (%)	46	$\frac{(j-e)}{(k)} \times 100$
Nitrogen efficiency of a sow and her offspring		
Nitrogen intake (kg) (l)	199	$c + d + (27 \times k)$
Nitrogen retention (kg) (m)	85	$h + (27 \times j)$
Nitrogen efficiency (%)	43	m/l

¹Estimation based on commercially published feed recommendations and average performances reported by Government of Flanders (Belgium).

² Measured nitrogen content of gilts of different BW.

reduce N excretion mainly focus on the fattening phase and why this paper focuses exclusively on the fattening phase.

Nitrogen efficiency in pork production has improved through the application of scientific knowledge gained since

the 1980s; further improvements can be expected by implementing recent knowledge. The aim of the current paper is to critically evaluate existing knowledge and assumptions on strategies to maximize N efficiency in pork production.

We first discuss three major management strategies frequently linked with improved N efficiency: genetic selection, castration and slaughter weight. With every management strategy, adapted feeding is important for a correct evaluation of N efficiency. Because nutrition is the most important factor affecting N excretion, we then discuss nutritional strategies to maximize N efficiency. Although individual adapted feeding may yield benefits and represents a paradigm shift in pig production that some authors claim to be necessary (Andretta *et al.*, 2016), most pigs throughout the world are still housed and fed in groups. Improving the feeding practices for group-fed animals may therefore have the highest impact now and in the near future. Therefore, the main focus of this paper is on increasing efficiency in group-fed pigs. Based on the currently available scientific knowledge we estimate which level of N efficiency may be achieved with these group-fed pigs in the short term.

Calculating and expressing protein efficiency

On a fattening pig farm, piglets and nutrients can be considered as inputs. Outputs can be defined in different ways: kg live weight, kg carcass, kg lean meat or kg nutrients (e.g. kg N).

Theoretically, economic and environmental optimization of nutrient efficiency should be done per 'animal unit' on the farm while accounting for all trade-offs between inputs and outputs. Within a group, animals vary in their individual requirements and performances (Ferguson *et al.*, 1997; Pomar *et al.*, 2003). Under most practical (group housing) circumstances, the individual differences between pigs cannot be measured or managed. Optimization of individual nutrient efficiency, although desirable, may not be feasible in current practice. Nevertheless, nutrient efficiency can still be improved by taking measures at the population (barn) level.

Although stochasticity should be considered when adapting livestock management strategies (Pomar *et al.*, 2003), reasoning at the level of the 'average' animal results in robust yet simple calculations. This 'average' pig is a theoretical animal whose requirements, performances and efficiency can be directly calculated from measured performances at either pen or farm level.

In this 'average' pig, whole body N efficiency can be defined as the amount of N retained in the body divided by the amount of N ingested by the animal. Because the major aim of raising pigs is to produce meat, a more functional approach is to use the amount of N needed to produce one kg lean meat. Because protein needs are expressed relative to lysine (LYS) (see below), we propose using standardized ileal digestible (SID) LYS/kg lean meat as a functional measure for N efficiency.

Management strategies to improve nitrogen efficiency

Genetic selection

Genetic selection may affect N efficiency of group-fed pigs via two mechanisms: first, by directly selecting for increased efficiency, and second, by selecting for homogeneous groups of pigs.

Improving feed energy efficiency is a major objective of current animal breeding programs (Shirali *et al.*, 2012). Traditionally, improving energy efficiency was obtained by selecting for a lower feed conversion ratio. However, this approach may result in a reduction of feed intake, which in turn may limit further improvement of growth (Shirali, 2014). Therefore, residual feed intake (RFI) has been used as measure of feed efficiency, which is theoretically independent of lean growth (Shirali, 2014). Residual feed intake is the difference between the amount of feed (energy) the animal eats and the amount it is expected to eat based on requirements for maintenance and production (Lefaucheur *et al.*, 2011).

Without changing the diet, it is clear that pigs with a low feed conversion ratio (FCR) or low residual feed intake (LRFI) consume less, and hence excrete less N per kg of gain. However, while it seems logical to assume that dietary protein efficiency can be beneficially affected by genetic selection, selection has primarily focused on energy efficiency rather than on protein efficiency. Hammond (1947) stated that when environmental conditions limit the development of a character, it is not possible to select for genes that can be expressed when not hindered by environmental factors. Therefore, most breeding programs provide a diet designed to allow the animal to express its full genetic potential. When animals are fed diets where protein and amino acid (AA) level do not limit growth, increased protein efficiency is at best a side effect of selection for improved energy efficiency.

Regardless, that side effect does appear to be a reality: Moehn *et al.* (2004) observed that the rate of inevitable lysine catabolism decreases with increasing pig growth potential. A recent study reports that LRFI pigs have better protein efficiency than high RFI pigs (Cruzen *et al.*, 2013). The authors state that decreased muscle protein turnover may be an important reason for improved feed efficiency in LRFI pigs, based on measured enzyme activity. Still, in experiments with genetically different pig lines on feeds that were shown to limit growth, the marginal efficiency of protein use did not differ between breeds (Kyriazakis *et al.*, 1994; Susenbeth *et al.*, 1999). Marginal efficiency can be defined as the proportion of each increment in protein intake that is retained in the body. Trials with at least two (preferably more) protein levels are needed for this. Caution is needed when interpreting differences in muscle metabolism between pigs from different genetic lines that are fed only one type of diet, because the amount of AA intake relative to their requirement may differ considerably and may therefore evoke different responses. A higher degree of protein restriction should result in more efficient use of the feed, accompanied by lower muscle protein turnover. When selecting for increased N efficiency, it might be good to use diets with AA concentrations that limit growth, as we hypothesize that these diets should favor animals that use protein more efficiently. Apart from the marginal efficiency of N use described, a higher proportion of muscle N in the body would probably be linked to the amount of meat per kg of

N input and the ratio of muscle to maintenance N. Thus, the amount of SID lysine/kg feed should be higher, but the amount of SID lysine intake/kg lean meat is probably lower in lean compared with fat pigs.

Direct selection for increased N efficiency is an option but selection for lower variance in feed intake and protein accretion potential may also be a useful strategy to improve N efficiency of group-fed pigs. Ibanez-Escriche *et al.* (2008) stated that environmental variance of slaughter weight at 175 days in pigs may be partly genetically determined. The heritability estimates for the SD of BW at birth and at 3 weeks of age are around 0.1, hence worth selecting for (Canario *et al.*, 2010). With lower variation, more animals in the group can be fed adequately, thus decreasing inefficiencies in group-fed pigs (see below).

Castration

In most countries, castration of male piglets has been common practice until recently. Now societal pressure is leading many farmers to raise entire male pigs or immunocastrates (Millet *et al.*, 2011a). The pig sector in the EU has committed to ban surgical castration of male piglets by 2018. Boars have higher protein deposition capacity than either gilts or barrows. In terms of feed consumption, immunocastrates can be considered boars until the second vaccination (Millet *et al.*, 2011a), after which their feed intake drastically increases. Differences in N efficiency between barrows and boars are especially visible when using a reductionist approach with one type of diet characterized by adequate AA levels. For example, Van den Broeke *et al.* (2016) performed a study where four types of animals (boars, gilts, barrows, immunocastrates) received the same diets, formulated to fulfill the AA requirements of boars. In doing so, barrows consumed dietary protein in excess of their requirements, which was reflected in higher serum urea levels in barrows compared with boars. Similarly, a tremendous increase in serum urea level was observed after the second GnRH dosis in immunocastrates, in accordance with their increased feed intake, which also resulted in protein intake in excess of their requirements. While differences in nutrient requirements between genders are well established, little is known about gender-specific differences in marginal protein efficiency. In accordance with Moehn *et al.* (2004) who observed that the rate of inevitable lysine catabolism decreases with increasing pig growth potential, higher marginal lysine efficiency can be expected in boars compared with barrows. Moreover, as boars are leaner than barrows (Quiniou and Noblet, 1995) the ratio of muscle protein to total body protein and the ratio of protein for growth *v.* protein for maintenance may also be higher in boars *v.* barrows. Therefore, one could expect a general decrease in the level of SID lysine intake per kg lean meat when raising entire males *v.* barrows.

Slaughter weight

Shirali *et al.* (2012) stated that N excretion per BW gain rises with increasing BW. The question arises whether this is a result of decreasing marginal efficiency or non-adapted

feeding. Ghimire *et al.* (2016) observed no significant difference in lysine efficiency between growing and finishing pigs. Moehn *et al.* (2000 and 2004) stated that inevitable lysine catabolism and the marginal efficiency of using available lysine is independent of BW. In contrast, according to the National Research Council (NRC, 2012), empirical results suggest that the marginal efficiency of using SID lysine intake for protein deposition decreases with increasing BW, from 0.68 at 20 kg to 0.57 at 120 kg BW. Furthermore, maintenance AA requirements increase with increasing BW. Both imply a higher need for AA per kg of lean gain as BW increases.

While reduction of slaughter weight seems to lead to improved N efficiency, there is a trade-off between the input of piglets and feed (Van Meensel *et al.*, 2010). Lower slaughter weights imply a higher feed efficiency but also a higher number of piglets to produce 1000 kg of pork, and a higher number of sows. Increasing the slaughter weight decreases the number of fattening rounds per year and thus lowers the number of pigs, but it does imply increased feed costs (both economic and environmental). Increased BW also increases carcass yield (Wagner *et al.*, 1999; Correa *et al.*, 2006; Serrano *et al.*, 2008) without a clear effect on lean meat percentage (Correa *et al.*, 2006; Serrano *et al.*, 2008). Therefore, pigs that are too light or too heavy may both require higher amounts of AA per kg lean gain. In simulations of Morel and Wood (2005), where N excretion was taken into account (in contrast to economic optimization alone) the optimal slaughter weight decreased, especially in fat genotypes (115.2 kg with economic optimization alone *v.* 96.6 kg when placing a large emphasis on reducing N excretion). Note, however, that the authors assumed only one type of finisher feed independent of slaughter date.

Based on the above information, the effect of increased slaughter weight on N efficiency may be overestimated in practice, caused primarily by other factors such as excess protein supply in comparison with the requirements at higher BWs.

Nutritional strategies: feeding for maximal N efficiency in group-fed pigs

The three abovementioned management strategies (genetic selection, castration and slaughter weight) affect the amount of SID lysine per kg lean meat. When using a reductionist approach (i.e. the same diet for all experimental pigs) to study the effects of these management factors, a large part of the observed differences in studies can be attributed to the diet. Obviously, when feeding barrows the same diet as gilts or when feeding 150 kg pigs the same diet as 100 kg pigs, the intrinsic differences will be exaggerated. Therefore, it is important to feed the right diet for each type of animal, both in practice and in experimental studies. To optimise N efficiency, several nutritional concepts should be taken into account. Briefly, the amount of excreted N and the route of excretion (fecal or urinary) depends on the amount of ingested N, the fraction of absorbed N, the animal's AA requirements and the AA balance in the diet.




Figure 1 Schematic illustration of the dietary nutrient concentration of three feeding systems differing in the subdivision of feeding phases: one-phase (—), three-phase (.....) and multiphase (.....) feeding. The figure on the top (a) shows the theoretical concept, while the figure on the bottom (b) shows the translation into practice.

Meeting the requirements of group-fed pigs throughout different stages of growth

Phase feeding (adapting the dietary AA content to the physiological needs of an animal during its different life stages) is a recognized strategy to lower N inputs and outputs while maintaining maximal performance (Han *et al.*, 2000). Terms such as one-phase, three-phase and multiphase feeding all share the same aim of providing the pig with sufficient nutrients at each time point (Figure 1a). As the number of phases increases, theoretically the amount of ingested protein will decrease and will therefore better match the animal's nutritional requirements over time. Pomar *et al.* (2014) estimated a 12% reduction in N excretion by switching from three-phase feeding to daily multiphase feeding in individually fed pigs. Of course, the amount of reduction depends on the control treatment to which it is compared. In their study, whole body N efficiency was 37% on three-phase feeding and 40% on the daily multiphase feeding strategy. Andretta *et al.* (2016) reached up to 57% N efficiency in individually fed pigs given a daily-phase feeding program designed to meet 80% of the estimated nutritional requirements, compared with 45% in a three-phase feeding program. In a recent trial at Flanders Research Institute for Agriculture, Fisheries and Food (ILVO, Melle, Belgium), 54% N efficiency was obtained in group-fed boars in a three-phase feeding system with diets formulated in line with commercial practice (Van den Broeke *et al.*, 2017). However, there is a large difference between theoretical and practical phase-feeding systems. While, in theory, the AA

levels proposed for three-phase feeding are sufficient at the beginning and in excess at the end of the feeding phase (Figure 1a), in the commercial three-phase feeding system nutrients are limiting at the beginning and in excess at the end of the feeding phase (Figure 1b).

As stated above, group housing is common practice for piglets and fattening pigs on farms. Pigs are fed and housed per age group. However, individual pigs of the same age group also differ in protein deposition capacity and hence may differ in AA requirement. This variation is important when formulating recommendations for feeding pigs in groups and may explain differences in research results on individual or group level. When feeding to evoke optimal responses of a group of pigs, many of the pigs receive excess nutrients. Therefore, even when feeding below the requirement for optimal performance of a group in a commercial three-phase feeding strategy, a considerable number of the pigs in the group are still likely to have their nutritional needs met at all times. Furthermore, compensatory growth mechanisms may be at play: several authors report that pigs subjected to early dietary AA restrictions may compensate and decrease N excretion during both the restriction and re-alimentation phases (Fabian *et al.*, 2004; Millet *et al.*, 2011b; Millet and Aluwé, 2014). Because results among studies do not always agree (De Greef *et al.*, 1992; Chiba *et al.*, 2002), it is difficult to generate general recommendations for maintaining profitability while minimizing N excretion through short-term dietary protein deficiencies. Despite clear compensatory growth responses after AA restriction in gilts (Millet *et al.*, 2011b) and barrows (Millet and Aluwé, 2014), the best (numerical) feed efficiency was still observed in the pigs that had never been restricted, while the highest lysine efficiency was seen in piglets that were fed an AA restricted diet throughout their life. In these studies, the lowest amount of total lysine per kg lean meat gain reached was 45.4 g in barrows and 42.1 g in gilts. Similar improvements at low lysine levels were found by Ghimire *et al.* (2016), where higher efficiency of lysine utilization was observed at lower levels of lysine intake. Moehn *et al.* (2004) only observed a reduced rate of lysine catabolism at the lowest lysine intake level (40% below requirements). Those two studies used data on individual piglets. As stated above, feeding pigs below the group optimum may increase lysine efficiency by decreasing the variance in lysine utilization. In conclusion, daily multiphase feeding adapted to the individual animal's needs is probably the most efficient in terms of N efficiency. For animals housed in groups, phase feeding appears to be required for optimizing N efficiency on group level. When combined with periods of temporary AA restriction, this efficiency can be further improved.

Optimal standardized ileal digestible lysine : digestible crude protein ratio

The increasing availability of feed grade AA makes it possible to decrease the CP content in the diet. While it is generally accepted that animals need AA rather than CP, the question remains whether there is a lower limit to protein provision.

Table 2 Maximal dietary standardized ileal digestible (SID)¹ lysine:CP ratios reported in scientific literature that can be used without negative effects on performance

BW range	SID LYS (g/kg)	LYS (g/kg)	CP (g/kg)	SID LYS : CP	LYS : CP	LYS : ATTD CP	Reference
19 to 48 kg	7.38 ²	8.2	120	0.062 ²	0.068	0.085 ²	Figuroa <i>et al.</i> (2002)
8 to 25 kg	10.0	11.1 ²	160	0.063	0.069 ²	0.087 ²	Jansman <i>et al.</i> (2016)
7 to 19 kg	13	14.3	194	0.067	0.074	0.092 ²	Nemeček <i>et al.</i> (2014)
12 to 20 kg	11.5	12.5	165	0.070	0.076	0.095 ²	Gloaguen <i>et al.</i> (2014) (experiment 1)
12 to 20 kg	9.2	10	134	0.069	0.075	0.093 ²	Gloaguen <i>et al.</i> (2014) (experiment 2)
8 to 24 kg	11.5	12.8	180	0.064	0.071	0.089 ²	Millet <i>et al.</i> (2017) ³ (linear plateau)
8 to 24 kg	12.9	14.2	180	0.072	0.079	0.099 ²	Millet <i>et al.</i> (2017) ³ (quadratic plateau)

¹The first five studies have been performed with a fixed SID LYS content and varying CP level.

²Values in italics have been estimated based on following assumptions: SID LYS : LYS = 0.9; apparent total tract digestible (ATTD) CP/CP = 0.8.

³In the study of Millet *et al.* (2017), CP was fixed at 180 g/kg and SID LYS varied.

Wu (2014) stated that minimal levels are also required for non-essential AA. But N itself can also be limiting; Mansilla *et al.* (2015) showed that N absorbed from the large intestine can be used when animals are fed diets deficient in dispensable AA nitrogen. Some studies have been performed on the optimal essential : total N ratio (Mitchell *et al.*, 1968; Heger *et al.*, 1998; Lenis *et al.*, 1999). However, essential AA are only essential up to the point where they no longer limit growth. Essential AA in excess can be deaminated and utilized for the synthesis of non-essential AA (Lenis *et al.*, 1999). Therefore, the ratio of SID lysine to apparent total tract digestible (ATTD) CP may be a more helpful measure to improve N efficiency; 25 years ago, Henry and Dourmad (1993) suggested that the crude lysine : protein ratio should not exceed 0.065 to 0.068. This ratio was suggested to limit the risk for deficiencies in non-essential AA or in essential AA that were not taken into account. Since then, a large body of work has further clarified AA requirements. As knowledge increases about the requirement of all the essential AA, it may become possible to decrease dietary CP content even further. Recently, several studies have tested decreases of CP level while maintaining (SID) lysine (LYS) level (Table 2). When the corresponding SID LYS : CP ratio was calculated, the maximal ratio varied between 0.062 and 0.070 (total lysine : CP between 0.068 and 0.076). Using this maximum in feed formulation may enable a decrease in CP during the growing-finishing period. If this ratio is used during the piglet phase (first weeks after weaning), the CP level may determine the SID LYS level. Indeed, a decreased CP level in piglet rations helps to maintain intestinal health (Nyachoti *et al.*, 2006). In two recent trials (Millet *et al.*, 2017), performances improved linearly with an SID LYS level between 8.5 and 13.5 and a corresponding CP level varying between 201 and 210 g/kg. In contrast, when CP was fixed at 180 g/kg, the SID LYS level for optimal FCR was 11.4 based on a linear plateau model and 12.9 based on a quadratic plateau model. Assuming that CP was limiting performance at the highest SID LYS levels, this would yield a maximal SID LYS : CP level of 0.064 or 0.072 (Millet *et al.*, 2017, Table 2). Further research is needed to determine the maximal SID LYS : CP level that does not negatively affect performance in different growth phases. Given the current knowledge, a ratio of

0.063 (0.07 LYS : CP) may be a safe choice. As N absorbed from the large intestine can also be used (Mansilla *et al.*, 2015), ideally ATTD CP should be determined in these trials and the maximum expressed as SID LYS : ATTD CP. Assuming an ATTD CP digestibility of 80%, with the studies mentioned in Table 2, the calculated SID LYS : ATTD CP in the studies mentioned in Table 2 would be between 0.077 and 0.087.

Optimal amino acid balance

Animal protein requirements are based on intake of a complete set of AA instead of CP. AAs given in excess are deaminated and the resulting urea is excreted in the urine (van Milgen and Dourmad, 2015). Decreasing the dietary CP content while maintaining optimal SID AA concentrations has been proven successful to reduce N input per kg of lean meat gain. This can be obtained by combining highly digestible AA sources and formulation of feeds for an optimal AA composition. Single AA deficiencies lead to inefficient use of the other AA, which are in turn deaminated and excreted in the urine, causing suboptimal growth. This knowledge has led to the well-known concept of 'ideal protein,' which varies with physiological state and level of productivity of the animal (NRC, 2012) and has been extensively discussed elsewhere (Boisen *et al.*, 2000; van Milgen and Dourmad, 2015). Now that feed grade crystalline AA are available, feed can be formulated close to the ideal AA pattern to be used in research and commercial practice. Although the ideal protein concept is clear, the methods to deduce the optimal balance between AA are still under discussion and different methodologies still yield (slightly) different results.

Estimation of maximal nitrogen efficiency attainable in group-fed pigs

Estimations of the maximal N efficiency for producing marketable pork meat were calculated from data presented in Table 3. With 45 g of LYS/kg lean meat gain as value we observed in several studies (Millet *et al.*, 2011b; Millet and Aluwé, 2014) and 0.07 as a safe LYS : CP ratio, we calculate an N efficiency of 57% for pigs between 8 and 110 kg (total lysine was used in accordance with other studies; Table 3). With further research, the amount of lysine per kg lean meat

Table 3 Estimate of nitrogen efficiency attainable with group-fed pigs

Parameters	Estimate	Calculation	References
Lean meat in 8 kg pig (kg) (a)	3.6	45% of live weight	Susenbeth and Keitel (1988)
Lean meat in 110 kg pig with 62% meat (kg) (b)	53.2	BW (110 kg) × carcass yield (78%) × lean meat percentage	
Lean meat growth (kg) (c)	49.6	$b - a$	
Minimum lysine ingested (g) (d)	2232	$c \times 45$	Millet <i>et al.</i> (2011b), Millet and Aluwé (2014) ¹
Minimum CP ingested (kg) (e)	31.9	$\frac{c}{0.07} \times 1000$	Table 2 ²
Nitrogen ingested (kg) (f)	5.1	$\frac{e}{6.25}$	
Nitrogen retained (kg) (g)	2.9		Table 1 ³
Nitrogen efficiency (%)	57	$\frac{g}{f} \times 100$	

¹Grams of lysine needed per kg of lean meat growth that have been observed in studies at ILVO. This is probably an overestimation of the minimum.

²Maximal dietary lysine: CP ratio that still allows maximal performance reported in literature ranges between 0.068 and 0.079. In this calculation, 0.070 was chosen as a conservative estimate.

³Nitrogen content of a 110 kg pig minus nitrogen content of an 8 kg piglet.

may decrease and the LYS:CP ratio may be increased, leading to an even higher efficiency. The availability of different protein sources or commercially available AA may interfere with reaching a 0.07 LYS:CP ratio while maintaining a correct AA balance. On the other hand, when applying the management choices of precision feeding, genetic selection and raising entire male pigs, the amount of lysine per kg lean meat gain can probably be reduced even further. More research is needed before this can be achieved. Furthermore, economic studies related to the management choices and the cost of technology are required. Although the proposed efficiency is much higher than the 33% reported for growing pigs under practical circumstances in the Netherlands, France and Denmark (Dourmad *et al.*, 1999) and the 46% estimated for contemporary Belgian pig production (Table 1), N efficiency close to 60% appears to be achievable in the near future in group-fed fattening pigs.

Acknowledgments

Thanks to Miriam Levenson for language correction.

References

- Andretta I, Pomar C, Rivest J, Pomar J and Radunz J 2016. Precision feeding can significantly reduce lysine intake and nitrogen excretion without compromising the performance of growing pigs. *Animal* 10, 1137–1147.
- Boisen S, Hvelplund T and Weisbjerg MR 2000. Ideal amino acid profiles as a basis for feed protein evaluation. *Livestock Production Science* 64, 239–251.
- Canario L, Lundgren H, Haandlykken M and Rydhmer L 2010. Genetics of growth in piglets and the association with homogeneity of body weight within litters. *Journal of Animal Science* 88, 1240–1247.
- Chiba LI, Kuhlers DL, Frobish LT, Jungst SB, Huff-Lonergan EJ, Lonergan SM and Cummins KA 2002. Effect of dietary restrictions on growth performance and carcass quality of pigs selected for lean growth efficiency. *Livestock Production Science* 74, 93–102.
- Correa JA, Faucitano L, Laforest JP, Rivest J, Marcoux M and Garipey C 2006. Effects of slaughter weight on carcass composition and meat quality in pigs of two different growth rates. *Meat Science* 72, 91–99.
- Cruzen SM, Harris AJ, Hollinger K, Punt RM, Grubbs JK, Selsby JT, Dekkers JCM, Gabler NK, Lonergan SM and Huff-Lonergan E 2013. Evidence of decreased muscle protein turnover in gilts selected for low residual feed intake. *Journal of Animal Science* 91, 4007–4016.
- CVB 2016. Feed Table for pigs. Voedernormen Varkens en voederwaarden voedermiddelen voor Varkens. Federatie Nederlandse Diervoederketen, Rijswijk, and Wageningen Livestock Research, Wageningen, the Netherlands.
- De Greef KH, Kemp B and Verstegen MWA 1992. Performance and body-composition of fattening pigs of 2 strains during protein-deficiency and subsequent realimentation. *Livestock Production Science* 30, 141–153.
- Dourmad JY, Seve B, Latimier P, Boisen S, Fernandez J, van der Peet-Schwering C and Jongbloed AW 1999. Nitrogen consumption, utilisation and losses in pig production in France, The Netherlands and Denmark. *Livestock Production Science* 58, 261–264.
- Fabian J, Chiba LI, Frobish LT, McElhenney WH, Kuhlers DL and Nadarajah K 2004. Compensatory growth and nitrogen balance in grower-finisher pigs. *Journal of Animal Science* 82, 2579–2587.
- Ferguson NS, Gous RM and Emmans GC 1997. Predicting the effects of animal variation on growth and food intake in growing pigs using simulation modelling. *Animal Science* 64, 513–522.
- Figuerola JL, Lewis AJ, Miller PS, Fischer RL, Gomez RS and Diedrichsen RM 2002. Nitrogen metabolism and growth performance of gilts fed standard corn-soybean meal diets or low-crude protein, amino acid-supplemented diets. *Journal of Animal Science* 80, 2911–2919.
- Ghimire S, Pomar C and Remus A 2016. Variation in protein content and efficiency of lysine utilisation in growing-finishing pigs. In *Energy and protein metabolism and nutrition: 5th EAAP International Symposium on Energy and Protein Metabolism and Nutrition*, 12 to 15 September 2016, Krakow, Poland, pp. 351–352.
- Gloaguen M, Le Floc'h N, Corrent E, Primot Y and van Milgen J 2014. The use of free amino acids allows formulating very low crude protein diets for piglets. *Journal of Animal Science* 92, 637–644.
- Hammond J 1947. Animal breeding in relation to nutrition and environmental conditions. *Biological Reviews of the Cambridge Philosophical Society* 22, 195–213.
- Han IK, Lee JH, Kim JH, Kim YG, Kim JD and Paik IK 2000. Application of phase feeding in swine production. *Journal of Applied Animal Research* 17, 27–56.
- Heger J, Mengesha S and Vodehnal D 1998. Effect of essential:total nitrogen ratio on protein utilization in the growing pig. *British Journal of Nutrition* 80, 537–544.
- Henry Y and Dourmad J 1993. Feeding strategy for minimizing nitrogen output in pigs. In *Nitrogen flow in pig production and environmental consequences*. (ed. MWA Verstegen, LA den Hartog, GJM van Kempen, JHM Metz), pp. 137–150. EAAP Publication, Wageningen, the Netherlands.
- Ibanez-Escriche N, Varona L, Sorensen D and Noguera JL 2008. A study of heterogeneity of environmental variance for slaughter weight in pigs. *Animal* 2, 19–26.

- Jansman A, van Diepen J, Rovers M and Corrent E 2016. Lowering the dietary protein content in piglets: How far can we go? In *Energy and protein metabolism and nutrition: 5th EAAP International Symposium on Energy and Protein Metabolism and Nutrition*, 12 to 15 September 2016, Krakow, Poland, pp. 161–162.
- Kyriazakis I, Dotas D and Emmans GC 1994. The effect of breed on the relationship between feed composition and the efficiency of protein-utilization in pigs. *British Journal of Nutrition* 71, 849–859.
- Lefaucheur L, Leuret B, Ecolan P, Louveau I, Damon M, Prunier A, Billon Y, Sellier P and Gilbert H 2011. Muscle characteristics and meat quality traits are affected by divergent selection on residual feed intake in pigs. *Journal of Animal Science* 89, 996–1010.
- Leip A, Weiss F, Lesschen JP and Westhoek H 2014. The nitrogen footprint of food products in the European Union. *Journal of Agricultural Science* 152, S20–S33.
- Lenis NP, van Diepen HT, Bikker P, Jongbloed AW and van der Meulen J 1999. Effect of the ratio between essential and nonessential amino acids in the diet on utilization of nitrogen and amino acids by growing pigs. *Journal of Animal Science* 77, 1777–1787.
- Mansilla WD, Columbus DA, Htoo JK and de Lange CFM 2015. Nitrogen absorbed from the large intestine increases whole-body nitrogen retention in pigs fed a diet deficient in dispensable amino acid nitrogen. *Journal of Nutrition* 145, 1163–1169.
- Millet S and Aluwé M 2014. Compensatory growth response and carcass quality after a period of lysine restriction in lean meat type barrows. *Archives of Animal Nutrition* 68, 16–28.
- Millet S, Aluwé M, De Paepe M, De Brabander DL and Van Oeckel MJ 2010. Effect of decreasing ideal protein levels on performance results and nitrogen efficiency of growing-finishing gilts. *Archives of Animal Nutrition* 64, 1–11.
- Millet S, Aluwé M, Le Gall E, Corrent E, Lambert W, De Sutter J, Ampe B and De Campeneere S 2017. The optimal standardised ileal lysine requirement of piglets between 8 and 24 kg. *Journées de la Recherche Porcine* 49, 121–122.
- Millet S, Gielkens K, De Brabander D and Janssens GPJ 2011a. Considerations on the performance of immunocastrated male pigs. *Animal* 5, 1119–1123.
- Millet S, Langendries K, Aluwé M and De Brabander DL 2011b. Effect of amino acid level in the pig diet during growing and early finishing on growth response during the late finishing phase of lean meat type gilts. *Journal of the Science of Food and Agriculture* 91, 1254–1258.
- Mitchell JR, Becker DE, Harmon BG, Norton HW and Jensen AH 1968. Some amino acid needs of the young pig fed a semisynthetic diet. *Journal of Animal Science* 27, 1322–1326.
- Moehn S, Gillis AM, Moughan PJ and de Lange CFM 2000. Influence of dietary lysine and energy intakes on body protein deposition and lysine utilization in the growing pig. *Journal of Animal Science* 78, 1510–1519.
- Moehn S, Ball RO, Fuller MF, Gillis AM and de Lange CFM 2004. Growth potential, but not body weight or moderate limitation of lysine intake, affects inevitable lysine catabolism in growing pigs. *Journal of Nutrition* 134, 2287–2292.
- Morel P and Wood G 2005. Optimisation of nutrient use to maximise profitability and minimise nitrogen excretion in pig meat production systems. *Acta Horticulturae* 674, 269–275.
- Nemeček JE, Tokach MD, Drits SS, Goodband RD and DeRouchey JM 2014. Evaluation of standardized ileal digestible valine:lysine, total lysine:crude protein, and replacing fish meal, meat and bone meal, and poultry byproduct meal with crystalline amino acids on growth performance of nursery pigs from seven to twelve kilograms. *Journal of Animal Science* 92, 1548–1561.
- National Research Council 2012. *Nutrient requirements of swine*, 11th revised edition. National Academy Press, Washington, DC, USA.
- Nyachoti CM, Omogbenigun FO, Rademacher M and Blank G 2006. Performance responses and indicators of gastrointestinal health in early-weaned pigs fed low-protein amino acid-supplemented diets. *Journal of Animal Science* 84, 125–134.
- Pomar C, Kyriazakis I, Emmans G and Knap P 2003. Modeling stochasticity: dealing with populations rather than individual pigs. *Journal of Animal Science* 81, E178–E186.
- Pomar C, Pomar J, Dubeau F, Joannopoulos E and Dussault JP 2014. The impact of daily multiphase feeding on animal performance, body composition, nitrogen and phosphorus excretions, and feed costs in growing-finishing pigs. *Animal* 8, 704–713.
- Quiniou N and Noblet J 1995. Prediction of tissular body-composition from protein and lipid deposition in growing pigs. *Journal of Animal Science* 73, 1567–1575.
- Serrano MP, Valencia DG, Fuentetaja A, Lazaro R and Mateos GG 2008. Effect of gender and castration of females and slaughter weight on performance and carcass and meat quality of Iberian pigs reared under intensive management systems. *Meat Science* 80, 1122–1128.
- Shirali M 2014. *Improvement of energy and nitrogen utilisation in pork production: genetics and growth models*. PhD thesis, Wageningen University, Wageningen, the Netherlands.
- Shirali M, Doeschl-Wilson A, Knap PW, Duthie C, Kanis E, van Arendonk JAM and Roehe R 2012. Nitrogen excretion at different stages of growth and its association with production traits in growing pigs. *Journal of Animal Science* 90, 1756–1765.
- Susent Beth A and Keitel K 1988. Partition of whole-body protein in different body fractions and some constants in body-composition in pigs. *Livestock Production Science* 20, 37–52.
- Susent Beth A, Dickel T, Diekenhorst A and Hohler D 1999. The effect of energy intake, genotype, and body weight on protein retention in pigs when dietary lysine is the first-limiting factor. *Journal of Animal Science* 77, 2985–2989.
- Van den Broeke A, Leen F, Aluwé M, Ampe B, Van Meensel J and Millet S 2016. The effect of GnRH vaccination on performance, carcass, and meat quality and hormonal regulation in boars, barrows, and gilts. *Journal of Animal Science* 94, 2811–2820.
- Van den Broeke A, Leen F, Aluwé M, Van Meensel J and Millet S 2017. Effect of slaughter weight and sex on carcass composition, N- and P- efficiency of pigs. Paper presented at the 68th Annual Meeting of the European Federation of Animal Science (EAAP), 28 August to 1 September 2017, Tallinn, Estonia.
- Van Meensel J, Lauwers L and Van Huylbroeck G 2010. Communicative diagnosis of cost-saving options for reducing nitrogen emission from pig finishing. *Journal of Environmental Management* 91, 2370–2377.
- van Milgen J and Dourmad JY 2015. Concept and application of ideal protein for pigs. *Journal of Animal Science and Biotechnology* 6, 15.
- Wagner JR, Schinckel AP, Chen W, Forrest JC and Coe BL 1999. Analysis of body composition changes of swine during growth and development. *Journal of Animal Science* 77, 1442–1466.
- Warnants N, Van Oeckel MJ, De Paepe M, Millet S and De Brabander DL 2006. Evolutie van de lichaamssamenstelling van big tot slachtvarken. In *Infonamiddag ILVO-DIER "Actuele onderzoeksresultaten bij éénmagigen"*, Melle (BE), pp. 35–50.
- Wu GY 2014. Dietary requirements of synthesizable amino acids by animals: a paradigm shift in protein nutrition. *Journal of Animal Science and Biotechnology* 5, 34.