maintenance actions performed. For the equipment in the administrative, support and research areas, the function and physical risk criteria were replaced by the safety and by the risks to the quality of service criteria. The evaluation is carried out by a multidisciplinary team. The tool categorizes the equipment into low, medium and high criticality.

Conclusions. The tool prioritized the equipment based on objective criteria evaluated by the departments' multidisciplinary team comprising experts who use the equipment in their activities, the department administrator and clinical engineers, and provided transparency regarding the decision-making of the hospital's Investment Committee. In 2019, the limited financial resources were invested only in the replacement of highly critical equipment. We believe the tool can be reproduced in hospitals in low and middle-income countries.

PP499 Disinvestment Supported By A Hospital-Based Health Technology Assessment Unit: A Case Of A Teaching Hospital In Brazil

Rosane Paixão Schlatter (rschlatter@hcpa.edu.br), Katiuce Tomazi Kny and Maria Angelica Pires Ferreira

Introduction. The Hospital de Clínicas de Porto Alegre (HCPA), a public teaching hospital, has a Hospital-based Health Technology Assessment (HB-HTA) unit to support the decisionmaking process on technology incorporation, rationalization or disinvestment. In 2017, the plastic adhesive drape was standardized at HCPA for use in cardiovascular, digestive, orthopedic, and neurological surgery for the purpose of preventing surgical site infection (SSI). This study evaluated whether the plastic adhesive drape technology is more effective than the no adhesive drapes in the surgical procedures in which it is used in the HCPA, so as to support the medical board's decision regarding the rationalization of use.

Methods. The primary outcome was the surgical site infection rate (SSI). Searches were performed in PubMed, Cochrane and national and international health agencies: World Health Organization (WHO), National Institute for Health and Care Excellence (NICE), Centers for Disease Control and Prevention (CDC), Society for Healthcare Epidemiology of America (SHEA), Brazilian National Commission for the Incorporation of Technologies (CONITEC) and Brazilian National Health Surveillance Agency (ANVISA) databases. The search strategy combined terms related to the technology and types of surgery in which it is used in the HCPA. The quality of the included studies was assessed. Additionally, data on technology utilization and costs in the hospital were analyzed.

Results. Technology assessment followed AdHopHTA project recommendations. Data from the hospital showed that the technology has been used in fifteen surgical specialties, different from the proposed incorporation, with a progressive increase in consumption from 2017 to 2018. The literature review included a systematic review with seven clinical trials, which concluded that the plastic adhesive drape lacks benefits, with potential for increased risk of SSI. The evidence was of moderate quality.

Conclusions. The expenses associated with the use of the technology were considered unjustified as it is not reimbursed by the Brazilian Ministry of Health and its disinvestment was recommended. The Medical Board approved the disinvestment of the technology based on the evidence found by the HB-HTA unit, and the medical staff complied with the decision.

PP501 Inclusion Of Key Stakeholders' Views When Developing A mHealth Assessment Tool: Focus Groups And Health Consensus Results

Elisa Puigdomènech, Elisa Poses, César Velasco, Xavier Garcia (xavier.garciacusco@gencat.cat) and Mireia Espallargues

Introduction. The Agency for Health Quality and Assessment of Catalonia (AQuAS) is developing an evaluation tool for mobile health (mHealth) solutions to be used by health technology assessment (HTA) agencies and evaluation experts. In order to have a practical and comprehensive tool taking into account the particularities and challenges of mobile interventions, we considered the views and opinions of key stakeholders. The objective was to present the final selection of general aspects (dimensions) to be assessed in the evaluation, as well as the specific items (criteria) to be included in each of these topics, as a result of different co-design approaches with health professionals, developers, hospital managers, HTA agencies and patients.

Methods. A list of criteria used for health apps evaluation were drawn from a literature review. The initial list included eightynine criteria items grouped in nine domains. Those criteria and domains were discussed during four focus groups (FG). The importance of the criteria that were not considered as mandatory were later rated through a Delphi online sub-study, in a scale from one to six points, taking as consensus value when median value (median 6, Interquartile range, 0–1) was reached.

Results. FG reduced domains and criteria from nine to seven and from eighty-nine to thirty-three, respectively. Most mandatory criteria were related with security, user experience, and clinical effectiveness. Fifty-seven individuals (53.7% of 106 invited to participate) were registered in the online platform (50.1% women, 68.4% 35–64years old and 42.1% from HTA agencies). From fifty non-mandatory criteria under consensus, ten criteria reached consensus (most from solution's content and health problem covered domains) concluding with a 43/7 criteria/domain tool.

Conclusions. Insights from main stakeholders on the content of the tool for mHealth assessment were considered through the FG and Delphi technique. The dimensions of security and privacy, clinical effectiveness, solutions' content, technological aspects, users' experience and costs were considered mandatory. The dimension related to the impact on the organization was appraised as a secondary domain for evaluation. A workshop with AQuAS research team and HTA external researchers will help to define: the assessment methods (type of instrument,