

the keywords “entrepreneurship”, “health education” and “health personnel”, on March 8th, 2018. Results were screened by title, abstract and full text by a team of three calibrated researchers, based upon pre-defined exclusion and inclusion criteria. The final list of papers was reviewed using an extraction tool to identify demographics, details of the paper, and its attitudes and perceptions towards entrepreneurship and innovation. **Results:** After screening, 59 papers were identified for qualitative analysis. These papers ranged from 1970–2018, mainly from the USA (n = 36). Most papers were commentaries/opinions (n = 35); 11 papers described specific innovations. Entrepreneurship was viewed positively in 45 papers, negatively in 2 papers, and mixed in 12 papers. Common specialties discussed were surgery (n = 9), internal medicine (n = 3), and not specified (n = 44). Emergency medicine was described in one paper. Major themes were: entrepreneurial environment (n = 29), funding and capital (n = 12), idea generation (n = 9), and teaching entrepreneurship (n = 6). Of the 11 innovation papers, the discussion was focused on educational (n = 6) or system (n = 5) innovations. These innovations related to surgery (n = 1), public health (n = 1) and palliative care (n = 1). None of these innovations were specific to emergency medicine. **Conclusion:** This review indicates a small number of programs focused on promoting innovation and entrepreneurship amongst trainees, but no programs specific to the emergency department. There may be benefit for educators in emergency medicine to consider how to foster a greater innovative spirit in our speciality, so our next generation of physicians can help tackle problems affecting patient care.

Keywords: entrepreneurship, health education, innovation

P127

Health information technology and the Ontario emergency department return visit quality program - A population level continuous quality improvement program

A. Taher, MD, MPH, E. Bunker, MPH, MSc, L. Chartier, MD, MPH, H. Ovens, MD, B. Davis, MBA, M. Schull, MD, MSc, University of Toronto, Toronto, ON

Introduction: Emergency department (ED) return visits are used for quality monitoring. Health information technology (HIT) has historically supported return visit programs in the same hospital or hospital system. The Emergency Department Return Visit Quality Program (EDRVQP) is a novel population level continuous quality improvement (QI) program connecting EDs across Ontario that leverages HIT. We sought to describe the EDRVQP HIT architecture, experience of participants, enabling program factors and barriers. **Methods:** The Informatics Stack conceptual framework was used to describe the HIT architecture. A literature review of peer-reviewed background literature, and stakeholder organization reports was conducted. Purposive sampling identified key informants. Semi-structured interviews were conducted until saturation. Common themes were identified by inductive qualitative thematic analysis. **Results:** Twenty-three participants from 15 organizations were interviewed. The EDRVQP architecture description is presented across the Informatics Stack. The levels from most comprehensive to most basic are world, organization, perspectives/roles, goals/functions, workflow/behaviour/adoption, information systems, modules, data/information/knowledge/algorithms, and technology. Enabling factors were a high rate of EHR adoption, provincial legislative mandate for data collection and program membership, use of functional and data standards, local variability, phased deployment, and QI and patient safety culture. Two main barriers were increased case turnaround time and privacy legislation.

Conclusion: The Informatics Stack framework provides a robust approach to thoroughly describe the HIT architecture of this population health programs. The EDRVQP is a population health program that illustrates the pragmatic use of continuous QI methodology across a population (provincial) level.

Keywords: emergency department, information technology, public health informatics

P128

Describing variability in treatment of THC hyperemesis in the emergency department: a health records review

J. Teeffy, MD, J. Blom, PhD, K. Woolfrey, MD, M. Riggan, MD, J. Yan, MD, London Health Sciences Centre, London, ON

Introduction: Cannabis Hyperemesis Syndrome (CHS) is a new and poorly understood phenomenon with a subset of patients presenting to emergency departments (ED) for symptomatic control of their refractory nausea and vomiting. Currently, there is a lack of agreement and considerable practice variability on initial treatment modalities for CHS. The objective of this study was to describe the treatment modalities for patients presenting to ED with cannabis-related sequelae.

Methods: This was a health records review of patients ≥ 18 years presenting to one of two tertiary care EDs (annual census: 150,000) with a discharge diagnosis including cannabis use with one of abdominal pain or nausea/vomiting using ICD-10 codes. Trained research personnel collected data from medical records including demographics, clinical history, results of investigations, and utilization of treatment options within the ED. Descriptive statistics are presented where appropriate. **Results:** From April 2014 to June 2016, 203 unique ED patients had a discharge diagnosis including cannabis use with abdominal pain or nausea/vomiting. Sixty-nine (33.4%) received any treatment during their visit with 28 (40.6%) receiving IV fluids, of which 24 (85.7%) received normal saline. Anti-emetics were used in 21 (30.4%) patients with ondansetron being the first-line agent in 11 (52.4%) patients followed by dimenhydrinate in 6 (28.6%) and haloperidol in 2 (9.5%) cases. Six patients required two doses of anti-emetics, favouring ondansetron in 3 cases followed by haloperidol, dimenhydrinate, and metoclopramide each used once. Thirteen (19%) patients required analgesia, with the first-line preference being non-opioid medications in 11 versus opioids in 2 cases. Seven patients required multiple modes of analgesia, favouring opioid medications in 4 patients. Twenty-eight (40.6%) patients required anxiolytics with lorazepam being used primarily in 16 (57.1%) patients followed by lorazepam/haloperidol in 5 (17.9%) cases. **Conclusion:** This ED-based study demonstrates variability of practice patterns for symptomatic treatment of cannabis related ED presentations. Despite knowledge of haloperidol being useful in patients with suspected CHS, physicians opted for ondansetron as first line anti-emetics. Future research should focus on studying various treatment modalities of patients with suspected CHS in the ED to optimize symptomatic treatment.

Keywords: cannabis, nausea, pain

P129

Safer transitions in the care of the elderly: identification of essential information in transitional care

S. Trivedi, MD, S. Beckett, BSc, A. Dick, BSc, SCBScN, R. Hartmann, MSc, MD, C. Roberts, MD, K. Lyster, MD, J. Stempien, MD, Royal University Hospital, Saskatoon, SK

Introduction: When presenting to the Emergency Department (ED), the care of elderly patients residing in Long Term Care