

Effects of befriending on depressive symptoms: a precautionary note on promising findings

Mead *et al*¹ recently meta-analysed data on the effectiveness of befriending interventions on reducing depressive symptoms. Befriending was defined as a non-professional intervention that provides clients with non-directive, emotionally focused support by one or more individuals; was not psychoeducational or mentoring in nature; and did not constitute formal psychotherapy. Mead *et al* found that befriending interventions had a modest, statistically significant effect on depressive symptoms within 12 months of randomisation (standardised mean difference 0.27, 95% CI 0.06 to 0.48, nine studies) and a slightly smaller effect on longer-term outcomes (standardised mean difference 0.18, 95% CI 0.05 to 0.32, five studies).

As the authors noted, the effect sizes for befriending were essentially equivalent to effect sizes from collaborative care depression interventions in primary care. In a 2006 meta-analysis, Gilbody *et al*² reported a short-term (within 6 months) standardised mean difference effect size for symptom reduction from collaborative or enhanced depression care of 0.25 (95% CI 0.18 to 0.32, 35 studies) and longer-term effect sizes of 0.15 at 2 years post-randomisation (95% CI -0.03 to 0.32, 9 studies) and 0.15 at 5 years post-randomisation (95% CI 0.001 to 0.30, 2 studies). As Mead and colleagues note, the implications of this are important. Befriending or social support interventions could provide a less expensive and potentially 'less medicalised' option of care for patients with mild to moderate symptoms of depression in primary care. Indeed, collaborative care is a complex, multifaceted, expensive organisational intervention that can be difficult to implement outside of research settings.^{3,4}

There are caveats, however. As noted by Mead *et al*, only a small set of heterogeneous studies have examined the effects of befriending interventions on depressive symptoms. Furthermore, funnel plot asymmetry suggested that publication bias may have influenced the estimate of the degree to which befriending may affect depressive symptoms. The authors did not assess the degree to which publication bias may have influenced the results of the meta-analysis. However, if only studies with statistical power of at least 0.70 among the studies with short-term outcomes evaluated by Mead *et al* are analysed, the resulting synthesised effect estimate is 0.08 (95% CI -0.06 to 0.21, four studies), a substantially smaller estimate than that produced by all nine studies (0.27, 95% CI 0.06 to 0.48). Thus, as noted by Mead *et al*, more high-quality research is needed on befriending in order to determine the likely benefit to patients in clinical practice.

Meanwhile, the results of the meta-analysis suggest that future research on collaborative care should use a befriending or attention control group. Up to now, collaborative care interventions have been compared with usual care, and it is not known to what degree the effects that have been reported are due to specific effects of the collaborative care intervention versus effects that may come from the substantially increased attention and support received by patients in collaborative care.

- 1 Mead N, Lester H, Chew-Graham C, Gask L, Bower P. Effects of befriending on depressive symptoms and distress: systematic review and meta-analysis. *Br J Psychiatry* 2010; **196**: 96–101.
- 2 Gilbody S, Bower P, Fletcher J, Richards D, Sutton AJ. Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. *Arch Intern Med* 2006; **166**: 2314–21.
- 3 Katon WJ, Seelig M. Population-based care of depression: team care approaches to improving outcomes. *J Occup Environ Med* 2008; **50**: 459–67.
- 4 Katon W, Unützer J, Wells K, Jones L. Collaborative depression care: history, evolution and ways to enhance dissemination and sustainability. *Gen Hosp Psychiatry* 2010; in press.

Ghassan El-Baalbaki, Department of Psychiatry, McGill University, and Department of Psychiatry, Jewish General Hospital, 4333 Côte Ste Catherine Road, Montreal, Quebec H3T 1E4, Canada. Email: ghassan.elbaalbaki@videotron.ca; **Erin Arthurs**, **Brooke Levis**, Department of Psychiatry, McGill University, and Jewish General Hospital, Montreal; **Brett D. Thombs**, Departments of Psychiatry, Epidemiology & Biostatistics, Occupational Health, and Medicine (Division of Rheumatology), McGill University, and Department of Psychiatry, Centre for Clinical Epidemiology and Community Studies, and Division of Rheumatology, Jewish General Hospital, Montreal, Quebec, Canada

doi: 10.1192/bjp.197.3.247

Authors' reply: We thank Dr El-Baalbaki and colleagues for their thoughtful comments on our recent paper. We agree with their cautious interpretation of the results of our meta-analysis.

We were interested in their suggestion that befriending serve as a comparator condition for more structured treatments such as collaborative care, to tease out the specific benefits of the latter over and above the general effects of increased attention and support, and to explore the cost-effectiveness of these complex organisational interventions. Although this makes good sense in design terms, it does, however, relegate befriending to the status of comparator rather than active intervention. The recent Mental Health Foundation report *The Lonely Society?* (www.mentalhealth.org.uk/campaigns/loneliness-and-mental-health/) highlights the impact of loneliness on health, and its findings are supported by the Royal College of Psychiatrists.¹ We would therefore want to complement Dr El-Baalbaki and colleagues' suggestion with further research specifically exploring the role of befriending as a potential alternative therapeutic intervention for certain groups such as isolated older adults.

- 1 Royal College of Psychiatrists. RCPsych comments on MHF report 'The Lonely Society?' Royal College of Psychiatrists, May 2010 (<http://www.rcpsych.ac.uk/press/pressreleases2010/thelonelysociety.aspx>).

Nicola Mead, NIHR School for Primary Care Research, 5th Floor, Williamson Building, University of Manchester, Oxford Road, Manchester M13 9PL, UK. Email: nicki.mead@manchester.ac.uk. **Helen Lester**, **Carolyn Chew-Graham**, **Linda Gask**, NIHR School for Primary Care Research, University of Manchester; **Peter Bower**, National Primary Care Research and Development Centre, University of Manchester, UK.

doi: 10.1192/bjp.197.3.247a

Correction

Help-seeking and receipt of treatment among UK service personnel. *BJP*, 197, 149–155. Authors' affiliations/correspondence details should read:

Amy C. Iversen, MRCPsych, Lauren van Staden, MA, King's Centre for Military Health Research, King's College London; Jamie Hacker Hughes, PsychD, Academic Centre

for Defence Mental Health, King's College London; Tess Browne, BSc, King's Centre for Military Health Research, King's College London; Neil Greenberg, MD, Academic Centre for Defence Mental Health, King's College London; Matthew Hotopf, PhD, Roberto J. Rona, FFPH, Simon Wessely, FMedSci, King's Centre for Military Health Research, King's College London; Graham Thornicroft, PhD, Health Service