

ON MODULES OF FINITE PROJECTIVE DIMENSION

S. P. DUTTA

Abstract. We address two aspects of finitely generated modules of finite projective dimension over local rings and their connection in between: embeddability and grade of order ideals of minimal generators of syzygies. We provide a solution of the embeddability problem and prove important reductions and special cases of the order ideal conjecture. In particular, we derive that, in any local ring R of mixed characteristic $p > 0$, where p is a nonzero divisor, if I is an ideal of finite projective dimension over R and $p \in I$ or p is a nonzero divisor on R/I , then every minimal generator of I is a nonzero divisor. Hence, if P is a prime ideal of finite projective dimension in a local ring R , then every minimal generator of P is a nonzero divisor in R .

In this note we consider two aspects of finitely generated modules of finite projective dimension over any local ring: embeddability and grade of order ideals of minimal generators of its syzygies (minimal). In regard to embeddability, Auslander and Buchweitz [1] proved that any finitely generated module on a Gorenstein local ring can be embedded in a module of finite projective dimension such that the cokernel is Cohen–Macaulay. This result has several applications in solving homological questions and conjectures in commutative algebra, including Serre’s χ_i -conjectures and their generalizations on intersection multiplicity (see [5]). For all these conjectures, one is usually concerned with finitely generated modules M which are of finite projective dimension over a local ring R but are not so over R/xR , x being a nonzero divisor in the annihilator of M in R . However, a similar result has been absent for non-Gorenstein local rings until now. In this note we prove the following with respect to embeddability for modules of finite projective dimension.

THEOREM 1. *Let (R, m) be a local ring, and let M be a finitely generated module of finite projective dimension with positive grade over R .*

Received March 4, 2013. Revised March 5, 2014. Accepted August 13, 2014.

2010 Mathematics Subject Classification. Primary 13D02, 13D22; Secondary 13C15, 13D25, 13H05.

© 2015 by The Editorial Board of the *Nagoya Mathematical Journal*

Let $\mathbf{x} = \{x_1, \dots, x_t\}$ be any R -sequence contained in the annihilator of M (henceforth $\text{ann}_R M$), and let $\overline{R} = R/(x_1, \dots, x_t)$. Then there exists a short exact sequence of finitely generated \overline{R} -modules

$$0 \rightarrow M \rightarrow Q \rightarrow T \rightarrow 0,$$

where $\text{pd}_{\overline{R}} Q < \infty$ and $\text{pd}_R T = t$.

From the construction of Q , it would follow that support of Q over $\overline{R} = \text{Spec}(\overline{R})$ and Q satisfies the strong intersection conjecture due to Peskine and Szpiro [15]. Since T is perfect, T also possesses the above property. The effect of this theorem on generalizations of Serre’s conjectures on intersection multiplicities for arbitrary local rings, in particular for Cohen–Macaulay rings, will be the subject of a future paper. This paper focuses on its effect on the order ideal conjecture, which is our next topic.

The order ideal conjecture stems from Evans and Griffith’s work [7] on grade of order ideals of minimal generators of syzygies in equicharacteristic. The statement of the conjecture is the following.

ORDER IDEAL CONJECTURE. *Let (R, m) be a local ring. Let M be a finitely generated module of finite projective dimension over R , and let S_i denote its i th syzygy for $i > 0$. If β is a minimal generator of S_i , then the order ideal $\mathcal{O}_{S_i}(\beta)$ has grade at least i .*

Let us recall that $\mathcal{O}_{S_i}(\beta) = \{f(\beta) \mid f \in \text{Hom}_R(S_i, R)\}$.

We say that a module M satisfies the order ideal conjecture if order ideals of minimal generators of all its syzygies satisfy the respective grade inequalities mentioned above.

Evans and Griffith [7], [9] proved the above conjecture for equicharacteristic local rings in order to solve the syzygy problem over the above class of rings. The existence of big Cohen–Macaulay modules, due to Hochster [12], played an important role in their proof. Later in [10] they proved a graded version of the above conjecture for a certain class of graded rings in mixed characteristic. We also refer the reader to [3, Theorem 9.5.2] for a more general version of the order ideal theorem in the equicharacteristic case. Actually, for their proof of the syzygy theorem in equicharacteristic, Evans and Griffith needed to prove the above conjecture only for modules which are locally free on the punctured spectrum of $\text{spec}(R)$, R being regular local. And they reduced the proof of this case to what is now known as the *improved new intersection conjecture*. Later Hochster [13] showed

that the canonical element conjecture implies the improved new intersection conjecture. The equivalence of these two conjectures was established in [4]. In our most recent work [6], we have shown that *a particular case of the order ideal conjecture implies the monomial conjecture* and therefore also implies all its equivalent forms—for example, the direct summand conjecture, the canonical element conjecture, and the improved new intersection conjecture. Thus, the order ideal conjecture now occupies a central position among several homological conjectures in commutative algebra.

First, let us mention that in order to prove the order ideal conjecture on arbitrary local rings R , *it is enough to concentrate on first syzygies* of modules of finite projective dimension (see Lemma 2.1). Theorem 2.3 shows that *for the validity of the order ideal conjecture it is enough to prove that every minimal generator of ideals of height 2, grade 2, and of finite projective dimension over R is a nonzero divisor*. In Theorem 2.5, we prove the following: *Given a module of finite projective dimension M on a local ring R , there exists an R -sequence x_1, \dots, x_h , $h = \text{pd}_R M$, such that for any R -module N for which x_1, \dots, x_h form an N -sequence, $\text{Tor}_j^R(M, N) = 0$ for $j > 0$.*

This theorem leads us to the following (see Corollary 2.7).

Let (R, m) be a local ring, and let M be a finitely generated module of finite projective dimension over R . Let x_1, \dots, x_h be an R -sequence as mentioned above. If R has mixed characteristic $p > 0$, we assume that p, x_1, \dots, x_h form a part of a system of parameters of R . Then for every minimal generator β of $S_1 = \text{Syz}_R^1(M)$, the grade of $\mathcal{O}_{S_1}(\beta) \geq 1$.

The statement of our main theorem in Section 2 is the following.

THEOREM 2.11. *Let (R, m) be a local ring of mixed characteristic $p > 0$. Let M be a finitely generated module of finite projective dimension over R , and let β be a minimal generator of S_i , the i th syzygy (minimal) of M , for $i > 0$. We assume that either p is a nonzero divisor in R or p is nilpotent. We have the following.*

- (a) *If p is nilpotent, then the order ideal conjecture is valid on R .*
- (b) *If $pM = 0$, then the grade of $\mathcal{O}_{S_1}(\beta) \geq 1$.*
- (c) *Suppose that balanced or complete almost Cohen–Macaulay algebras exist over complete local domains. If $pM = 0$ and $p \in m - m^2$, then the grade of $\mathcal{O}_{S_i}(\beta) \geq i$, $\forall i \geq 1$.*

- (d) Assume that every element in $m - m^2$ is a nonzero divisor and that the order ideal conjecture is valid over R/xR for $x \in m - m^2$. If $\text{ann}_R M \cap (m - m^2) \neq \phi$ or $\text{depth}_R M > 0$, then $\text{grade } \mathcal{O}_{S_i}(\beta) \geq i, \forall i \geq 1$.

As a consequence of the above theorem, we have the following corollaries.

COROLLARY 2.12. *For any ideal I of finite projective dimension over R of mixed characteristic $p > 0$, where p is a nonzero divisor in R , if $p \in I$ or p is a nonzero divisor on R/I , then every minimal generator of I is a nonzero divisor. In particular, if P is a prime ideal of finite projective dimension over R , then every minimal generator of P is a nonzero divisor in R .*

COROLLARY 2.13. *Let (R, m) be a regular local ring of dimension n , and assume that the order ideal conjecture is valid for regular local rings of dimension $(n - 1)$. If M is a finitely generated R -module such that either M is annihilated by a regular parameter or $\text{depth}_R M > 0$, then M satisfies the order ideal conjecture.*

The two main ingredients of our proof of Theorem 2.11 are Theorem 1.2 and Shimomoto’s theorem (Theorem 2.10) on the existence of almost Cohen–Macaulay algebras (see [17, Theorem 5.3]).

Throughout this note, *local* means Noetherian local, $\text{pd}_R M$ denotes projective dimension of M over R , and $\text{Syz}_R^i(M)$ denotes the i th syzygy in a minimal free resolution of M . (For definitions of standard notions and their basic properties, see [3].)

§1.

First we would like to mention the following proposition.

PROPOSITION 1.1 ([4, Proposition 1.1]). *Let A be a local ring, and let $F_\bullet : \rightarrow A^{s_i} \rightarrow A^{s_{i-1}} \rightarrow \dots \rightarrow A^{s_0} \rightarrow 0$ be a free complex with $H_0(F_\bullet) = M$. Let $N \subset M$ be a submodule of M . Then we can construct a free complex $G_\bullet : \rightarrow A^{d_i} \rightarrow A^{d_{i-1}} \rightarrow \dots \rightarrow A^{d_0} \rightarrow 0$ with $H_0(G) = N$ and a map $\phi_\bullet : G_\bullet \rightarrow F_\bullet$ such that ϕ_0 induces the inclusion $N \hookrightarrow M$ and such that the mapping cone of ϕ_\bullet is a free resolution of M/N ; that is, ϕ_\bullet induces an isomorphism $H_i(G_\bullet) \xrightarrow{\sim} H_i(F_\bullet)$ for $i > 0$. Moreover, by our construction, G_\bullet is minimal.*

For a proof we refer the reader to [4, Proposition 1.1].

Next we prove the main theorem of this section.

THEOREM 1.2. *Let (R, m) be a local ring, and let M be a finitely generated module of finite projective dimension over R . Assume that $\text{grade}_R M > 0$. Let \mathfrak{x} denote the ideal generated by an R -sequence of length t contained in $\text{ann}_R M$, and let $\overline{R} = R/\mathfrak{x}R$. Then there exists a short exact sequence of finitely generated \overline{R} -modules*

$$0 \rightarrow M \rightarrow Q \rightarrow T \rightarrow 0$$

such that $\text{pd}_{\overline{R}} Q < \infty$ and $\text{pd}_R T = t$.

Proof. First suppose that $\text{pd}_R M = 1$. Let us recall that $\text{grade}_R M \leq \text{pd}_R M$. Since $\text{grade}_R M > 0$, it follows that $\text{grade}_R M = 1$. Let $x \in \text{ann}_R M$ be a nonzero divisor. Consider a minimal free resolution: $0 \rightarrow R^{t_1} \xrightarrow{f} R^{t_0} \rightarrow M \rightarrow 0$ of M over R . Tensoring this resolution with R/xR , we obtain an exact sequence: $0 \rightarrow \text{Tor}_1^R(M, R/xR) \rightarrow (R/xR)^{t_1} \xrightarrow{\overline{f}} (R/xR)^{t_0} \rightarrow M \rightarrow 0$. Since $x \in \text{ann}_R M$, $\text{Tor}_1^R(M, R/xR) \simeq M$. Let $T = \text{im } \overline{f}$. Hence, we obtain the short exact sequence $0 \rightarrow M \rightarrow (R/xR)^{t_1} \rightarrow T \rightarrow 0$, and our assertion follows. So we can assume that $\text{pd}_R M \geq 2$.

Let $(F_\bullet, d_\bullet) : 0 \rightarrow R^{r_n} \xrightarrow{d_n} R^{r_{n-1}} \rightarrow \dots \rightarrow R^{r_1} \xrightarrow{d_1} R^{r_0} \rightarrow 0$ be a minimal projective resolution of M over R , and let $(L_\bullet, \phi_\bullet) : \overline{R}^{s_n} \xrightarrow{\phi_n} \overline{R}^{s_{n-1}} \rightarrow \dots \rightarrow \overline{R}^{s_1} \xrightarrow{\phi_0} \overline{R}^{s_0} \rightarrow 0$ be a minimal projective resolution of M over \overline{R} . Since $\text{pd}_R M = n$, $\text{grade Ext}_R^i(M, R) \geq i$ for $1 \leq i \leq n$, $\text{Ext}_R^n(M, R) \neq 0$, and $\text{Ext}_R^i(M, R) = 0$ for $i > n$. Hence, $\text{grade Ext}_{\overline{R}}^i(M, \overline{R}) \geq i$ for $1 \leq i \leq n - t$, $\text{Ext}_{\overline{R}}^{n-t}(M, \overline{R}) \neq 0$ ($\simeq \text{Ext}_R^n(M, R)$), and $\text{Ext}_{\overline{R}}^i(M, \overline{R}) = 0$ for $i > (n - t)$.

Applying $\text{Hom}_{\overline{R}}(-, \overline{R})$ to L_\bullet , we obtain the following free complex L_\bullet^* :

$$L_\bullet^* : 0 \rightarrow \overline{R}^{s_0^*} \rightarrow \overline{R}^{s_1^*} \rightarrow \dots \rightarrow \overline{R}^{s_{n-t}^*} \rightarrow 0.$$

(For any $R(\overline{R})$ module V , $V^* = \text{Hom}_R(M, R)(\text{Hom}_{\overline{R}}(V, \overline{R}))$.)

Let $G = \text{Coker } \phi_{n-t}^*$. We have a short exact sequence

$$(1) \quad 0 \rightarrow \text{Ext}_{\overline{R}}^{n-t}(M, \overline{R}) \xrightarrow{c} G \xrightarrow{\eta} \text{Im } \phi_{n-t+1}^* \rightarrow 0.$$

By the above proposition, there exists a minimal free complex $(P_\bullet, \alpha_\bullet) : \overline{R}^{b_{n-t}} \xrightarrow{\alpha_{n-t}} \overline{R}^{b_{n-t-1}} \rightarrow \dots \xrightarrow{\alpha_1} \overline{R}^{b_0} \rightarrow 0$ with $H_0(P_\bullet) = \text{Ext}_{\overline{R}}^{n-t}(M, \overline{R})$ and a map $\Psi_\bullet : P_\bullet \rightarrow L_\bullet^*$ such that Ψ_\bullet induces the injection c in (1) and the mapping cone of Ψ_\bullet is a free resolution of $\text{Im } \phi_{n-t+1}^*$. We have the following

commutative diagram:

$$\begin{array}{ccccccccccc}
 (2) & P_\bullet : & \longrightarrow & \overline{R}^{b_{n-t+1}} & \xrightarrow{\alpha_{n-t+1}} & \overline{R}^{b_{n-t}} & \longrightarrow & \overline{R}^{b_{n-t-1}} & \cdots & \longrightarrow & \overline{R}^{b_1} & \longrightarrow & \overline{R}^{b_0} & \longrightarrow & 0 \\
 & & & & & \downarrow \Psi_{n-t} & & \downarrow \Psi_{n-t-1} & & & \downarrow \Psi_1 & & \downarrow \Psi_0 & & & \\
 & L_\bullet^* : & 0 & \longrightarrow & \overline{R}^{s_0^*} & \longrightarrow & \overline{R}^{s_1^*} & \cdots & \longrightarrow & \overline{R}^{s_{n-t-1}^*} & \xrightarrow{\phi_{n-t}^*} & \overline{R}^{s_{n-t}^*} & \longrightarrow & 0
 \end{array}$$

Applying $\text{Hom}(-, \overline{R})$ to (2), we obtain the following commutative diagram of exact complexes:

$$\begin{array}{ccccccccccc}
 (3) & & \longrightarrow & \overline{R}^{s_{n-t}} & \longrightarrow & \overline{R}^{s_{n-t-1}} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{s_1} & \longrightarrow & \overline{R}^{s_0} & \longrightarrow & M & \longrightarrow & 0 \\
 & & & \downarrow \Psi_0^* & & \downarrow \Psi_1^* & & & & \downarrow \Psi_{n-t-1}^* & & \downarrow \Psi_{n-t}^* & & \downarrow \Psi & & \\
 & 0 & \longrightarrow & \overline{R}^{b_0^*} & \longrightarrow & \overline{R}^{b_1^*} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{b_{n-t-1}^*} & \xrightarrow{\alpha_{n-t}^*} & \overline{R}^{b_{n-t}^*} & \longrightarrow & Q & \longrightarrow & 0
 \end{array}$$

where $Q = \text{Coker } \alpha_{n-t}^*$ and $\Psi : M \rightarrow Q$ is induced by Ψ_{n-t}^* . Since $\text{grade Ext}_{\overline{R}}^i(M, \overline{R}) \geq i, 1 \leq i \leq n-t$, the bottom row of (3) provides a minimal free resolution of Q , and hence $\text{pd}_{\overline{R}} Q = n-t$. Moreover, by construction, $\text{Ext}_{\overline{R}}^i(Q, \overline{R}) \xrightarrow{\sim} \text{Ext}_{\overline{R}}^i(M, \overline{R})$ for $i > 0$.

Next we want to prove that $\Psi : M \rightarrow Q$ in (3) is injective.

Let $\overline{F}_\bullet = F_\bullet \otimes \overline{R}, S_i = \text{Syz}_R^i(M)$, and let $\overline{S}_i = S_i/\mathbf{x}S_i$, for $i > 0$. Since \mathbf{x} is generated by an R -sequence of length t in $\text{ann}_R M$, we have $\text{Tor}_t^R(M, \overline{R}) \xleftarrow{\sim} M$ and $\text{Tor}_i^R(M, \overline{R}) = 0$ for $i > t$. Tensoring the exact sequence $0 \rightarrow S_t \rightarrow R^{t-1} \rightarrow S_{t-1} \rightarrow 0$ by \overline{R} , we obtain an exact sequence

$$0 \rightarrow M \xrightarrow{j} \overline{S}_t \rightarrow \overline{R}^{t-1} \rightarrow \overline{S}_{t-1} \rightarrow 0.$$

Let $\theta_\bullet : L_\bullet \rightarrow \overline{F}_\bullet$ be a lift of $j : M \hookrightarrow \overline{S}_t$. We have the following commutative diagram:

$$\begin{array}{ccccccccccc}
 (4) & L_\bullet : & \longrightarrow & \overline{R}^{s_{n-t}} & \longrightarrow & \overline{R}^{s_{n-t-1}} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{s_0} & \longrightarrow & 0 \\
 & & & \downarrow \theta_{n-t} & & \downarrow \theta_{n-t-1} & & & & \downarrow \theta_0 & & & & & & \\
 & \overline{F}_\bullet : & 0 & \longrightarrow & \overline{R}^{r_n} & \longrightarrow & \overline{R}^{r_{n-1}} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{r_t} & \longrightarrow & \overline{R}^{r_{t-1}} & \longrightarrow & \cdots
 \end{array}$$

The mapping cone of $\{\theta_\bullet\}$ in (4) is a free resolution of $\text{Im } \overline{d}_t$ over \overline{R} , and θ_0 induces the isomorphism $\tilde{\theta}_0 : M \xrightarrow{\sim} \text{Tor}_t^R(M, \overline{R})$ via $j : M \hookrightarrow \overline{S}_t$.

(8)

$$\begin{array}{ccccccc}
 \overline{F}_\bullet^* & : & \cdots & \longrightarrow & \overline{R}^{r_j^*} & \xrightarrow{\overline{d}_{j+1}^*} & \overline{R}^{r_{j+1}^*} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{r_{n-1}^*} & \longrightarrow & \overline{R}^{r_n^*} & \longrightarrow & 0 \\
 & & & & \swarrow^{h_j} & \downarrow^{\theta_{j-1}^*} & \searrow^{h_{j+1}} & & & & \downarrow^{h_n} & & & & & \\
 L_\bullet^* & : & \cdots & \longrightarrow & \overline{R}^{s_{j-2}^*} & \longrightarrow & \overline{R}^{s_{j-1}^*} & \longrightarrow & \overline{R}^{s_j^*} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{s_{n-t-1}^*} & \longrightarrow & \overline{R}^{s_{n-1}^*} & \longrightarrow & 0 \\
 & & & & \downarrow^{\eta_{j-2}} & & \downarrow^{\eta_{j-1}} & & & & \downarrow & & & & \downarrow & & \\
 H_\bullet & : & \cdots & \longrightarrow & \overline{R}^{b_{n-j}} \oplus \overline{R}^{s_{j-2}^*} & \xrightarrow{\delta_{n-j}} & \overline{R}^{b_{n-j-1}} \oplus \overline{R}^{s_{j-1}^*} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{b_0} \oplus \overline{R}^{s_{n-t-1}^*} & \longrightarrow & \overline{R}^{s_{n-1}^*} & & & \\
 & & & & \downarrow^{\gamma_{n-j}} & & \downarrow^{\gamma_{n-j-1}} & & & & \downarrow & & & & & & \\
 P_\bullet(-1) & : & \cdots & \longrightarrow & \overline{R}^{b_{n-j}} & \xrightarrow{\alpha_{n-j}} & \overline{R}^{b_{n-j-1}} & \longrightarrow & \cdots & \longrightarrow & \overline{R}^{b_0} & \longrightarrow & 0 & & & \\
 & & & & & & \delta_{n-j} = (\alpha_{n-j}, (-1)^{n-j-1}\Psi_{n-j} + \phi_{j-1}^*) & & & & & & & & & &
 \end{array}$$

$$\begin{array}{ccccccc}
 \overline{F}_\bullet^* & : & \cdots & \longrightarrow & \overline{R}^{r_j^*} & \longrightarrow & \overline{R}^{r_{j+1}^*} \\
 & & & & \swarrow^{k_j} & \downarrow^{\beta_j} & \searrow^{k_{j+1}} \\
 P_\bullet & : & \cdots & \longrightarrow & \overline{R}^{b_{n-j}} & \longrightarrow & \overline{R}^{b_{n-j-1}} \\
 & & & & \downarrow^{\theta_{j-1}^*} & \downarrow^{\Psi_{n-j}} & \downarrow^{\theta_j^*} \\
 L_\bullet^* & : & \longrightarrow & \overline{R}^{s_{j-2}^*} & \longrightarrow & \overline{R}^{s_{j-1}^*} & \longrightarrow & \overline{R}^{s_j^*}
 \end{array}$$

Hence, $\{\theta_\bullet^*\}$ and $\{\Psi_\bullet\beta'_\bullet\}$ define identical maps (modulo \pm sign) on homologies of F_\bullet^* and L_\bullet and, consequently, for the homologies of their respective duals. If $\tilde{\Psi} : M \rightarrow H^{n-t}(P_\bullet^*)$, $\tilde{\beta} : H^{n-t}(P_\bullet^*) \rightarrow \text{Tor}_t^R(M, \overline{R})$, and $\tilde{\theta}_0 : M \xrightarrow{\sim} \text{Tor}_t^R(M, \overline{R})$ denote the maps induced by Ψ_{n-t}^* , β_t^* , and $(\theta_0^*)^* = \theta_0$, respectively, then $\tilde{\theta}_0 = \tilde{\beta} \cdot \tilde{\Psi}$ (modulo a sign). Since $\tilde{\theta}_0$ is an isomorphism, $\Psi = (\text{inclusion of } H^{n-t}(P_\bullet^*) \hookrightarrow Q) \cdot \tilde{\Psi} : M \hookrightarrow Q$ is injective.

Let $\text{Coker } \psi = T$; then $0 \rightarrow M \xrightarrow{\Psi} Q \rightarrow T \rightarrow 0$ is exact. Since the mapping cones of $\Psi_\bullet^*, \Psi_\bullet$ are free resolutions of T , $\text{Im } \phi_{n-t+1}^*$, respectively, and $\text{Ext}_R^i(M, \overline{R}) \xrightarrow{\sim} \text{Ext}_R^i(Q, \overline{R})$ for $i > 0$, it follows that $\text{Ext}_R^i(T, \overline{R}) = 0$ for $i > 0$; that is, $\text{Ext}_R^i(T, R) = 0$ for $i > t$. Since $\text{pd}_R M < \infty$, we have $\text{pd}_R T < \infty$, and hence $\text{pd}_R T = t$. This completes our proof. \square

COROLLARY 1.3. *Let R be a local ring, and let M be a finitely generated module of finite projective dimension with $\text{pd}_R M > 1$ and $\text{grade } R > 0$. Given any nonzero divisor $x \in \text{ann}_R M$, M can be embedded in a finitely generated module Q of finite projective dimension over R/xR in such a*

way that, if $(G_\bullet, \gamma_\bullet), (F_\bullet, d_\bullet)$ are minimal free resolutions of M and Q over R , respectively, and $\phi_\bullet : G_\bullet \rightarrow F_\bullet$ is a lift of $i : M \hookrightarrow Q$, then ϕ_\bullet induces an isomorphism between $(G_\bullet, \gamma_\bullet)_{i \geq 2}$ and $(F_\bullet, d_\bullet)_{i \geq 2}$, $\phi_1(G_1)$ is either a summand of or isomorphic to F_1 , and ϕ_0 is an injection. Moreover, $\text{syz}_R^1(M) \oplus R^t \simeq \text{syz}_R^1(Q)$ for some $t \geq 0$.

Proof. By Theorem 1.2 we have an exact sequence of R/xR modules

$$0 \rightarrow M \xrightarrow{i} Q \xrightarrow{\eta} T \rightarrow 0,$$

where $\text{pd}_{R/xR} Q < \infty$ and $\text{pd}_R T = 1$.

The proof of the corollary now follows directly either by constructing a minimal free resolution of Q from minimal free resolutions of M and T or by extracting a minimal free resolution of M from the mapping cone of $\eta_\bullet : F_\bullet \rightarrow L_\bullet$, where F_\bullet, L_\bullet are minimal free resolutions of Q and T , respectively. (For details of such basic constructions, the reader is referred to [3].) □

REMARK. Since $\text{grade}_R T = \text{pd}_R T = r$, T is perfect. It can be easily checked from the above construction that $\text{grade}_{\overline{R}} Q = 0$, and since $\text{pd}_{\overline{R}} Q < \infty$, $\text{support}_{\overline{R}} Q = \text{Spec}(\overline{R})$. Moreover, the strong intersection conjecture (see [12], [15]) is valid for both Q and T . For details on this observation we refer the reader to [15, Section 4, Chapitre II].

§2.

LEMMA 2.1. *Let (R, m) be a local ring of dimension of n . Assume that the order ideal conjecture is valid for local rings of dimension $(n - 1)$. Then, for the validity of the order ideal conjecture on R , it is enough to prove the validity of the assertion for the first syzygies of modules of finite projection. In particular, for cyclic modules of finite projective dimension over R , it is enough to prove that every minimal generator of any ideal in R of finite projective dimension over R is a nonzero divisor in R .*

Proof. Let (F_\bullet, d_\bullet) be a minimal free resolution of M where $F_i = R^{r_i}$ for $i \geq 0$. Let S_i denote the i th syzygy of M for $i \geq 1$, and let β be a minimal generator of S_i for $i > 1$. Then $\beta = d_i(e)$ for some free generator e of F_i , and we have $\beta = \begin{pmatrix} a_1 \\ \vdots \\ a_{r_{i-1}} \end{pmatrix} \in R^{r_{i-1}}$. Let J denote the ideal generated by $a_1, \dots, a_{r_{i-1}}$. Let x be a nonzero divisor on R ; then x is a nonzero divisor on

S_1 . Let $\overline{R} = R/xR$, and let $\overline{S}_i = S_i/xS_i$, $i \geq 1$. We have that \overline{S}_i is of finite projective dimension over \overline{R} , and $\overline{S}_i = \text{Syz}_{\overline{R}}^{i-1}(\overline{S}_1)$ for $i > 1$. By induction hypothesis, $\text{grade}_{\overline{R}}(J + xR)/xR \geq (i - 1)$; this implies that $\text{grade}_R J \geq (i - 1)$. Let $y \in J$ be a nonzero divisor on R . Let $\tilde{R} = R/yR$, and let $\tilde{S}_i = S_i/yS_i$ for $i \geq 1$. Then, again by arguing as above, $\text{grade}_{\tilde{R}} J/(y) \geq (i - 1)$. Hence, $\text{grade}_R J \geq i$. The second assertion now follows readily. \square

LEMMA 2.2. *Let M be a finitely generated module of finite projective dimension over a local ring (R, m) . Suppose that $\text{rank}_R M = s$. Then there exists a free submodule $F = R^s$, generated by a part of a minimal set of generators of M , such that M/F has positive grade.*

Proof. We induct on s . Since $\text{pd}_R M$ is finite, if $s = 0$, then for every associated prime P of R , $M_P = 0$, and hence $\text{grade}_R M > 0$. Now suppose that $s > 0$. Since $\text{pd}_R M$ is finite, by basic element method (see [8, Lemma 2.1]) there exists a minimal generator α of M such that the image of α is a part of a basis of M_P for every associate prime P of R . Hence, we have a short exact sequence

$$0 \rightarrow R \rightarrow M \xrightarrow{\phi} M' \rightarrow 0 \quad (1 \rightarrow \alpha).$$

Then $\text{pd}_R M' < \infty$ and $\text{rank}_R M' = \text{rank}_R M - 1 = s - 1$. By induction, there exists a free submodule $F' = R^{s-1}$ of M' , generated by a part of a minimal set of generators of M' , such that $M'' = M'/F'$ has positive grade. We have the following short exact sequence:

$$0 \rightarrow F' \xrightarrow{i} M' \xrightarrow{\psi} M'' \rightarrow 0.$$

Since F' is free, we can lift $i : F' \rightarrow M'$ to $\eta : F' \rightarrow M$ such that $\phi \cdot \eta = i$. Let $\theta = \psi \cdot \phi$. It can be easily checked that θ is surjective and that $\text{Ker } \theta = R \oplus F'$. Hence the lemma follows. \square

Our next theorem reduces the order ideal conjecture to the assertion that every minimal generator of a certain class of ideals of finite projective dimension must be a nonzero divisor.

THEOREM 2.3. *Let (R, m) be a local ring of dimension n . Assume that the order ideal conjecture is valid for local rings of dimension $(n - 1)$. Then, for the validity of the order ideal conjecture over R , it is enough to prove that every minimal generator of any ideal of grade 2, height 2, and of finite projective dimension over R is a nonzero divisor in R .*

Proof. Let M be a finitely generated module of finite projective dimension over R . Due to Lemma 2.1, for the validity of the order ideal conjecture it is enough to consider minimal generators of $S = \text{Syz}_R^1(M)$. If $\text{grade}_R M = 0$, then $\text{ann}_R M = 0$. Let $s_0 = \text{rank}_R M$. By Lemma 2.2, there exists a free submodule $F = R^{s_0}$ generated by a part of a minimal set of generators of M such that $M' = M/F$ has positive grade. From the commutative diagram

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 & & & R^{s_0} & & R^{s_0} & \\
 & & & \downarrow & & \downarrow & \\
 (1) & 0 & \rightarrow & S & \rightarrow & R^{r_0} & \xrightarrow{\eta} & M & \rightarrow & 0 \\
 & & & \downarrow & & \downarrow & & \downarrow & & \\
 & 0 & \rightarrow & S & \rightarrow & R^{r_0-s_0} & \rightarrow & M' & \rightarrow & 0 \\
 & & & \downarrow & & \downarrow & & \downarrow & & \\
 & & & 0 & & 0 & & & &
 \end{array}$$

it follows that, without any loss of generality, we can assume that $\text{grade}_R M > 0$. Let x be a nonzero divisor in $\text{ann}_R M$. By Corollary 1.3 we can assume that M has finite projective dimension over R/xR and that $\text{support}(M) = \text{support}(R/xR)$. Let $\overline{R} = R/xR$, and let $\overline{S} = S/xS$. By tensoring $0 \rightarrow S \rightarrow R^{r_0} \rightarrow M \rightarrow 0$ with \overline{R} , we obtain the short exact sequences

$$(2) \quad 0 \rightarrow M \xrightarrow{\theta} \overline{S} \rightarrow T \rightarrow 0, \quad 0 \rightarrow T \rightarrow \overline{R}^{r_0} \rightarrow M \rightarrow 0,$$

where $T = \text{Syz}_{\overline{R}}^1(M)$. If γ is a minimal generator of M , and if $e \in R^{r_0}$, a free generator of R^{r_0} , is such that $\eta(e) = \gamma$, then $\theta(\gamma) = \text{Im}(xe)$ in \overline{S} . Let $\alpha_i \in S$ denote the lifts of a minimal set of generators $\overline{\alpha}_i$, $1 \leq i \leq h$, of T . By induction, $\text{grade}_{\mathcal{O}_{T,\overline{R}}}(\overline{\alpha}_i) \geq 1$; hence, $\text{grade}_{\mathcal{O}_S}(\alpha_i) \geq 1$ for $1 \leq i \leq h$. Due to the exact sequences in (2) we obtain a minimal set of generators $xe_1, \dots, xe_a, \alpha_1, \dots, \alpha_h$ of S where e_1, \dots, e_a form a part of a basis of R^{r_0} . If (F_\bullet, d_\bullet) ($F_i = R^{r_i}$) denote a minimal free resolution of M over R , then there exist $\tilde{e}_1, \dots, \tilde{e}_a, \dots, \tilde{e}_{a+j}, \dots$ a basis of R^{r_1} such that $d_1(\tilde{e}_i) = xe_i$, $1 \leq i \leq a$, and $d_1(\tilde{e}_{a+j}) = \alpha_j$, $1 \leq j \leq h$. Any minimal generator of S is of the form $\sum_{i=1}^a c_i xe_i + \sum_{j=1}^h d_j \alpha_j$, where at least one of c_i 's or d_j 's is a unit. If any d_j in the above expression is a unit, then we are done by induction. Thus, it is easy to check that, in order to show that for any minimal generator β of S , $\text{grade}_{\mathcal{O}_S}(\beta) \geq 1$, it is enough to consider $\beta = xe - \sum \lambda_i \alpha_i$, where $e \in \{e_1, \dots, e_a\}$, $\sum \lambda_i \overline{\alpha}_i \neq 0$ in T . Due to the second exact sequence in (2),

if any λ_i is a unit, then we are done by induction. Hence, we can assume that all $\lambda_i \in m$ in the above expression of β . Let $t = \text{rank}_{\overline{R}} M$. By arguing as in (1) we obtain the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \rightarrow & R^t & \xrightarrow{x} & R^t & \rightarrow & \overline{R}^t \\
 & & \downarrow \psi & & \downarrow \phi_0 & & \downarrow \phi \\
 (3) & & 0 & \rightarrow & R^{r_0} & \rightarrow & M \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \rightarrow & S' & \rightarrow & R^{r_0-t} & \rightarrow & M' \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Let $\{\overline{e}_j\}, 1 \leq j \leq t$, denote a basis of \overline{R}^t such that $\gamma_j = \phi(\overline{e}_j), 1 \leq j \leq t$, in (3) is a part of a minimal set of generators of M (see Lemma 2.2). By commutativity of (3), $\psi(e_j) = xe_j, 1 \leq j \leq t$. Note that none of xe_j may be a part of $\{xe_1, \dots, xe_a\}$ —part of a minimal basis of S mentioned above. We want to prove the following.

CLAIM. For any $j, 1 \leq j \leq t$, the grade of the ideal generated by the entries of $(xe_j - \sum \lambda_i \alpha_i)$ is greater than or equal to 1 (here $xe_j = \psi(e_j)$).

Proof. Let $\gamma_1, \dots, \gamma_t, \gamma_{t+1}, \dots, \gamma_{r_0}$ be a minimal set of generators of M where $\gamma_j = \phi(\overline{e}_j), 1 \leq j \leq t$ as above, and $t = \text{rank}_{\overline{R}} M$. Then $\text{grade}_{\overline{R}}(M') \geq 1$; that is, $\text{grade}_R(M') \geq 2$. Let $y \in \text{ann}_{\overline{R}} M'$ be a nonzero divisor in \overline{R} ; then $\forall k > t$, and $y\gamma_k = \sum_{j=1}^t a_{kj}\gamma_j$; hence, $ye_k - \sum a_{kj}e_j \in S$ for $k > t$. Let $P \in \text{Ass}_R(R)$, and let $q \in \text{Ass}_{\overline{R}}(\overline{R})$ containing P . By construction, q is an associated prime of M (recall that $\text{ann}_{\overline{R}} M = 0$); then M_q and T_q are free \overline{R}_q modules of rank t and $r_0 - t$, respectively. Since $\text{pd}_{R_q} M_q = 1, S_q$ is also a free R_q -module. We have the following short exact sequences:

$$(4) \quad 0 \rightarrow M_q \rightarrow \overline{S}_q \rightarrow T_q \rightarrow 0, \quad 0 \rightarrow T_q \rightarrow \overline{R}_q^{r_0} \rightarrow M_q \rightarrow 0.$$

Let $\overline{\beta}_k = \overline{e}_k - \sum_{j=1}^t \frac{a_{kj}}{y} \overline{e}_j$. Then $\{\overline{\beta}_k\}, t+1 \leq k \leq r_0$, is a basis of T_q . Let $\beta_k = e_k - \sum_{j=1}^t \frac{a_{kj}}{y} e_j, t+1 \leq k \leq r_0$. Then it follows from (4) that $\{xe_1, \dots, xe_t, \beta_{t+1}, \dots, \beta_{r_0}\}$ form a basis of S_q . In S_q , we have

$$(5) \quad \sum \lambda_i \alpha_i = \sum_{i=1}^t \frac{c_i}{b} xe_i + \sum_{k=t+1}^{r_0} \frac{d_k}{b} \beta_k, \quad b \notin q.$$

If $\sum \lambda_i \alpha_i \notin qS_q$, then $\sum \lambda_i \bar{\alpha}_i \notin qT_q$, which implies that some $d_k \notin q$. Hence, $\text{Im}(xe_j - \sum \lambda_i \alpha_i)$ is a minimal generator of T_q which is free, and thus $xe_j - \sum \lambda_i \alpha_i \notin PR^{r_0}$. Now suppose that $\sum \lambda_i \alpha_i \in qS_q$; then all $c_i, d_i \in q$ and $x \nmid d_k$ for at least one k in (5). Hence, $xe_j - \sum \lambda_i \alpha_i = (1 - \frac{c_j}{b})xe_j - \sum_{i \neq j} \frac{c_i}{b}xe_i - \sum \frac{d_{ik}}{b}\beta_k$. Since $1 - \frac{c_j}{b}$ is a unit in R_q , $xe_j - \sum \lambda_i \alpha_i$ is a minimal generator of S_q . Since S_q is a free R_q module, $xe_j - \sum \lambda_i \alpha_i \notin PR_q^{r_0}$, and hence $xe_j - \sum \lambda_i \alpha_i \notin PR^{r_0}$. This completes the proof of our claim. \square

Due to the commutative diagram (3) and Lemma 2.2, we can assume, without any loss of generality, that $\text{grade}_R M \geq 2$. Now we appeal to the following result due to Smoke [18, Lemma 4.1, Theorem 4.2]. Given a finitely generated R -module M of grade greater than or equal to 2, we can construct an exact sequence

$$(6) \quad 0 \rightarrow M \rightarrow R/I \rightarrow R/J \rightarrow 0,$$

where R/J has a filtration whose successive quotients are isomorphic to cyclic modules of the form $R/(u, v)$, where $\{u, v\}$ form an R -sequence. The proof in [18] shows that, by choosing the R -sequences of length 2 in m (annihilator of the corresponding module over R), this exact sequence can be constructed in such a way that, if $\delta : S \rightarrow I$ denotes the corresponding map on first syzygies via (6), then δ is surjective and $\bar{\delta} : S/mS \rightarrow I/mI$ is an isomorphism. Since $\text{pd}_R M < \infty, \text{pd}_R R/I < \infty$, and from the construction it follows that height of $I = \text{grade of } I = 2$. Thus, the proof of our theorem is complete. \square

Now let us state the following lemma.

LEMMA 2.4. *Let (R, m) be a local ring, and let M be a finitely generated module of projective dimension $n < \infty$. Let N be an R -module such that $\text{ann}_R M$ contains an N -sequence of length r . Then $\text{Tor}_{n-i}^R(M, N) = 0$ for $0 \leq i < r$.*

We leave the proof of this lemma to the reader.

In our next theorem we indicate the vanishing of Tor from an altogether different perspective.

THEOREM 2.5. *Let (R, m) be a local ring, and let M be a finitely generated module of finite projective dimension over R . Let $\text{pd}_R M = h$. Then there exists an R -sequence x_1, \dots, x_h of length h such that for any R -module N for which x_1, \dots, x_h form an N -sequence, $\text{Tor}_i^R(M, N) = 0$ for $i > 0$.*

Proof. If $\text{rank}_R M = s_0 > 0$, by Lemma 2.2 there exists a free submodule $F = R^{s_0}$ generated by a part of a minimal set of generators of M such that $M' = M/F$ has positive grade. From the commutative diagram

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 & & & R^{s_0} & & R^{s_0} & \\
 & & & \downarrow & & \downarrow & \\
 (1) & 0 & \rightarrow & S & \rightarrow & R^{r_0} & \rightarrow & M & \rightarrow & 0 \\
 & & & \downarrow & & \downarrow & & \downarrow & & \\
 & 0 & \rightarrow & S & \rightarrow & R^{r_0 - s_0} & \rightarrow & M' & \rightarrow & 0 \\
 & & & \downarrow & & \downarrow & & \downarrow & & \\
 & & & 0 & & 0 & & 0 & &
 \end{array}$$

it follows that $S = \text{Syz}_R^1(M) = \text{Syz}_R^1(M')$. If $\text{rank}_R M = 0$, then $M' = M$.

Let $x_1 \in \text{ann}_R M'$ be a nonzero divisor on R , let $R_1 = R/x_1R$, and let $S_1 = S/x_1S$. By Theorem 1.2, we have an exact sequence of R_1 -modules

$$(2) \quad 0 \rightarrow M' \rightarrow M_1 \rightarrow V_1 \rightarrow 0$$

such that $\text{pd}_{R_1} M_1 < \infty$ ($= \text{pd}_R M - 1$) and $\text{pd}_R V_1 = 1$. It is also clear from Corollary 1.3 that $S(M_1) = \text{Syz}_{R_1}^1(M_1) = S \oplus R^{t_1} = \text{Syz}_{R_1}^1(M') \oplus R^{t_1} = \text{Syz}_R^1(M) \oplus R^{t_1}$ for some $t_1 \geq 0$. Tensoring the short exact sequence $0 \rightarrow S(M_1) \rightarrow R^{\ell_1} \rightarrow M_1 \rightarrow 0$ with R_1 , we obtain the following short exact sequences:

$$(3) \quad 0 \rightarrow M_1 \rightarrow S(M_1) \otimes R_1 \rightarrow T_1 \rightarrow 0, \quad 0 \rightarrow T_1 \rightarrow R_1^{\ell_1} \rightarrow M_1 \rightarrow 0.$$

Here $T_1 = \text{Syz}_{R_1}^1(M_1)$. Recall that $\text{pd}_{R_1} M_1 < \infty$ and that $\text{support}(M_1) = \text{Spec } R_1$. Now we start with M_1 over R_1 and repeat the process described in (1), (2), and (3). We continue this process $(h - 2)$ times and obtain an R -sequence $x_1, \dots, x_i, 1 \leq i \leq h - 1$, modules M_i of finite projective dimension over $R_i = R/(x_1, \dots, x_i)$, and short exact sequences

$$(2i) \quad 0 \rightarrow M'_{i-1} \rightarrow M_i \rightarrow V_i \rightarrow 0,$$

$$(1i) \quad 0 \rightarrow F_i \rightarrow M_i \rightarrow M'_i \rightarrow 0,$$

where $\text{pd}_{R_i} M_i = \text{pd}_R M - i$, $\text{support}(M_i) = \text{support}(R_i)$, $\text{pd}_{R_{i-1}} V_i = 1$, $M'_{i-1} = M_{i-1}/F_{i-1}$, F_{i-1} a free R_{i-1} -module as constructed in (1), and M'_i is a module over R_{i+1} . We also have short exact sequences of R_i -modules

$$(3i) \quad 0 \rightarrow M_i \rightarrow S(M_i) \otimes R_i \rightarrow T_i \rightarrow 0, \quad 0 \rightarrow T_i \rightarrow R_i^{\ell_i} \rightarrow M_i \rightarrow 0.$$

We note that $\text{grade } M_i = i$ and that $\text{pd}_R M_i = h$. Let $x_h \in m$ be such that $\text{im}(x_h)$ in R_{h-1} is a nonzero divisor contained in $\text{ann}_{R_{h-1}} M'_{h-1}$. Then x_1, \dots, x_h form an R -sequence, and this is our required sequence.

Since projective dimension $\text{pd}_R M'_{h-1} = h$, if N is an R -module such that x_1, \dots, x_h form an N -sequence, then it follows from Lemma 2.4 that $\text{Tor}_i^R(M'_{h-1}, N) = 0$ for $i > 0$. Now it follows from the above short exact sequences starting from M'_{h-1} and tracing back to M that $\text{Tor}_j^R(M, N) = 0$ for $j \geq 1$. If $\text{grade}_R M = r > 0$ and x_1, \dots, x_r is an R -sequence contained in $\text{ann}_R M$, we start with Q —an $\overline{R}(= R/(x_1, \dots, x_r))$ -module as in Theorem 1.2—and construct an \overline{R} -sequence x_{r+1}, \dots, x_h by the above method. Then x_1, \dots, x_h form an R -sequence satisfying the required vanishing property of Tor . Since $\text{pd}_R M < \infty$, if $\text{grade}_R M = 0$, then $\text{ann}_R M = 0$, and hence $\text{rank}_R M > 0$. Thus we are back to diagram (1). \square

COROLLARY 2.6. *Let (R, m) be a local ring, and let M be a finitely generated module of finite projective dimension h over R . Let x_1, \dots, x_h be an R -sequence as mentioned in the above proposition. Suppose that for every $P \in \text{Ass}_R(R)$ there exists an R/P -module N such that x_1, \dots, x_h form an N -sequence and $N \neq mN$. Then, for every minimal generator β of $S_1 = \text{Syz}_R^1(M)$, $\text{grade } \mathcal{O}_{S_1}(\beta) \geq 1$.*

Proof. Let (F_\bullet, d_\bullet) be a minimal resolution of M over R . If possible, let $\text{grade } \mathcal{O}_{S_1}(\beta) = 0$. Then there exists an associated prime P such that $\mathcal{O}_{S_1}(\beta) \subset P$. Let $\overline{R} = R/P$, and let $\overline{F}_\bullet = F_\bullet \otimes R/P$. Consider the sequence

$$\overline{F}_2 \xrightarrow{\overline{d}_2} \overline{F}_1 \xrightarrow{\overline{d}_1} \overline{F}_0 \rightarrow 0.$$

Since $\mathcal{O}_{S_1}(\beta) \subset P$, $\overline{d}_1(\beta) = 0$. This implies that, for any \overline{R} -module N , $\text{Tor}_1^{\overline{R}}(M, N) \neq 0$. However, by hypothesis, there exists an \overline{R} -module N such that x_1, \dots, x_h form an N -sequence. Then, by Theorem 2.5, we have $\text{Tor}_j^{\overline{R}}(M, N) = 0$ for $j > 0$, which leads to a contradiction. Hence, $\text{grade } \mathcal{O}_{S_1}(\beta) \geq 1$. \square

COROLLARY 2.7. *Let (R, m) be a local ring, and let M be a finitely generated module of finite projective dimension h over R . Let x_1, \dots, x_h be an R -sequence as mentioned in Theorem 2.5. If R has mixed characteristic $p > 0$, we assume that p, x_1, \dots, x_h form a part of a system of parameters of R . Then for every minimal generator β of $S_1 = \text{Syz}_R^1(M)$, $\text{grade } \mathcal{O}_{S_1}(\beta) \geq 1$.*

Proof. If possible, let $\text{grade } \mathcal{O}_{S_1}(\beta) = 0$. Then there exists an associated prime P of R such that $\mathcal{O}_{S_1}(\beta) \subset P$. If R is equicharacteristic, then there exists a big Cohen–Macaulay R/P -module N such that x_1, \dots, x_h form a regular N -sequence. If R has mixed characteristic $p > 0$, then there exists a big Cohen–Macaulay $R/(P + pR)$ -module N such that x_1, \dots, x_h form a regular N -sequence. Hence, we are done by Corollary 2.6. \square

REMARK. Due to the existence of Cohen–Macaulay algebras over local domains of dimension less than or equal to 3 (see [14]), it can be checked from the above arguments that finitely generated modules of projective dimension less than or equal to 3 satisfy the order ideal conjecture.

LEMMA 2.8. *Let $(R, m, K = R/m)$ be an equicharacteristic complete local ring of dimension d , and let x_1, \dots, x_d be a system of parameters of R . Let M be a big Cohen–Macaulay R -module such that $\mathbf{x}M \neq M$ and x_1, \dots, x_d form a maximal M -sequence. Then $\widehat{M} =$ the m -adic completion of M is a flat $K[[x_1, \dots, x_d]]$ -module.*

Proof. Let $S = K[[x_1, \dots, x_d]]$. Then S is a complete power series ring in d variables, R is a module-finite extension of S , and $\mathbf{x} = \{x_1, \dots, x_d\}$ form a regular system of parameters of S . Moreover, \mathbf{x} is \widehat{M} -regular (see [3, Theorem 8.5.1]), and \widehat{M} is a balanced big Cohen–Macaulay module (see [3, Corollary 8.5.3]). Hence, $\psi : \frac{\widehat{M}}{\mathbf{x}\widehat{M}}[X_1, \dots, X_d] \rightarrow \bigoplus_{n=0}^{\infty} \frac{\mathbf{x}^n \widehat{M}}{\mathbf{x}^{n+1} \widehat{M}}$ is an isomorphism. Since \mathbf{x} is a regular system of parameters of S , $K[X_1, \dots, X_d] \simeq \bigoplus_{n=0}^{\infty} \frac{\mathbf{x}^n S}{\mathbf{x}^{n+1} S}$. Hence, $\frac{\mathbf{x}^n S}{\mathbf{x}^{n+1} S} \otimes_K \widehat{M}/\mathbf{x}\widehat{M} \simeq \frac{\mathbf{x}^n \widehat{M}}{\mathbf{x}^{n+1} \widehat{M}}$, and $\frac{\widehat{M}}{\mathbf{x}\widehat{M}}$ is a nonnull vector spacing over $K = S/\mathbf{x}S$. Thus, it follows, by [2, Theorem 1, Section 5.2], that \widehat{M} is S -flat. \square

THEOREM 2.9 ([11, Theorem 6.9]). *Let (R, m, K) be an equicharacteristic complete local ring, and let N be a finitely generated module of finite projective dimension over R . Let I be an ideal of height h , and let M be a big Cohen–Macaulay module over R/I . Let \widehat{M} be the m -adic completion of M . Then $\text{Tor}_i^R(\widehat{M}, N) = 0$ for $i > h$.*

Proof. For a proof, we refer the reader to [8, Theorem 1.11] or [11], where the existence of a big Cohen–Macaulay R/I -module Q such that $\text{Tor}_i^R(Q, N) = 0$, for $i > h$, has been demonstrated. (It was first proved by Foxby [11].) In these proofs, it was required that such a Q be free over a certain complete regular local ring S contained in R/I such that R/I is a

module-finite extension of S . By Lemma 2.8, the completion \widehat{M} of any big Cohen–Macaulay R/I module is flat over certain complete regular subrings of R/I . This flatness is enough to ensure the validity of arguments provided in [8, Theorem 1.11] or in [11, Theorem 6.9] for proving our assertion. \square

Next we recall Shimomoto’s theorem.

THEOREM 2.10 ([17, Theorem 5.3]). *Let (R, m) be a complete local domain of mixed characteristic $p > 0$. Then there exists some system of parameters p, x_2, \dots, x_d of R and an almost Cohen–Macaulay quasilocal algebra B over R^+ in the sense that*

- (1) $(p, x_2, \dots, x_d)B \neq B$,
- (2) x_2, \dots, x_d form a regular sequence on B/pB , and
- (3) p is not nilpotent in B and the ideal $(0 : p)_{\mathbf{B}}$ is annihilated by p^ϵ for any rational $\epsilon > 0$.

Actually, Shimomoto’s construction shows that such a B can be constructed for any system of parameters of the form p, x_2, \dots, x_d of R .

DEFINITION. An almost Cohen–Macaulay algebra B as above is called *balanced* if B/pB is a balanced Cohen–Macaulay algebra.

Now we are ready to prove our final theorem.

THEOREM 2.11. *Let (R, m) be a local ring of mixed characteristic $p > 0$. We assume that either p is nilpotent or that p is a nonzero divisor in R . Let M be a finitely generated module of finite projective dimension over R , and let β be a minimal generator of S_i , the i th syzygy of M (minimal), for $i > 0$. We have the following.*

- (a) *If p is nilpotent, the order ideal conjecture is valid on R .*
- (b) *If $pM = 0$, then the grade of $\mathcal{O}_{S_1}(\beta) \geq 1$.*
- (c) *Suppose that balanced or complete almost Cohen–Macaulay algebras exist over complete local domains. If $pM = 0$ and $p \in m - m^2$, then the grade of $\mathcal{O}_{S_i}(\beta) \geq i$, $\forall i \geq 1$.*
- (d) *Assume that every element in $m - m^2$ is a nonzero divisor and that the order ideal conjecture is valid over R/xR for any $x \in m - m^2$. If $\text{ann}_R M \cap (m - m^2) \neq \phi$ or $\text{depth}_R M > 0$, then $\text{grade } \mathcal{O}_{S_i}(\beta) \geq i$, $\forall i \geq 1$.*

Proof. (a) If p is nilpotent, the proof follows immediately by similar arguments as in [9, Theorem 2.4] due to the existence of big Cohen–Macaulay

modules on equicharacteristic local domains R/P for every prime ideal P in $\text{Spec } R$. One may also use [3, Lemma 9.1.8] to prove the assertion.

(b) Since p is a nonzero divisor on R , by Corollary 1.3 we can assume that $\text{pd}_{R/pR} M < \infty$. We write $\overline{R} = R/pR$. Consider the short exact sequence

$$(1) \quad 0 \rightarrow S_1 \xrightarrow{g} R^{r_0} \xrightarrow{\eta} M \rightarrow 0.$$

Tensoring this sequence with \overline{R} , we obtain the exact sequences

$$(2) \quad \mathcal{O} \rightarrow M \xrightarrow{j} \overline{S}_1 \xrightarrow{b} T_1 \rightarrow 0,$$

$$(3) \quad \mathcal{O} \rightarrow T_1 \xrightarrow{\gamma} \overline{R}^{r_0} \xrightarrow{\overline{\eta}} M \rightarrow 0,$$

where $T_1 = \text{Syzy}_{\overline{R}}^1(M)$ and $\overline{S}_1 = S_1 \otimes_R \overline{R}$. For any minimal generator ϑ of M we have $\vartheta = \eta(e)$, e being a minimal generator of R^{r_0} and $j(\vartheta) = \text{class of } pe \text{ in } \overline{S}$. Recall that if I is an ideal of R such that $\text{grade}_R I = 0$, then for any nonzero divisor x in R , $\text{grade}_{\overline{R}}(I + xR)/xR$ is also 0. Hence, for any minimal generator β of S_1 , if $b(\beta)$ is a minimal generator of T_1 , then our assertion follows due to the validity of the order ideal conjecture on equicharacteristic local rings. If $b(\beta) = 0$, then $\beta \in j(M)$; that is, $\beta = j(\vartheta)$, where ϑ is a minimal generator of M . Let $\{\overline{\alpha}_j\}_{1 \leq j \leq t}$ be a minimal set of generators of T_1 , and let $\{\alpha_j\}$ denote a lift of $\{\overline{\alpha}_j\}$ in S_1 . Then there exists a basis $e_1, \dots, e_i, \dots, e_{r_0}$ of R^{r_0} such that $\{pe_1, \dots, pe_i, \alpha_j s\}$ form a minimal set of generators of S_1 . Moreover, to show that, for any minimal generator β of S_1 , $\text{grade}_{\mathcal{O}_{S_1}}(\beta) \geq 1$, it is enough to take $\beta = pe - \sum_{j=1}^t \lambda_j \alpha_j$, where $e \in \{e_1, \dots, e_i\}$, $\lambda_j s \in m$, and $\sum \lambda_j \overline{\alpha}_j \neq 0$ in T_1 .

If possible, let $\text{grade}_{\mathcal{O}_{S_1}}(\beta) = 0$, that is, let $\mathcal{O}_{S_1}(\beta) \subset P$, for some $P \in \text{Ass}_R(R)$. Let $\text{pd}_R M = h$; then $\text{pd}_{\overline{R}} M = (h - 1)$. By Theorem 2.5, corresponding to M over \overline{R} , there exists an \overline{R} sequence $\overline{x}_1, \dots, \overline{x}_{h-1}$ satisfying the assertion mentioned in Theorem 2.5. Let $\tilde{R} = R/pR$, let $\tilde{p} = \text{im}(p)$ in \tilde{R} , and let $\tilde{x}_i = \text{im}(x_i)$ in \tilde{R} , $1 \leq i \leq h - 1$. Then $\tilde{p}, \tilde{x}_1, \dots, \tilde{x}_{h-1}$ form a part of a system of parameters of \tilde{R} . By Corollary 2.7 there exists an almost Cohen–Macaulay \tilde{R}^+ -algebra B such that $(\tilde{p}, \dots, \tilde{x}_{h-1})B \neq B$ and $\tilde{x}_1, \dots, \tilde{x}_{h-1}$ form a regular sequence on B/pB . Then, by Theorem 2.5 or by [11, Theorem 2.4], we have $\text{Tor}_j^{\overline{R}}(M, B/pB) = 0$ for $j > 0$.

We consider the following part of a minimal resolution F_\bullet of M over R :

$$\begin{array}{ccccccc} R^{r_2} & \xrightarrow{d_2} & R^{r_1} & \xrightarrow{d_1} & R^{r_0} & \rightarrow & 0 \\ \parallel & & \parallel & & \parallel & & \\ F_2 & \rightarrow & F_1 & \rightarrow & F_0 & \rightarrow & 0 \end{array}$$

Let $\tilde{F}_i = F_i \otimes \tilde{R}$, $0 \leq i \leq 2$. Tensoring the above sequence with \tilde{R} , we obtain a sequence

$$\tilde{F}_2 \xrightarrow{\tilde{d}_2} \tilde{F}_1 \xrightarrow{\tilde{d}_1} \tilde{F}_0 \rightarrow 0,$$

where $\tilde{\beta} = \text{im}(\tilde{\beta})$ in $\tilde{F}_0 = 0$.

Tensoring the above sequence with B and writing $F_{iB} = F_i \otimes B$, we have

$$(4) \quad F_{2B} \xrightarrow{d_{2B}} F_{1B} \xrightarrow{d_{1B}} F_{0B} \rightarrow 0,$$

where $\beta_B = \text{im } \tilde{\beta}$ in $F_{0B} = 0$. Hence,

$$(5) \quad pe_B = \sum \lambda_j \alpha_{jB} \quad \text{in } B^{r_0} = F_{0B},$$

where $\alpha_{jB} = \text{im}(\alpha_j \otimes 1_B)$ in F_{0B} , $\alpha_j \otimes 1_B \in S_1 \otimes B$. Since $\text{Tor}_j^{\overline{R}}(M, B/pB) = 0$, for $j > 0$, tensoring (2) and (3) with B/pB over \overline{R} we obtain the following short exact sequences

$$(6) \quad 0 \rightarrow M \otimes B/pB \xrightarrow{j \otimes 1_{B/pB}} \overline{S}_1 \otimes B/pB \rightarrow T_1 \otimes B/pB \rightarrow 0$$

and

$$(7) \quad \mathcal{O} \rightarrow T_1 \otimes B/pB \rightarrow (B/pB)^{r_0} \rightarrow M \otimes B/pB \rightarrow 0.$$

Let $\overline{\alpha}_{jB}$ = the image of $(\overline{\alpha}_j \otimes 1_{B/pB})$ in $T_1 \otimes B/pB$ in $(B/pB)^{r_0} = \text{im } \alpha_{jB}$ in $(B/pB)^{r_0}$. Due to (5) and (7), we have $\sum \lambda_j \overline{\alpha}_{jB} = 0$ in $(B/pB)^{r_0}$, and hence $\sum \lambda_j (\overline{\alpha}_j \otimes 1_{\overline{B}}) = 0$ in $T_1 \otimes B/pB$. Since $\lambda_j s \in m$, this implies, due to the exact sequence (6) and the definition of j in (2), that

$$(8) \quad \sum \lambda_j (\alpha_j \otimes 1_B) = \sum a_i (pe_i \otimes 1_B) + p \left(\sum b_i (pe_i \otimes 1_B) + \sum \mu_i (\alpha_i \otimes 1_B) \right)$$

in $S_1 \otimes B$, $a_i s \in m_B$, where m_B is the maximal ideal of B .

Hence, we have from (5) and (8)

$$pe_B = \sum (a_i + pb_i) pe_{iB} + p \sum \mu_i \alpha_{iB}$$

in B^{r_0} ; that is,

$$(*) \quad p \left\{ e_B - \left[\sum (a_i + pb_i) e_{iB} + \sum \mu_i \alpha_{iB} \right] \right\} = 0.$$

Since $a_i \in m_B$, entries of $\alpha_{i\beta} \in m_B$, $e_B - [\sum(a_i + pb_i)e_i + \sum \mu_i \alpha_{iB}]$ is a free generator of B^{r_0} ; hence, p cannot annihilate it. Thus, $(*)$ leads to a contradiction. Hence, grade $\mathcal{O}_{S_1}(\beta)$ must be greater than or equal to 1.

(c) We assume that $p \in m - m^2$ and that $pM = 0$. Let $F_\bullet : 0 \rightarrow R^{r_n} \xrightarrow{d_n} R^{r_{n-1}} \rightarrow \dots \rightarrow R^{r_1} \xrightarrow{d_1} R^{r_0} \rightarrow 0$ be a minimal free resolution of M over R , and let $P_\bullet : 0 \rightarrow \overline{R}^{s_{n-1}} \rightarrow \dots \rightarrow \overline{R}^{s_1} \rightarrow \overline{R}^{s_0} \rightarrow 0$ be a minimal free resolution of M over $\overline{R}(=R/pR)$. Shamash [16] has shown that P_\bullet can be obtained from F_\bullet via the homotopy maps $h_\bullet : F_\bullet \rightarrow F_\bullet(+1)$ induced by the 0-map on M due to multiplication by p . Actually, each F_i , for $i > 0$, can be decomposed into two parts: $F_i = F'_i \oplus F''_i$, where $h_i(F''_i) = 0$, every free generator e' of F'_{i-1} is such that $e = h_{i-1}(e')$ is a free generator of F''_i , and $d_i(e) = pe' - h_{i-2}d_{i-1}(e')$. Moreover, it follows from Shamash's theorem that (i) there exists $\phi_\bullet : P_\bullet \rightarrow \overline{F_\bullet(+1)}$ ($\overline{F_\bullet} = F_\bullet \otimes_R \overline{R}$), where ϕ_0 induces the inclusion map $M = \text{Tor}_1^R(M, R/p) \xrightarrow{j} \overline{S}_1$, and (ii) ϕ_\bullet induces a splitting on each component of P_\bullet and $P_\bullet(+1)$ can be extracted from the mapping cone of ϕ_\bullet . Let $S_i = \text{Syz}_R^i(M)$, let $T_i = \text{Syz}_{\overline{R}}^i(M)$, and let $\overline{S}_i = S_i \otimes \overline{R}$. This leads to the following short exact sequences for $i > 1$:

$$(1') \quad 0 \rightarrow T_{i-1} \xrightarrow{\psi} \overline{S}_i \xrightarrow{\eta} T_i \rightarrow 0,$$

$$(2') \quad 0 \rightarrow T_i \rightarrow \overline{R}^{r_{i-1}} \rightarrow T_{i-1} \rightarrow 0,$$

where ψ is induced by ϕ_\bullet and $\psi(\text{Im } \eta') = \overline{pe - h_{i-2}d_{i-1}(e')}$ is a minimal generator of \overline{S}_i due to the splitting property of ϕ_\bullet . Let $\gamma = pe' - h_{i-2}d_{i-1}(e')$, and let

$$(3') \quad \overline{\gamma} = \overline{pe' - h_{i-2}d_{i-1}(e')}.$$

Then γ and $\overline{\gamma}$ are minimal generators of S_i and \overline{S}_i , respectively. □

CLAIM. *In the above situation, grade $\mathcal{O}_{S_i}(\gamma) \geq i$.*

Proof. If possible let P be a prime ideal of height $(i - 1)$ containing $\mathcal{O}_{S_i}(\gamma)$. Since $e' \in F'_{i-1}$ and $h_{i-2}(F_{i-2}) = F''_{i-1}$, it follows that $p \in P$. Let p, x_2, \dots, x_{i-1} be a maximal R -sequence contained in $\mathcal{O}_{S_i}(\gamma)$ —we denote it by \mathbf{x} . Then $\text{Tor}_j^R(M, R/\mathbf{x}) = 0$ for $j \geq i$. Let $R' = R/\mathbf{x}$, let $S'_i = S_i \otimes R'$, and so forth. We have an exact sequence

$$0 \rightarrow S'_i \rightarrow R'^{r_{i-1}} \rightarrow S'_{i-1} \rightarrow 0.$$

Then S'_{i-1} is a module of finite projective dimension over R' , and $S'_i = \text{Syz}_{R'}^1(S'_{i-1})$ has a minimal generator $\gamma' = \text{im } \gamma$ in S'_i such that $\mathcal{O}_{S'_i}(\gamma')$ has grade 0 in R' . This contradicts part (b) of our theorem, and hence the claim is established.

Let $\{\bar{\alpha}_j\}$ form a minimal set of generators of T_i , and let $\{\alpha_j\}$ denote their lifts in S_i . Since characteristic of $\bar{R} = p > 0$, by the theorem by Evans and Griffith [9, Theorem 2.4], $\text{grade}_{\bar{R}}(\mathcal{O}_{T_i}(\bar{\alpha}_j)) \geq i$. Since $(1') \otimes R/m$ is exact due to the above claim, for the purpose of proving our theorem it would be enough to establish that, for every minimal generator β of S_i of the form $\beta = \gamma - \sum \lambda_j \alpha_j$, $\text{grade } \mathcal{O}_{S_i}(\beta) \geq i$, where $\bar{\gamma} = \psi(\text{im } \bar{e}')$ for some free generator e' for F_{i-1} and $\lambda_j s \in m$. If possible, let $\mathcal{O}_{S_i}(\beta) \subset P$ — a prime ideal of height $(i - 1)$. If $p \in P$, there exists a maximal Cohen–Macaulay R/P -algebra $((R/P)^+$ -algebra) \bar{B} . If $p \notin P$, then by assumption there exists a balanced almost Cohen–Macaulay R/P -algebra $((R/P)^+$ -algebra) B such that p is not nilpotent on B and such that $\bar{B} = B/pB$ is a maximal Cohen–Macaulay algebra over $R/(P + pR)$. Hence, in either case, by [11, Theorem 2.4], $\text{Tor}_j^{\bar{R}}(M, \bar{B}) = 0$ for $j \geq i$. Tensoring $(1')$ and $(2')$ with \bar{B} , we get the exact sequences

$$(6') \quad \mathcal{O} \rightarrow T_{i-1} \otimes \bar{B} \xrightarrow{\psi \otimes 1_{\bar{B}}} \bar{S}_i \otimes \bar{B} \rightarrow T_i \otimes \bar{B} \rightarrow 0$$

and

$$(7') \quad \mathcal{O} \rightarrow T_i \otimes \bar{B} \rightarrow \bar{B}^{r_{i-1}} \rightarrow T_{i-1} \otimes \bar{B} \rightarrow 0.$$

Let e be a free generator of F_i such that $d_i(e) = \beta$.

From F_\bullet , we consider the exact sequence

$$F_{i+1} \xrightarrow{d_{i+1}} F_i \xrightarrow{d_i} F_{i-1}.$$

Let $\tilde{R} = R/P$, and let $\tilde{F}_i = F_i \otimes_R \tilde{R}$. Tensoring the above sequence with \tilde{R} , we obtain a complex

$$\tilde{F}_{i+1} \xrightarrow{\tilde{d}_{i+1}} \tilde{F}_i \xrightarrow{\tilde{d}_i} \tilde{F}_{i-1},$$

where $\tilde{d}_i(\tilde{e}) = 0$. Let $e_B = e \otimes 1_B$. Then in the complex $F_{i+1} \otimes B \rightarrow F_i \otimes B \rightarrow F_{i-1} \otimes B$, we have $d_{iB}(e_B) = 0$. Let $\gamma_B = \text{im}(\gamma \otimes 1_B)$, and let $\alpha_{jB} = \text{im}(\alpha_j \otimes 1_B)$ in $B^{r_{i-1}} = F_{i-1} \otimes B$. Then

$$(5') \quad d_{iB}(e_B) = 0 \Rightarrow \gamma_B - \sum \lambda_j \alpha_{jB} = 0.$$

Let $\bar{\alpha}_{jB} = \text{Im } \alpha_{jB}$ in $\bar{B}^{r_{i-1}}$. Due to (5') and (6'), we have $\sum \lambda_j \bar{\alpha}_{jB} = 0$ in $\bar{B}^{r_{i-1}}$; hence, $\sum \lambda_j (\bar{\alpha}_j \otimes 1_B) = 0$ in $T_i \otimes \bar{B}$. Since $\lambda_j s \in m$, this implies, due to the exact sequence (6') and definition of ψ in (1'), that

$$(8') \quad \begin{aligned} \sum \lambda_j (\alpha_j \otimes 1_B) &= \sum a_i (\gamma_i \otimes 1_B) \\ &+ p \left(\sum b_i (\gamma_i \otimes 1_B) + \sum \mu_j (\alpha_j \otimes 1_B) \right) \end{aligned}$$

in $S_i \otimes B, a_i s \in m_B$. Hence, from (5') we have

$$\gamma_B = \sum (a_i + pb_i) (\gamma_{iB}) + p \sum \mu_j \alpha_{jB}$$

in $B^{r_{i-1}}$. Since $\gamma_B = pe'_B - h_{i-2} d_{i-1} (e'_B)$ and γ_{iB} 's also have similar expressions, comparing the e'_B th coordinate in $B^{r_{i-1}}$, we obtain from above

$$(**) \quad p \left[e'_B - \sum (a_i + pb_i) e_{B'} - \sum \delta_j \right] = 0,$$

where δ_j is the e'_B th coordinate of $\mu_j \alpha_{jB}$. Since $a_i, \delta_j \in m_B$, the term within brackets in (**) is a free generator of $B^{r_{i-1}}$, and hence p cannot annihilate it. Thus, (**) leads to a contradiction. Hence, $\text{grade } \mathcal{O}_{S_i}(\beta) \geq i$, and our proof is complete.

(d) We assume that every element in $m - m^2$ is a nonzero divisor and that for any such element x the order ideal conjecture is valid for R/xR . Let M be a finitely generated module of finite projective dimension such that either $\text{ann}_R M \cap (m - m^2) \neq \phi$ or $\text{depth}_R M > 0$.

First, let us assume that $\text{ann}_R M \cap (m - m^2) \neq \phi$. Let $x \in \text{ann}_R M \cap (m - m^2)$, and let $\bar{R} = R/xR$. Since $\text{pd}_R M < \infty$ and $x \in m - m^2$, $\text{pd}_{\bar{R}} M < \infty$. By hypothesis, M satisfies the order ideal conjecture as an \bar{R} -module. Let (F_\bullet, d_\bullet) and (P_\bullet, S_\bullet) denote minimal free resolutions of M over R and \bar{R} , respectively; let $S_i = \text{Syz}_R^i(M)$, and let $T_i = \text{Syz}_{\bar{R}}^i(M)$. Arguing as in the proof of part (b) of the theorem ((1), (2), (3), etc.), we construct a minimal set of generators $\{xe_1, \dots, xe_i, \alpha_j s\}$ such that $\{e_1, \dots, e_j\}$ form a part of a basis of F_0 , and $\bar{\alpha}_j (= \text{im } \alpha_j \text{ in } T_1)$ form a minimal set of generators of T_1 . To show that, for any minimal generators β of S_1 , $\text{grade } \mathcal{O}_{S_1}(\beta) \geq 1$, due to inductive hypothesis, it is enough to take $\beta = xe - \sum \lambda_i \alpha_i, e \in \{e_1, \dots, e_i\}$, and $\lambda_i \in m$. Then β_e is the e th coordinate of $\beta = x - \sum \lambda_i \alpha_{ie}$. Since $\lambda_i s \in m, \alpha_{ie} s \in m$, and $x \in m - m^2$, we have $\beta_e \neq 0, \beta_e \in m - m^2$, and hence by assumption β_e is a nonzero divisor in R . Thus, the conclusion follows for $i = 1$.

Now consider $i > 1$. Arguing as in part (c) above we see from (1'), (2'), (3'), and so on, that it is enough to consider a minimal generator β of S_i of the form $\beta = \gamma - \sum \lambda_i \alpha_i$, where $\gamma = xe' - h_{i-2}h_{i-1}(e') = d_i(e)$, $e = h_{i-1}(e')$, e' a minimal generator of F'_{i-1} , and $\bar{\alpha}_j (= \text{im } \alpha_j \in T_i)$ form a minimal set of generators of T_i . Similar arguments as in part (c) show that the grade of $\mathcal{O}_{S_i}(\gamma) \geq i$ and, by hypothesis, that $\text{grade } \mathcal{O}_{T_i}(\bar{\alpha}_j) \geq i$. Let J denote the ideal generated by entries of β . Recall that $F_{i-1} = F'_{i-1} \oplus F''_{i-1}$, that e' is a minimal generator of F'_{i-1} , and that $h_{i-2}d_{i-1}(e') \in F''_{i-1}$. Let $y =$ the e' th coordinate of $\beta = x - \sum \lambda_i \alpha_{ie'}$; since $\lambda_i s \in m$ and $\alpha_{ie'} s \in m$, $y \in m - m^2$, and hence, by assumption, y is a nonzero divisor in R . Let $\bar{R} = R/yR$, and let $\bar{S}_i = S_i/yS_i$ for $i \geq 1$; then $\bar{S}_i = \text{Syz}_{\bar{R}}^{i-1}(\bar{S}_1)$. By hypothesis J/yR has grade $\geq (i - 1)$ in \bar{R} . This implies that $\text{grade}_R J \geq i$, and our proof is complete.

Now assume that $\text{depth}_R M > 0$. We can find $x \in m - m^2$ such that x is a nonzero divisor on M . Let $\bar{R} = R/xR$, and let $\bar{M} = M/xM$. Since $\text{pd}_{\bar{R}} \bar{M} = \text{pd}_R M < \infty$, by hypothesis, \bar{M} satisfies the order ideal conjecture over \bar{R} . Hence M satisfies the order ideal conjecture over R . □

COROLLARY 2.12. *Let (R, m) be a local ring of mixed characteristic $p > 0$ such that p is a nonzero divisor in R . Let I be an ideal of finite projective dimension over R . If $p \in I$ or p is a nonzero divisor on R/I , then every minimal generator of I is a nonzero divisor in R . In particular, if P is a prime ideal of finite projective dimension over R , then every minimal generator of P is a nonzero divisor.*

Proof. If $p \in I$, the proof follows from Theorem 2.11(b); if p is a nonzero divisor on R/I , the result follows from the validity of the order ideal conjecture on R/pR (see [9]). □

COROLLARY 2.13. *Let (R, m) be a regular local ring of dimension n , and assume that the order ideal conjecture is valid for regular local rings of dimension $(n - 1)$. If M is a finitely generated R -module such that either M is annihilated by a regular parameter or $\text{depth}_R M > 0$, then M satisfies the order ideal conjecture.*

The proof follows from Theorem 2.11(d). □

REMARK. After going through Theorem 1.2 of this paper, one of the referees pointed out that this theorem could be achieved by combining an embedding theorem for modules of finite G -dimension due to Christensen and

Iyengar (“Gorenstein dimension of modules over homomorphisms”) with a couple of results from “complete intersection dimension” by Avramov, Gasharov, and Peeva. This author was not aware of this observation. It is also clear from the comments that the other referee and most of the experts are also unaware of this observation. After taking a glance at the proposed proof via this observation, it seems that this approach would require several other notions and is rather technical and not constructive. In comparison, our proof is much more direct and to the point, and it exploits only the Ext characterization of modules of finite projective dimension.

Acknowledgments. I am thankful to the referees for their suggestions toward improving the exposition in this paper.

REFERENCES

- [1] M. Auslander and R.-O. Buchweitz, “The homological theory of maximal Cohen–Macaulay approximations” in *Colloque en l’honneur de Pierre Samuel (Orsay, 1987)*, Mém. Soc. Math. Fr. (N.S.) **38**, Soc. Math. Fr., Marseilles, 1989, 5–37. [MR 1044344](#).
- [2] N. Bourbaki, *Elements of Mathematics: Commutative Algebra*, Addison-Wesley, Reading, MA, 1972. [MR 0360549](#).
- [3] W. Bruns and J. Herzog, *Cohen–Macaulay Rings*, Cambridge Stud. Adv. Math. **39**, Cambridge University Press, Cambridge, 1993. [MR 1251956](#).
- [4] S. P. Dutta, *On the canonical element conjecture*, Trans. Amer. Math. Soc. **299**, no. 2 (1987), 803–811. [MR 0869233](#). [DOI 10.2307/2000525](#).
- [5] ———, *On negativity of higher Euler characteristics*, Amer. J. Math. **126** (2004), 1341–1354. [MR 2102398](#).
- [6] ———, *The monomial conjecture and order ideals*, J. Algebra **383** (2013), 232–241. [MR 3037977](#). [DOI 10.1016/j.jalgebra.2013.03.006](#).
- [7] E. G. Evans and P. Griffith, *The syzygy problem*, Ann. of Math. (2) **114** (1981), 323–333. [MR 0632842](#). [DOI 10.2307/1971296](#).
- [8] ———, *Syzygies*, London Math. Soc. Lecture Note Ser. **106**, Cambridge University Press, Cambridge, 1985. [MR 0811636](#). [DOI 10.1017/CBO9781107325661](#).
- [9] ———, “Order ideals” in *Commutative Algebra (Berkeley, CA, 1987)*, Math. Sci. Res. Inst. Publ. **15**, Springer, New York, 1989, 213–225. [MR 1015519](#). [DOI 10.1007/978-1-4612-3660-3_10](#).
- [10] ———, *A graded syzygy theorem in mixed characteristic*, Math. Res. Lett. **8** (2001), 605–611. [MR 1879804](#). [DOI 10.4310/MRL.2001.v8.n5.a2](#).
- [11] H.-B. Foxby, *On the μ^i in a minimal injective resolution, II*, Math. Scand. **41** (1977), 19–44. [MR 0476801](#).
- [12] M. Hochster, *Topics in the Homological Theory of Modules over Commutative Rings*, CBMS Reg. Conf. Ser. Math. **24**, Amer. Math. Soc., Providence, 1975. [MR 0371879](#).
- [13] ———, *Canonical elements in local cohomology modules and the direct summand conjecture*, J. Algebra **84** (1983), 503–553. [MR 0723406](#). [DOI 10.1016/0021-8693\(83\)90092-3](#).
- [14] ———, *Big Cohen–Macaulay algebras in dimension three via Heitmann’s theorem*, J. Algebra **254** (2002), 395–408. [MR 1933876](#). [DOI 10.1016/S0021-8693\(02\)00086-8](#).

- [15] C. Peskine and L. Szpiro, *Dimension projective finie et cohomologie locale*, Publ. Math. Inst. Hautes Études Sci. **42** (1973), 47–119. [MR 0374130](#).
- [16] J. Shamash, *The Poincaré series of a local ring*, J. Algebra **12** (1969), 453–470. [MR 0241411](#).
- [17] K. Shimomoto, *Almost Cohen–Macaulay algebras in mixed characteristic via Fontaine rings*, Illinois J. Math. **55** (2011), 107–125. [MR 3006682](#).
- [18] W. Smoke, *Perfect modules over Cohen–Macaulay local rings*, J. Algebra **106** (1987), 367–375. [MR 0880963](#). [DOI 10.1016/0021-8693\(87\)90002-0](#).

Department of Mathematics
University of Illinois
Urbana, Illinois 61801
USA
s-dutta@illinois.edu