

A VLA/MERLIN/VLBA FOR INTERMEDIATE SCALE LENSES AND THE DISCOVERY OF A NEW LENS SYSTEM?

P. AUGUSTO, P.N. WILKINSON AND I.W.A. BROWNE
*Nuffield Radio Astronomy Laboratories, Univ. of Manchester
Jodrell Bank, Macclesfield, Cheshire, SK11 9DL, U.K.*

We are searching for small lens systems (50-250 mas or $10^8 - 10^9 M_{\odot}$) in a sample of ~ 1800 flat spectrum radio sources. This is the first time a systematic search has been made “between” the VLA and VLBI resolutions. Finding any would indicate the existence of other than the “conventional” spiral/elliptical lenses (only $\sim 0.01\%$ chance - Turner et al. (1984)). For example, *faint galaxies* are numerous ($\sim 10^6$ gal/deg² - Lilly (1993), Glazebrook et al. 1995), compact (HST Medium Deep Survey (MDS) - Griffiths et al. 1994) and ideally placed for lensing ($\langle z \rangle \sim 0.6$ - MDS, Smail et al. (1994); *c.f.* Turner et al. 1984). Early-type *dwarf galaxies* (dE,N and cE), if extant at intermediate- z as favored by MDS are also obvious lens candidates. If no lenses are found, a limit 400 times better than the current one (Surdej et al. 1993), $\Omega_L < 0.001$, will be placed on the cosmological density of compact objects (*e.g.* black holes) for the above mass range.

From the JVAS (Patnaik et al. 1992) and the Cosmic Lens All Sky Survey (CLASS) (Browne et al. this volume) - we selected a parent sample of ~ 1800 objects satisfying $\alpha_{1.4-5} < 0.50$ ($S_{\nu} \propto \nu^{-\alpha}$), $S_{8.4}^{total} \geq 100$ mJy and $|b^{II}| \geq 10^{\circ}$. The 67 “candidates” were selected from the visibility plots (the sources > 50 mas) and constitute an interesting population: two thirds of the candidates have $\alpha \sim 0.3 - 0.4$ whereas the parent sample has $\langle \alpha \rangle = 0.0$; in addition, the candidate sample is clearly dominated by empty fields (40%) whereas the parent sample has 60% of QSOs.

Our 67 candidates were mapped with MERLIN “snapshots” at 5 GHz (4×15 min.). About 20 sources look very interesting for subsequent VLBA follow-up; the great majority of the remaining appear to be “core-jets”.

One arcsecond scale source, 2114+022, looked promising enough for long track MERLIN observations (see Fig.1) and it is a good lens candidate despite its very strange configuration; there will be a multi-frequency VLBA follow-up.




Figure 1. MERLIN map of 2114+022 at 5 GHz (11hrs. phase-referencing). The noise level is very low (< 0.1 mJy) and there are no signs of any extended structure; all the four images are unresolved. The (variable?) fluxes of the components are, starting from the east (left): 20 mJy (21 mJy - snapshot 10 days after), 13 mJy (10 mJy), 50 mJy (45 mJy), 66 mJy (60 mJy); the spectral indexes are: $\alpha_{1.6-5} = 0.32$ and $\alpha_{5-8.4} = 0.95$, $\alpha_{1.6-5} = -0.30$ and $\alpha_{5-8.4} = -0.87$, $\alpha_{1.6-5} = -0.65$ and $\alpha_{5-8.4} = 0.42$, $\alpha_{1.6-5} = 0.03$ and $\alpha_{5-8.4} = 0.78$. The optical identification (POSS) - 19^m elliptical red “blob” (optical counter-parts?) - is only 21° misaligned with the NE-SW radio-alignment.

Acknowledgements: P. Augusto is supported by a grant from JNICT/EC PRAXIS XXI.

References

- Glazebrook, K., Ellis, R., Colless, M.M., Broadhurst, T.J., Allington-Smith, J., & Tanvir, N., 1995, MNRAS, 273, 157
Griffiths, R.E., Ratnatunga, K.U., Neuschaefer, L.W., Casertano, S., et al., 1994, ApJ, 437, 67
Lilly, S.J., 1993, ApJ, 411, 501
Patnaik, A.R., Browne, I.W.A., Wilkinson, P.N., & Wrobel, J.M., 1992, MNRAS, 254, 655
Smail, I., Ellis, R.S., & Fitchett, M.J., 1994, MNRAS, 270, 245
Surdej, J., Claeskens, J.F., Crampton, D., Filipenko, A.V., et al., 1993, AJ, 105, 2064
Turner, E.L., Ostriker, J.P., Gott, J.R., 1984, ApJ, 284, 1