



## ETIOLOGY

### ORIGINAL RESEARCH PAPER

## Mental disorders in pregnancy and 5–8 years after delivery

P. H. C. Rondó<sup>1\*</sup>, R. F. Ferreira<sup>1</sup>, J. O. Lemos<sup>1</sup> and J. A. Pereira-Freire<sup>2</sup>

<sup>1</sup>Department of Nutrition, School of Public Health, University of São Paulo, Avenida Dr Arnaldo 715, São Paulo, CEP: 01246-904, Brazil

<sup>2</sup>Federal University of Piauí, Campus Senador Helvídio Nunes de Barros, Rua Cícero Eduardo s/n, Bairro Junco, Picos PI, CEP: 64600-000, Brazil

*Global Mental Health* (2016), 3, e31, page 1 of 8. doi:10.1017/gmh.2016.26

**Background.** Even though mental disorders represent a major public health problem for women and respective children, there remains a lack of epidemiological longitudinal studies to assess the psychological status of women throughout pregnancy and later in life. This epidemiological cohort study assessed the relationship between mental disorders of 409 Brazilian women in pregnancy and 5–8 years after delivery.

**Methods.** The women were followed from 1997 to 2000 at 17 health services, and subsequently from 2004 to 2006 at their homes. Mental disorders were investigated by the Perceived Stress Scale-PSS, General Health Questionnaire-GHQ and State-Trait Anxiety Inventories-STAI. The relationship between scores of the PSS, GHQ and STAI 5–8 years after delivery and in pregnancy was assessed by multivariate linear regression analysis, controlling for the following confounders: maternal age, education, *per capita* income, family size, work, marital status and body mass index.

**Results.** Scores of the PSS, GHQ and STAI 5–8 years after delivery were positively associated with scores of the PSS, GHQ and STAI in the three trimesters of pregnancy, and inversely associated with maternal age and *per capita* income (adj.  $R^2$  varied from 0.15 to 0.37). PSS, GHQ and STAI scores in the 3rd trimester of pregnancy were positively associated with scores of the PSS, GHQ and STAI in the 1st and 2nd trimesters of pregnancy (adj.  $R^2$  varied from 0.31 to 0.65).

**Conclusions.** The results of this study reinforce the urgency to integrate mental health screening into routine primary care for pregnant and postpartum women.

Received 13 May 2016; Revised 14 September 2016; Accepted 8 October 2016

**Key words:** Anxiety, maternal depression, postpartum, pregnancy, stress.

### Introduction

According to the WHO World Mental Health (WMH) Surveys (Kessler *et al.* 2015) involving 24 countries, approximately half of the individuals with major depressive disorder (MDD) had previously anxiety disorders. Women and formerly married people were the groups most at risk for elevated rates of MDD associated with anxiety disorders.

It is well recognized that during women's life pregnancy and postpartum are very sensitive periods, predisposing them to mental disorders like anxiety and depression (Howard *et al.* 2014). Mental disorders in pregnancy have been associated with prematurity and low birth weight (Rondó *et al.* 2003; Rondó, 2007; Grote *et al.* 2010). When present in the postpartum period, these disorders can have a negative impact on the mental health of the children (Buss *et al.* 2011; Davis & Sandman, 2012; Loomans *et al.* 2012; Ali *et al.* 2013; Rouse & Goodman, 2014), and even a few years later in life, they may interfere in the nutritional status of the children (Dunkel Schetter, 2011; Ali *et al.* 2013; Rondó *et al.* 2013).

\* Address for correspondence: Professor P. H. C. Rondó, Department of Nutrition, School of Public Health, University of São Paulo, Avenida Dr Arnaldo 715, São Paulo CEP: 01246-904, Brazil. (Email: [phcrondo@usp.br](mailto:phcrondo@usp.br))



Mental disorders, if firstly observed in pregnancy and in the postpartum periods, may be transitional, but can also persist throughout life (Kim *et al.* 2008; Mora *et al.* 2009; Sutter-Dallay *et al.* 2012; Kuo *et al.* 2014), bringing several problems for the women and the whole family.

Even though mental disorders represent a major public health problem for women and respective children, there remains a lack of epidemiological longitudinal studies to assess the psychological status of women throughout pregnancy and later in life. Most studies emphasized the pregnancy and/or the immediate post-partum periods (Haim *et al.* 2014; Radoš *et al.* 2015), but did not follow women years after delivery.

Therefore, the objective of this study was to follow women throughout pregnancy up to 5–8 years after delivery to assess the relationship between mental disorders in the three trimesters of pregnancy and later in life.

## Methods

This epidemiological cohort study involved 409 women in Jundiai city, São Paulo state, Brazil. It is derived from a cohort, carried out between 1997 and 2000, initially composed of 865 low-income pregnant women from Jundiai, to evaluate stress and distress as predictors of low birth weight, prematurity and intrauterine growth retardation. The women were recruited from all health units and hospitals in the city and were followed before the 16th week of pregnancy to the birth of their babies. All women were insured by the National Health Service (SUS) and were apparently healthy, considering that those who present any problem in pregnancy are usually reported to specialized antenatal services. Women with chronic infectious diseases, metabolic diseases, cardiopathy, mental diseases, hypertension/pre-eclampsia/eclampsia, vaginal bleeding and multiple deliveries were excluded from the study. Details of the cohort have been published previously (Rondó *et al.* 2003; Rondó & Souza, 2007).

The second prospective cohort study was carried out between November 2004 and December 2006 to assess mental disorders 5–8 years after delivery, and consisted of two phases. In the first phase, information from the questionnaire of the first cohort study was taken into account, and the women who at that time were living in Jundiai and nearby municipalities were located. The women were invited to participate in the present study through telephone contact or by visiting their homes if they did not have a telephone. Next, a home visit was made, during which the objective of the study was explained, and the ethical consent form was signed by the women. At the time of this

home visit, a general questionnaire was applied in order to assess demographic and socioeconomic factors. In the second phase of the study, the participants were contacted again by telephone to arrange to collect anthropometric measurements and data on mental disorders.

Out of the 865 women from the previous cohort, 745 women were located and invited to participate in the study, resulting in a sample of 649 women who signed a free informed consent form and answered a general questionnaire. However, 240 of them had incomplete data or did not participate in the second phase of the study, resulting in a final sample of 409 women.

Demographic and socioeconomic characteristics of the women were evaluated by the same general questionnaire used in the original cohort study. Their nutritional status was determined by the body mass index (BMI) and classified according to WHO recommendations (WHO, 2000). The women were weighed on a portable Sohnle<sup>®</sup> electronic scale (Sohnle, model 7500, Murrhardt, Germany), with a precision of 100 g. Height was measured with a SECA<sup>®</sup> wall-mounted stadiometer (Leicester Portable Height measure model, Hamburg, Germany), with a precision of 0.1 cm. The anthropometric measurements were performed according to the recommendations of Jelliffe & Jelliffe (1989).

Mental disorders were assessed by four psychologists who interviewed the women 3 times in pregnancy (at a gestational age <16 weeks, from 20 to 26 weeks and from 30 to 36 weeks) and 5–8 years after delivery, using versions of the Perceived Stress Scale (PSS), the General Health Questionnaire (GHQ) and the State-Trait Anxiety Inventories (STAI) validated in Brazil, respectively, by Luft *et al.* (2007), Mari & Williams (1985) and Biaggio *et al.* (1977). The PSS (Cohen *et al.* 1983) determined the degree to which situations in the last month had been considered as stressful, on a five-point scale ranging from 'never' to 'very often'. A 12-item version of the GHQ (Mari & Williams, 1985) assessed mental disorders in general, and classified the individuals in two groups: with low, 0–3 and high,  $\geq 4$  scores. The STAI (Spielberger *et al.* 1970) assessed anxiety by a well standardized, 40 item, self-report instrument designed to measure both state and trait anxiety. For State Anxiety (SA), subjects were asked how they felt at the time of being questioned, and for Trait Anxiety (TA), subjects were asked how they felt generally. A cut-off point greater than 40 was selected for both SAI and TAI.

The Stata version 10 software (College Station, TX, USA) was used for storage and statistical analysis of the data. The relationship between scores of the PSS, GHQ and STAI 5–8 years postpartum (dependent variables) and in the three trimesters of pregnancy (independent variables) was assessed by multivariate



linear regression analysis, using the backward stepwise selection method. The relationship between scores of the PSS, GHQ and STAI in the 3rd trimester of pregnancy (dependent variables) and in the 1st and 2nd trimesters (independent variables) was also assessed. The following confounding factors were included in the models: maternal age, education, *per capita* income, family size, work, marital status and BMI. A *p* value  $\leq 0.05$  was considered as statistically significant.

## Results

There were losses of 33.6% and 42.7% of the sample, respectively, considering all the mothers who were located ( $n=745$ ) in the first phase of the study, and the ones who participated in the original cohort ( $n=865$ ). Comparison of the characteristics between the mothers included in the cohort and those who did not conclude the study showed no significant differences.

Table 1 presents the characteristics of the women included in the study. Considering that there is no clear cut-off point for the scores of the PSS, they are presented in tertiles. High scores of the GHQ, SAI and TAI in the three trimesters of pregnancy and 5–8 years after delivery varied from 23.3% to 58.4%.

Table 2 shows four linear regression models considering scores of the PSS, GHQ, SAI and TAI 5–8 years after delivery as outcomes, and scores of the PSS, GHQ, SAI e TAI in the three trimesters of pregnancy as independent variables, controlling for confounders. Scores of the PSS 5–8 years after delivery were positively associated with scores of the PSS in the 1st and 3rd trimesters of pregnancy, and inversely associated with maternal age and *per capita* income (adj.  $R^2=0.21$ ). Scores of the GHQ 5–8 years after delivery were positively associated with scores of the GHQ in the three trimesters of pregnancy, and inversely associated with maternal age and *per capita* income (adj.  $R^2=0.18$ ). Scores of the SAI 5–8 years after delivery were positively associated with scores of the SAI in the 2nd and 3rd trimesters of pregnancy, and inversely associated with *per capita* income (adj.  $R^2=0.15$ ). Scores of the TAI 5–8 years after delivery were positively associated with scores of the TAI in the 1st and 2nd trimesters of pregnancy, and inversely associated with maternal age and *per capita* income (adj.  $R^2=0.37$ ).

Table 3 shows four linear regression models considering scores of the PSS, GHQ, SAI and TAI in the 3rd trimester of pregnancy as outcomes, and scores of the PSS, GHQ, SAI e TAI in the 1st and 2nd trimesters of pregnancy as independent variables, controlling for confounders. Scores of the PSS, GHQ, SAI and TAI in the 3rd trimester of pregnancy were positively associated with scores of the PSS, GHQ, SAI and TAI in the

1st and 2nd trimesters of pregnancy (adj.  $R^2$  varied from 0.31 to 0.65).

## Discussion

According to the results of this study, there were statistically significant associations between mental disorders (assessed by the PSS, GHQ and STAI scores) in the three trimesters of pregnancy, lower age and *per capita* income and mental disorders 5–8 years after delivery. The other socioeconomic and demographic factors investigated did not show statistically significant results. BMI, an indicator of the nutritional status of the women, was not associated with mental disorders; though Nagl *et al.* (2015) and Molyneaux *et al.* (2016) had referred that obese pregnant women might constitute a group vulnerable for anxiety and depression, respectively.

Apparently, women have a higher risk of mental disorders in the postpartum period if they have shown these disorders before (Bilszta *et al.* 2008; Witt *et al.* 2011; Kirkan *et al.* 2015; Patton *et al.* 2015) or during pregnancy (Onoye *et al.* 2013; Yazici *et al.* 2015). Bilszta *et al.* (2008) observed an association between history of psychopathology in the antenatal period and depression at 6–8 weeks postnatally in Australian women. Patton *et al.* (2015) observed that perinatal depressive symptoms were mostly preceded by mental health problems that began before pregnancy, in adolescence or young adulthood. Onoye *et al.* (2013) confirmed the associations between stress, depression and anxiety in the three trimesters of pregnancy and in early ( $<6$  weeks after delivery) and late ( $\geq 6$  weeks after delivery) postpartum in women from Hawaii. Kirkan *et al.* (2015) and Yazici *et al.* (2015) reported that a previous mental disorder or an untreated depressive disorder in the 1st trimester of pregnancy was an important predictor of depression in the sixth postpartum week.

Comparing the mean scores of the women in the highest PSS tertile with the mean scores of the whole population, and the mean scores of the populations studied by Cohen *et al.* (1983), we concluded that the women included in our study had perceived stress in pregnancy and 5–8 years after delivery. Interestingly, PSS mean scores increased significantly (*t* test,  $P < 0.001$ ; data not shown) 5–8 years after delivery compared with the mean scores in pregnancy, while the GHQ, SAI and TAI scores maintained the mean values in the four different periods investigated, indicating higher levels of perceived stress 5–8 years after delivery than in pregnancy. Schmied *et al.* (2013) observed that women's mood appears to be better in the first year after birth, when compared with a specific moment in pregnancy and 5 years later. The proportion of women reporting depressive symptoms in the



**Table 1.** Characteristics of the women included in the study (n = 409)

|                                      | N   | %    | Mean (s.d.) |
|--------------------------------------|-----|------|-------------|
| Age (years)                          |     |      | 30.7 (6.06) |
| Education (years) <sup>a</sup>       |     |      | 7.7 (3.1)   |
| Per capita income (MBW) <sup>b</sup> |     |      | 0.92 (0.67) |
| Family size (number of persons)      |     |      | 4.3 (1.4)   |
| Number of children                   |     |      | 1.8 (0.97)  |
| 1                                    | 209 | 51.1 |             |
| 2                                    | 116 | 28.4 |             |
| 3                                    | 61  | 14.9 |             |
| ≥4                                   | 23  | 5.6  |             |
| Work                                 |     |      |             |
| Yes                                  | 114 | 27.9 |             |
| No                                   | 295 | 72.1 |             |
| Marital status                       |     |      |             |
| Married/with partner                 | 313 | 76.5 |             |
| Others                               | 96  | 23.5 |             |
| BMI (weight/height <sup>2</sup> )    |     |      | 27.3 (4.1)  |
| 18.5–24.9 (normal)                   | 124 | 30.3 |             |
| 25.0–29.9 (overweight)               | 201 | 49.1 |             |
| 30.0–39.9 (obese I and II)           | 84  | 20.6 |             |
| 1st trimester                        |     |      |             |
| PSS                                  |     |      | 22.7 (6.6)  |
| 7–20                                 | 138 | 33.6 |             |
| 21–25                                | 125 | 30.4 |             |
| 26–43                                | 147 | 36.0 |             |
| GHQ                                  |     |      | 2.8 (3.1)   |
| ≤3                                   | 273 | 66.7 |             |
| ≥4                                   | 136 | 33.3 |             |
| SAI                                  |     |      | 37.3 (8.8)  |
| <40                                  | 282 | 68.9 |             |
| ≥40                                  | 127 | 31.1 |             |
| TAI                                  |     |      | 41.8 (10.6) |
| <40                                  | 199 | 48.7 |             |
| ≥40                                  | 210 | 51.3 |             |
| 2nd trimester                        |     |      |             |
| PSS                                  |     |      | 22.8 (6.9)  |
| 4–19                                 | 132 | 32.3 |             |
| 20–25                                | 138 | 33.6 |             |
| 26–46                                | 140 | 34.1 |             |
| GHQ                                  |     |      | 2.3 (2.6)   |
| ≤3                                   | 303 | 74.1 |             |
| ≥4                                   | 106 | 25.9 |             |
| SAI                                  |     |      | 35.9 (7.9)  |
| <40                                  | 314 | 76.7 |             |
| ≥40                                  | 95  | 23.3 |             |
| TAI                                  |     |      | 40 (9.9)    |
| <40                                  | 227 | 55.6 |             |
| ≥40                                  | 182 | 44.4 |             |
| 3rd trimester                        |     |      |             |
| PSS                                  |     |      | 22.1 (7.2)  |
| 2–19                                 | 147 | 35.8 |             |
| 20–25                                | 129 | 31.6 |             |
| 26–53                                | 134 | 32.6 |             |

(Continued)

**Table 1 (cont.)**

|                       | N   | %    | Mean (s.d.) |
|-----------------------|-----|------|-------------|
| GHQ                   |     |      | 2.5 (2.7)   |
| ≤3                    | 304 | 74.3 |             |
| ≥4                    | 105 | 25.7 |             |
| SAI                   |     |      | 36.8 (8.4)  |
| <40                   | 302 | 73.8 |             |
| ≥40                   | 107 | 26.2 |             |
| TAI                   |     |      | 39.5 (9.8)  |
| <40                   | 224 | 54.8 |             |
| ≥40                   | 185 | 45.2 |             |
| 5–8 years post-partum |     |      |             |
| PSS                   |     |      | 38.1 (7.4)  |
| 15–35                 | 143 | 34.8 |             |
| 36–41                 | 136 | 33.3 |             |
| 42–55                 | 131 | 31.9 |             |
| GHQ                   |     |      | 3.2 (3.2)   |
| ≤3                    | 252 | 61.6 |             |
| ≥4                    | 157 | 38.4 |             |
| SAI                   |     |      | 40.2 (10.1) |
| <40                   | 228 | 55.7 |             |
| ≥40                   | 181 | 44.3 |             |
| TAI                   |     |      | 43.7 (10.8) |
| <40                   | 170 | 41.6 |             |
| ≥40                   | 239 | 58.4 |             |

<sup>a</sup> n = 406.

<sup>b</sup> Minimum Brazilian Wage – MBW (1 MBW = R\$350.00 = approx. US\$77).

PSS, Perceived Stress Scale; GHQ, General Health Questionnaire; STAI, State-Trait Anxiety Inventories (SAI, State Anxiety Inventory; TAI-Trait Anxiety Inventory).

first year postpartum was between 10% and 20% and this remained stable over 25 years. However, the studies included in the meta-analysis carried out by Schmieid *et al.* (2013) did not assess mental health in the three trimesters of pregnancy. Agrati *et al.* (2015) found that anxiety followed a U-shaped pattern from pregnancy to 2 years postpartum, which was modified by early life experience of the women. Greater early adversity was associated with higher anxiety in pregnancy, followed by a marked decrease once the baby was born, and a subsequent increase during the later postpartum period.

Similar to other studies in the literature, lower age (Bottino *et al.* 2012; Räisänen *et al.* 2014; Siegel & Brandon, 2014) and inferior *per capita* income (Abdollahi *et al.* 2014; Hein *et al.* 2014; Alfayumi-Zeadna *et al.* 2015) were associated with mental disorders in the postpartum period. According to Bottino *et al.* (2012), maternal age was significantly associated with PPD. For each additional year, a reduction of 4% in the chance of developing PPD was anticipated, effect that was not modified by confounders.



**Table 2.** Linear regression models considering scores of the PSS, GHQ, SAI and TAI 5–8 years after delivery as outcomes

| Outcomes                         | Coefficient | S.E. | T     | 95% CI               |                       | p      |
|----------------------------------|-------------|------|-------|----------------------|-----------------------|--------|
| PSS (5–8 years after delivery)   |             |      |       |                      |                       |        |
| PSS (1st trimester)              | 0.20        | 0.08 | 2.54  | 0.05                 | 0.36                  | 0.01   |
| PSS (2nd trimester)              | 0.13        | 0.09 | 1.50  | –0.04                | 0.31                  | 0.14   |
| PSS (3rd trimester)              | 0.19        | 0.08 | 2.35  | 0.03                 | 0.35                  | 0.02   |
| Maternal age                     | –0.15       | 0.07 | –2.24 | –0.29                | –0.02                 | 0.03   |
| Per capita income                | –0.67       | 0.23 | –2.86 | –1.13                | –0.21                 | 0.005  |
| $R^2 = 0.23$ ; Adj. $R^2 = 0.21$ |             |      |       |                      |                       |        |
| GHQ (5–8 years after delivery)   |             |      |       |                      |                       |        |
| GHQ (1st trimester)              | 0.27        | 0.06 | 4.14  | 0.14                 | 0.39                  | <0.001 |
| GHQ (2nd trimester)              | 0.12        | 0.07 | 1.67  | –0.02                | 0.27                  | 0.09   |
| GHQ (3rd trimester)              | 0.14        | 0.07 | 2.10  | $0.9 \times 10^{-2}$ | 0.28                  | 0.04   |
| Maternal age                     | –0.07       | 0.03 | –2.78 | –0.12                | –0.02                 | 0.006  |
| Per capita income                | –0.24       | 0.09 | –2.55 | –0.42                | –0.05                 | 0.01   |
| $R^2 = 0.19$ ; Adj. $R^2 = 0.18$ |             |      |       |                      |                       |        |
| SAI (5–8 years after delivery)   |             |      |       |                      |                       |        |
| SAI (1st trimester)              | 0.10        | 0.06 | 1.59  | –0.02                | 0.23                  | 0.11   |
| SAI (2nd trimester)              | 0.16        | 0.08 | 2.10  | 0.01                 | 0.31                  | 0.04   |
| SAI (3rd trimester)              | 0.28        | 0.07 | 4.18  | 0.15                 | 0.42                  | <0.001 |
| Per capita income                | –0.84       | 0.29 | –2.92 | –1.41                | –0.27                 | 0.004  |
| $R^2 = 0.16$ ; Adj. $R^2 = 0.15$ |             |      |       |                      |                       |        |
| TAI (5–8 years after delivery)   |             |      |       |                      |                       |        |
| TAI (1st trimester)              | 0.29        | 0.06 | 5.00  | 0.18                 | 0.41                  | <0.001 |
| TAI (2nd trimester)              | 0.25        | 0.07 | 3.62  | 0.12                 | 0.39                  | <0.001 |
| TAI (3rd trimester)              | 0.12        | 0.08 | 1.53  | –0.03                | 0.26                  | 0.13   |
| Maternal age                     | –0.15       | 0.07 | –2.10 | –0.29                | $-0.9 \times 10^{-2}$ | 0.04   |
| Per capita income                | –0.90       | 0.27 | –3.34 | –1.43                | –0.37                 | 0.001  |
| $R^2 = 0.38$ ; Adj. $R^2 = 0.37$ |             |      |       |                      |                       |        |

PSS, Perceived Stress Scale; GHQ, General Health Questionnaire; SAI, State Anxiety Inventory; TAI-Trait Anxiety Inventory.

**Table 3.** Linear regression models considering scores of the PSS, GHQ, SAI and TAI in the 3rd trimester of pregnancy as outcomes

| Outcomes                         | Coefficient | S.E. | T     | 95% CI |      | p      |
|----------------------------------|-------------|------|-------|--------|------|--------|
| PSS (3rd trimester)              |             |      |       |        |      |        |
| PSS (1st trimester)              | 0.27        | 0.06 | 4.47  | 0.15   | 0.39 | <0.001 |
| PSS (2nd trimester)              | 0.54        | 0.06 | 8.95  | 0.42   | 0.66 | <0.001 |
| $R^2 = 0.47$ ; Adj. $R^2 = 0.47$ |             |      |       |        |      |        |
| GHQ (3rd trimester)              |             |      |       |        |      |        |
| GHQ (1st trimester)              | 0.20        | 0.05 | 4.38  | 0.11   | 0.29 | <0.001 |
| GHQ (2nd trimester)              | 0.49        | 0.05 | 10.17 | 0.39   | 0.59 | <0.001 |
| $R^2 = 0.35$ ; Adj. $R^2 = 0.35$ |             |      |       |        |      |        |
| SAI (3rd trimester)              |             |      |       |        |      |        |
| SAI (1st trimester)              | 0.23        | 0.05 | 4.97  | 0.14   | 0.33 | <0.001 |
| SAI (2nd trimester)              | 0.44        | 0.05 | 8.31  | 0.33   | 0.54 | <0.001 |
| $R^2 = 0.32$ ; Adj. $R^2 = 0.31$ |             |      |       |        |      |        |
| TAI (3rd trimester)              |             |      |       |        |      |        |
| TAI (1st trimester)              | 0.31        | 0.36 | 8.72  | 0.24   | 0.38 | <0.001 |
| TAI (2nd trimester)              | 0.56        | 0.04 | 14.54 | 0.48   | 0.63 | <0.001 |
| $R^2 = 0.65$ ; Adj. $R^2 = 0.65$ |             |      |       |        |      |        |

PSS, Perceived Stress Scale; GHQ, General Health Questionnaire; SAI, State Anxiety Inventory; TAI-Trait Anxiety Inventory.



Räisänen *et al.* (2014) referred that one of the risk profiles of major depression included adolescence or advanced maternal age. Primarily focusing upon depressive symptoms, findings from a large literature review showed that rates of depression in pregnant and postpartum adolescents are comparable with non-pregnant adolescents, but higher than those reported among pregnant adults (Siegel & Brandon, 2014). In fact, Aras *et al.* (2013) showed that early maternal age is one of the risk factors for depressive disorders even in non-perinatal reproductive age.

Abdollahi *et al.* (2014) identified low household income as the predictive factor with the highest risk (OR 3.57, 95% CI 1.49–8.5) for PPD. Hein *et al.* (2014) concluded that socioeconomic factors, including monthly income, define subgroups that have different depression scores during and after pregnancy. Alfayumi-Zeadna *et al.* (2015), using a validated Arabic translation of the 'Edinburgh Postnatal Depression Scale' found that low income was one of the socio-demographic factors associated with PPD, among Arab-Bedouin women.

A limitation of this study was the lack of assessment of mental disorders in the immediate post-partum period. Even though mental disorders had not been investigated in the immediate postpartum period, it was probably a problem for those pregnant women considering the associations of the PSS, GHQ and STAI scores between the 3rd and 1st and 2nd trimesters of pregnancy. Witt *et al.* (2011) referred that poor pre-pregnancy mental health and poor antepartum mental health, both independently, increased the risk of postpartum mental health problems in North American women. However, the authors did not use specific questionnaires to assess mental disorders. They used self-reports and symptoms of mental health conditions or global mental health ratings of 'fair' or 'poor'.

## Conclusion

Mental disorders in the three trimesters of pregnancy, lower age and *per capita* income are associated with mental disorders 5–8 years after delivery.

Referral of young, low-income pregnant women with mental disorders, particularly anxiety and depression, should be encouraged to psychological or psychiatric treatment. Finally, we reinforce the urgency to integrate mental health screening into routine primary care for pregnant and postpartum women.

## Acknowledgements

The authors acknowledge the field workers, particularly Alessandra L Santos and Josimara F Moura, and Fundação de Amparo à Pesquisa do Estado de São

Paulo – FAPESP, Brazil (grants no. 1998/00321-0, no. 2004/04109-8).

## Declaration of interest

All the authors declare no conflict of interest.

## Ethical standards

The study was approved by the Ethics Committee of the School of Public Health, University of Sao Paulo. It was carried out in accordance with the Declaration of Helsinki of 1975, as revised in 2008.

## References

- Abdollahi F, Zarghami M, Azhar MZ, Sazlina SG, Lye MS (2014) Predictors and incidence of post-partum depression: a longitudinal cohort study. *Journal of Obstetrics and Gynaecology Research* **40**, 2191–2200.
- Agrati D, Browne D, Jonas W, Meaney M, Atkinson L, Steiner M, Fleming AS, MAVAN Research Team (2015). Maternal anxiety from pregnancy to 2 years postpartum: transactional patterns of maternal early adversity and child temperament. *Archives of Women's Mental Health* **18**, 693–705.
- Alfayumi-Zeadna S, Kaufman-Shriqui V, Zeadna A, Lauden A, Shoham-Vardi I (2015). The association between sociodemographic characteristics and postpartum depression symptoms among Arab-Bedouin women in Southern Israel. *Depression and Anxiety* **32**, 120–128.
- Ali NS, Mahmud S, Khan A, Ali BS (2013). Impact of postpartum anxiety and depression on child's mental development from two peri-urban communities of Karachi, Pakistan: a quasi-experimental study. *BMC Psychiatry* **13**, 274.
- Aras N, Oral E, Aydin N, Gulec M (2013). Maternal age and number of children are risk factors for depressive disorders in non-perinatal women of reproductive age. *International Journal of Psychiatry in Clinical Practice* **17**, 298–306.
- Biaggio AMB, Natalicio L, Spielberger CD (1977). Desenvolvimento da forma experimental em português do Inventário de Ansiedade Traço-Estado (IDATE) de Spielberger. *Arquivos Brasileiros de Psicologia Aplicada* **29**, 31–44.
- Bilszta JL, Gu YZ, Meyer D, Buist AE (2008). A geographic comparison of the prevalence and risk factors for postnatal depression in an Australian population. *Australian and New Zealand Journal of Public Health* **32**, 424–430.
- Bottino MN, Nadanovsky P, Moraes CL, Reichenheim ME, Lobato G (2012). Reappraising the relationship between maternal age and postpartum depression according to the evolutionary theory: empirical evidence from a survey in primary health services. *Journal of Affective Disorders* **142**, 219–224.
- Buss C, Davis EP, Hobel CJ, Sandman CA (2011). Maternal pregnancy-specific anxiety is associated with child executive function at 6–9 years age. *Stress* **14**, 665–676.



- Cohen S, Kamarck T, Mermelstein R** (1983). A global measure of perceived stress. *Journal of Health and Social Behavior* **24**, 385–396.
- Davis EP, Sandman CA** (2012) Prenatal psychobiological predictors of anxiety risk in preadolescent children. *Psychoneuroendocrinology* **37**, 1224–1233.
- Dunkel Schetter C** (2011) Psychological science on pregnancy: stress processes, biopsychosocial models, and emerging research issues. *Annual Review of Psychology* **62**, 531–558.
- Grote NK, Bridge JA, Gavin AR, Melville JL, Iyengar S, Katon WJ** (2010) A meta-analysis of depression during pregnancy and the risk of preterm birth, low birth weight, and intrauterine growth restriction. *Archives of General Psychiatry* **67**, 1012–1024.
- Haim A, Sherer M, Leuner B** (2014) Gestational stress induces persistent depressive-like behavior and structural modifications within the postpartum nucleus accumbens. *European Journal of Neuroscience* **40**, 3766–3773.
- Hein A, Rauh C, Engel A, Häberle L, Dammer U, Voigt F, Fasching PA, Faschingbauer F, Burger P, Beckmann MW, Kornhuber J, Goecke TW** (2014). Socioeconomic status and depression during and after pregnancy in the Franconian Maternal Health Evaluation Studies (FRAMES). *Archives of Gynecology and Obstetrics* **289**, 755–763.
- Howard LM, Molyneaux E, Dennis CL, Rochat T, Stein A, Milgrom J** (2014). Non-psychotic mental disorders in the perinatal period. *Lancet* **384**, 1775–1788.
- Jelliffe DB, Jelliffe EFP** (1989). *Community Nutritional Assessment, with Special Reference to Less Technically Developed Countries*, 2nd edn. Oxford University Press: London.
- Kessler RC, Sampson NA, Berglund P, Gruber MJ, Al-Hamzawi A, Andrade L, Bunting B, Demyttenaere K, Florescu S, de Girolamo G, Gureje O, He Y, Hu C, Huang Y, Karam E, Kovess-Masfety V, Lee S, Levinson D, Medina Mora ME, Moskalewicz J, Nakamura Y, Navarro-Mateu F, Browne MA, Piazza M, Posada-Villa J, Slade T, Ten Have M, Torres Y, Vilagut G, Xavier M, Zarkov Z, Shahly V, Wilcox MA** (2015). Anxious and non-anxious major depressive disorder in the World Health Organization World Mental Health Surveys. *Epidemiology and Psychiatric Sciences* **24**, 210–226.
- Kim YK, Hur JW, Kim KH, Oh KS, Shin YC** (2008). Prediction of postpartum depression by sociodemographic, obstetric and psychological factors: a prospective study. *Psychiatry and Clinical Neurosciences* **62**, 331–340.
- Kirkan TS, Aydin N, Yazici E, Aslan PA, Acemoglu H, Daloglu AG** (2015). The depression in women in pregnancy and postpartum period: a follow-up study. *International Journal of Social Psychiatry* **61**, 343–349.
- Kuo SY, Chen SR, Tzeng YL** (2014) Depression and anxiety trajectories among women who undergo an elective cesarean section. *PLoS ONE* **9**, e86653.
- Loomans EM, van der Stelt O, van Eijsden M, Gemke RJ, Vrijkotte TG, Van den Bergh BR** (2012) High levels of antenatal maternal anxiety are associated with altered cognitive control in five-year-old children. *Developmental Psychobiology* **54**, 441–450.
- Luft CD, Sanches SdeO, Mazo GZ, Andrade A** (2007) Brazilian version of the Perceived Stress Scale: translation and validation for the elderly. *Revista de Saude Publica* **41**, 606–615.
- Mari JJ, Williams P** (1985) A comparison of the validity of two psychiatric screening questionnaires (GHQ-12 and SRQ-20) in Brazil, using relative operating characteristic (ROC) analysis. *Psychological Medicine* **15**, 651–659.
- Molyneaux E, Poston L, Khondoker M, Howard LM** (2016). Obesity, antenatal depression, diet and gestational weight gain in a population cohort study. *Archives of Women's Mental Health* (<http://link.springer.com/article/10.1007%2Fs00737-016-0635-3>). Accessed 14 September 2016.
- Mora PA, Bennett IM, Elo IT, Mathew L, Coyne JC, Culhane JF** (2009). Distinct trajectories of perinatal depressive symptomatology: evidence from growth mixture modeling. *American Journal of Epidemiology* **169**, 24–32.
- Nagl M, Linde K, Stepan H, Kersting A** (2015). Obesity and anxiety during pregnancy and postpartum: a systematic review. *Journal of Affective Disorders* **186**, 293–305.
- Onoye JM, Shafer LA, Goebert DA, Morland LA, Matsu CR, Hamagami F** (2013). Changes in PTSD symptomatology and mental health during pregnancy and postpartum. *Archives of Women's Mental Health* **16**, 453–463.
- Patton GC, Romaniuk H, Spry E, Coffey C, Olsson C, Doyle LW, Oats J, Hearps S, Carlin JB, Brown S** (2015). Prediction of perinatal depression from adolescence and before conception (VIHCS): 20-year prospective cohort study. *Lancet* **386**, 875–883.
- Radoš SN, Herman R, Tadinac M** (2015). Is the predictability of new-onset postpartum depression better during pregnancy or in the early postpartum period? A prospective study in Croatian women. *Health Care for Women International* **3**, 1–22.
- Räisänen S, Lehto SM, Nielsen HS, Gissler M, Kramer MR, Heinonen S** (2014). Risk factors for and perinatal outcomes of major depression during pregnancy: a population-based analysis during 2002–2010 in Finland. *BMJ Open* **4**, e004883.
- Rondó P, Souza MR** (2007). Maternal distress and intended breastfeeding duration. *Journal of Psychosomatic Obstetrics and Gynecology* **28**, 55–60.
- Rondó PH, Ferreira RF, Nogueira F, Ribeiro MC, Lobert H, Artes R** (2003). Maternal psychological stress and distress as predictors of low birth weight, prematurity and intrauterine growth retardation. *European Journal of Clinical Nutrition* **57**, 266–272.
- Rondó PH, Rezende G, Lemos JO, Pereira JA** (2013). Maternal stress and distress and child nutritional status. *European Journal of Clinical Nutrition* **67**, 348–352.
- Rondó PHC** (2007). Maternal stress/distress and low birth weight, preterm birth and intrauterine growth restriction – a review. *Current Women's Health Reviews* **3**, 13–29.
- Rouse MH, Goodman SH** (2014). Perinatal depression influences on infant negative affectivity: timing, severity, and co-morbid anxiety. *Infant Behavior and Development* **37**, 739–751.
- Schmied V, Johnson M, Naidoo N, Austin MP, Matthey S, Kemp L, Mills A, Meade T, Yeo A** (2013). Maternal mental health in Australia and New Zealand: a



- review of longitudinal studies. *Women and Birth* **26**, 167–178.
- Siegel RS, Brandon AR** (2014). Adolescents, pregnancy, and mental health. *Journal of Pediatric and Adolescent Gynecology* **27**, 138–150.
- Spielberger CD, Gorsuch RI, Lushene RE** (1970). *STAI Manual for the State – Trait Anxiety Inventory*. Consulting Psychologist Press: Palo Alto, CA.
- Sutter-Dallay AL, Cosnefroy O, Glatigny-Dallay E, Verdoux H, Rasclé N** (2012). Evolution of perinatal depressive symptoms from pregnancy to two years postpartum in a low-risk sample: the MATQUID cohort. *Journal of Affective Disorders* **139**, 23–29.
- Witt WP, Wisk LE, Cheng ER, Hampton JM, Creswell PD, Hagen EW, Spear HA, Maddox T, Deleire** (2011). Poor prepregnancy and antepartum mental health predicts postpartum mental health problems among US women: a nationally representative population-based study. *Women's Health Issues* **21**, 304–313.
- World Health Organization– WHO** (2000) *Consultation on Obesity. Obesity: Preventing and Managing the Global Epidemic*. WHO: Geneva, Switzerland.
- Yazici E, Kirkan TS, Aslan PA, Aydin N, Yazici AB** (2015). Untreated depression in the first trimester of pregnancy leads to postpartum depression: high rates from a natural follow-up study. *Neuropsychiatric Disease and Treatment* **11**, 405–411.