

Results: A total of 100 individuals participated, of whom, 80 were women and 20 were men. The median scores in HADS was 16 and the interquartile range was 12. Multilevel analysis shows that better physical functioning (sit to stand test) and worse sleep quality (Insomnia severity index) are predictors of worse affective state. The models explain 36.5% of the HADS variance.

Conclusions: It is relevant to take account these variables in the treatment of the affective state of patients with long covid.

Disclosure of Interest: None Declared

EPV0294

Evolution of the affective state of a cohort of people suffering from long covid and associated factors

B. Oliván-Blázquez^{1*}, S. León-Herrera¹, M. Samper-Pardo¹, F. Méndez-López², A. Aguilar-Latorre² and M. Martínez-Perrochan²

¹University of Zaragoza and ²Institute for Health Research Aragón, Zaragoza, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1643

Introduction: Long COVID patients have experienced a decline in their quality of life caused, in part but not wholly, by its negative emotional impact. Some of the most prevalent mental symptoms presented by Long COVID patients are anxiety, depression and sleep disorders.

Objectives: The objective of this study is to increase understanding of the affective state of people diagnosed with Long COVID, the evolution and associated factors.

Methods: Longitudinal study of three months of duration. The study population was 100 post-COVID-19 patients aged 18 years or older (80 women and 20 men). The main variable was the affective state through the Hospital Anxiety and Depression Scale (HADS) questionnaire. The rest of the collected variables were: Socio-demographic variables, number of residual symptoms, cognitive functioning using the Montreal Cognitive Assessment (MoCA), physical functioning variable measured by Sit to Stand Test and Sleep quality through the Insomnia Severity Index (ISI). A statistical analysis comparing baseline and 3months follow up measures were performed, using a Student T for related samples statistical. A lineal regression analysing associated factors to a reduction in HADS score was also performed. Ethics approval was granted by the Clinical Research Ethics Committee of Aragón (PI21/139 and PI21/454).

Results: At baseline the score in anxiety, depression and total score were 9,10 (SD: 4,67), 8,25 (SD: 4,51) and 17,35 (SD: 8,43) respectively, and 74% of the participants were considered cases. At three months, there is a slightly decrease but not significative in the score of HADS, both in anxiety, depression and total score (pvalue 0,465; 0,236; and 0,216 respectively). 64,4% of the participants had a positive diagnosis of depression/anxiety. About the rest of the variables there were also a slight decrease but without being significant There was not a predictive model that explained the decrease in the HADS score.

Conclusions: The evolution of the people suffering long covid is very slow along the time, and also the affective state.

Disclosure of Interest: None Declared

EPV0295

PSYCHOLOGICAL IMPACT OF COVID-19 LOCKDOWN ON A POPULATION WITH SERIOUS MENTAL DISORDER: DIAGNOSTIC GROUP ANALYSIS

B. Pedruzo*, C. Aymerich, A. Catalan, M. Pacho, M. Bordenave, O. Estevez, J. Herrero, M. Laborda, G. Mancebo, J. L. Perez and M. A. Gonzalez Torres

¹Service of Psychiatry. Basurto University Hospital, Osakidetza, Bilbao, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1644

Introduction: Since its emergence at the end of 2019, the COVID-19 virus has spread worldwide. In Spain, mandatory home confinement was established on March 15, 2020, and lasted 99 days. Previous studies on events that required isolation situations suggest a worsening in the mental health of general population, and in particular, of especially vulnerable groups such as individuals with severe mental disorder (SMD).

Objectives: The aim of this study is to evaluate the psychological effect (anxiety and depression) of confinement in patients with SMD and to study the dissimilarities among the different diagnostic groups.

Methods: In this study, assessments were performed using the IDER and STAI questionnaires, in order to evaluate symptoms of depression and anxiety, respectively. The evaluations were carried out in patients who had required at least one admission to the Psychiatric Hospitalization Unit of the University Hospital of Basurto. The Shapiro-Wilk test was used to verify the normality of the sample. ANOVA test was used to study differences among diagnostic groups. Posteriorly, Bonferroni correction was performed.

Results: 95 participants completed the IDER questionnaire, obtaining a mean score of 24.56 (SD=8.18) for the state and 23.57 (SD=8.14) for the trait. In the STAI questionnaire, a mean score of 27.86 (SD=15.19) was obtained for the state and 30.49 (SD=14.71) for the trait. ANOVA test indicated presence of differences among groups. However, differences did not persist after Bonferroni correction.

Conclusions: Increased levels of anxiety and depression were found in the sample studied with respect to the general population. No statistically significant differences were found among different diagnostic groups. Further studies should be performed in order to increase the knowledge around this research area.

Disclosure of Interest: None Declared

EPV0296

A case report of post-acute COVID-19 encephalopathy

S. Cekerinac¹, D. Sekulic¹ and B. Batinic^{2,3*}

¹Department of Psychiatry, General Hospital, Sremska Mitrovica;

²Department of Psychology, University of Belgrade, Faculty of Philosophy and ³Clinic of Psychiatry, University Clinical Centre of Belgrade, Belgrade, Serbia

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.1645