

Policy and Systems Commentary

*The online version of this article has been updated since original publication. A notice detailing the change has also been published.

Cite this article: Böhm B, Palma M, Ousley J, Keane G (2022). Competency-based mental health supervision: evidence-based tool needs for the humanitarian context. *Global Mental Health* 9, 221–222. <https://doi.org/10.1017/gmh.2022.23>

Received: 26 November 2021

Revised: 21 March 2022

Accepted: 28 March 2022

First published online: 10 May 2022

Key words:

Clinical supervision; competency-based supervision; mhGAP; task-sharing; task-shifting

Author for correspondence:

Bettina Böhm,

E-mail: bettina.boehm@freenet.de

Competency-based mental health supervision: evidence-based tool needs for the humanitarian context

Bettina Böhm , Miguel Palma, Janet Ousley and Gregory Keane

Médecins Sans Frontières, Paris, France

The mental health (MH) care needs in humanitarian crises often far outweigh the local workforce capacity, making task-sharing to non-specialist MH staff essential to provide care for people with severe and common mental disorders (Murray and Jordans, 2016; Singla *et al.*, 2017; Patel *et al.*, 2018; Carreño *et al.*, 2019). Yet, task-sharing strategies cannot (and should not) replace professional accreditation models involving years of training. Clinical supervision of MH staff in task-sharing contexts is widely accepted as a way to guarantee quality of care; however, these activities are not consistently prioritized by MH policy makers who may focus on quick fixes such as one-off trainings without follow-up strategies including supervision, or on the immediate needs of the health care system over the needs of frontline care providers (Kemp *et al.*, 2019). In low-resource and humanitarian settings (LRHS), this means that practitioners with limited experience often care for high numbers of extremely vulnerable patients after time-limited training (Faregh *et al.*, 2019), with expert support only provided occasionally (Kemp *et al.*, 2019). Typical capacity building approaches have focused on ensuring fidelity to intervention manuals. The emphasis must shift, however, to building practical clinical skills over time.

Clinical supervision from a senior MH professional is critical for effective knowledge and skills transfer to novice practitioners (Holt *et al.*, 2015; Faregh *et al.*, 2019), thereby ensuring that they can provide evidence-based interventions safely. Although not replacing the imperative need to scale up psychosocial support to MH professionals, clinical supervision is also central in helping supervisees manage their own reactions to patients by addressing issues such as emotional over-involvement in or disengagement from a therapeutic process (Brockhouse *et al.*, 2011), which can harm patients. Protecting staff psychosocial well-being is particularly important in conflict, post-conflict, epidemic and disaster settings where they themselves are exposed to substantial and often ongoing suffering. Leaving their occupational stress unaddressed may lead to staff loss and high turnover, with consequent drops in care quality, shortages in and interruptions of essential services, which ultimately affects service users.

Successful supervision in LRHS is complicated by the fact that evidence-based practices have largely been developed for professionals in high-resource environments with none of the realities or constraints of humanitarian crises, which can include limited expert clinical care and supervision. Firstly, supervisory senior staff may be the only formally trained clinician for a large population, and any time spent supervising others can take away from their own patient or client services. Secondly, consistency of supervision is a challenge. Regular staff turnover is exacerbated by supervisors having different theoretical orientations, disparate supervision styles, and sometimes limited experience supervising others. A corollary of these factors, inadequate or poorly conducted supervision therefore results, for example, in decreased empathic engagement in patient care (Rønnestad *et al.*, 2019) or a reduction in regular opportunities for shared reflection and improvement, harm to clinicians and clinical malpractice. It is therefore crucial to ensure structure and consistency to supervision with concrete guidance and practical tools. These should be evidence-based (Holt *et al.*, 2015), conducive to learning (Kemp *et al.*, 2019), culturally appropriate (Heim and Kohrt, 2019) and context-adapted to different global expressions of suffering and forms of healing (Faregh *et al.*, 2019; Heim and Kohrt, 2019; Kohrt *et al.*, 2020a). Finally, co-design and field testing of these instruments may enable an easier institutionalization of competency-based supervision models.

Médecins sans Frontières (MSF), a non-profit humanitarian organization, prioritizes clinical supervision as an essential component of MH and psychosocial support activities in LRHS, building upon existing competency-based training approaches (that have been applied by MSF to the mhGAP intervention guide, e.g.). However, with respect to competency-based supervision, we have observed a dearth of available resources relevant to the contexts where we have operations. To fill this gap, MSF has designed tools to support professionals delivering supervision. Evidence-based competencies for provision of psychiatric, psychological or counselling care were initially identified (see research manuscript). Prioritization was given to common factors of the therapeutic alliance rather than to the technical components of a particular intervention. Abstract concepts such as hope or empathy, which are common factors of quality

© The Author(s), 2022. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

therapeutic consultations (Singla et al., 2017; Pedersen et al., 2020; Kohrt et al., 2020a) featured in task-sharing manuals, had to be turned into tangible-specific behaviours. These represent standards with the goal of accommodating the multitude of professional, educational and cultural backgrounds found in LRHS. This frame allows supervisors with different theoretical backgrounds to set consistent learning objectives, and in doing so enables supervisees to build on previously developed skills. Additionally, the use of structured case formulation discussions helps to build understanding of vulnerability and resilience (Sørbye et al., 2019) to produce more patient-centred care. To our knowledge, no tool for monitoring case formulation within supervision is available for non-specialists.

We hope that MSF's recent effort is a start. Yet, while we realize that one ideal tool for every context may not be feasible, a broader availability of tools may enable supervisors to better address the specificities of their contexts while maintaining a focus on evidence-based treatment and incorporating standardized manuals (e.g. mhGAP). This should also reduce the need for duplication between organizations. We eagerly await WHO's EQUIP package for psychological interventions (Kohrt et al., 2020b) and invite critical feedback on the MSF toolkit, which can be found here: <https://scienceportal.msf.org/assets/7610>. We hope that sharing our experience can help refine current and future supervision approaches, and we invite other global MH actors to design and share their tools to improve quality of care for some of the world's most vulnerable patients.

Acknowledgements. We would like to thank everyone in the various MSF projects who contributed to the development of this tool: Da'ed Almezil, Emmanuel Ballah, Nadia Duarte Marini, Mahmoud Hebib, Jean-Clement Ishimwe, Moathe Khaleel Ismail, Anna Krickl, Virginia Lee, Devash Naidoo, Hassan Nasser, Simone Silberberg, Marieke van Nuenen, Saima Zai. Thank you also to Mariana Duarte for supervision guidance, and to Anne McPherson for deconstruction.

Financial support. The writing of this manuscript was not supported financially. MSF medical editor Janet Ousley reviewed both manuscripts and provided publishing support. Article processing charges will be covered by MSF. The development of a competency-based supervision approach was supported through time-limited project funding by the Medical Department of MSF France in 2019.

Conflict of interest. Author B.B. was employed by MSF France until June 2021, and now works for the Department of Health in Nuremberg, Germany. Authors G.K., M.P. and J.O. are currently employed by MSF France and MSF New York.

References

Brockhouse R, Msetfi RM, Cohen K and Joseph S (2011) Vicarious exposure to trauma and growth in therapists: the moderating effects of sense of

coherence, organizational support, and empathy. *Journal of Traumatic Stress* 24, 735–742.

Carreño C, Keane G, Echeverri C, Goodfriend M, Tijerino AM and Fagard S (2019) Requirements for quality task-shifting in care of people with psychiatric disorder: MSF's experience with mhGAP intervention guide 2.0, Lisbon.

Faregh N, Lencucha R, Ventevogel P, Dubale BW and Kirmayer LJ (2019) Considering culture, context and community in mhGAP implementation and training: challenges and recommendations from the field. *International Journal of Mental Health Systems* 13, 58.

Heim E and Kohrt BA (2019) Cultural adaptation of scalable psychological interventions: a new conceptual framework. *Clinical Psychology in Europe* 1, 22.

Holt H, Beutler LE, Kimpara S, Macias S, Haug NA, Shiloff N, Goldblum P, Temkin RS and Stein M (2015) 'Evidence-based supervision: tracking outcome and teaching principles of change in clinical supervision to bring science to integrative practice. *Psychotherapy* 52, 185–189.

Kemp CG, Petersen I, Bhana A and Rao D (2019) Supervision of task-shared mental health care in low-resource settings: a commentary on programmatic experience. *Global Health: Science and Practice* 7, 150–159.

Kohrt BA, Ottman K, Panter-Brick C, Konner M and Patel V (2020a) Why we heal: the evolution of psychological healing and implications for global mental health. *Clinical Psychology Review* 82, Online Preprint, 101920.

Kohrt BA, Schafer A, Willhoite A, Van't Hof E, Pedersen GA, Watts S, Ottman K, Carswell K and van Ommeren M (2020b) Ensuring quality in psychological support (WHO EQUIP): developing a competent global workforce. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)* 19, 115–116.

Murray LK and Jordans MJD (2016) Rethinking the service delivery system of psychological interventions in low and middle income countries. *BMC Psychiatry* 16, 234.

Patel V, Saxena S, Lund C, Thornicroft G, Baingana F, Bolton P, Chisholm D, Collins PY, Cooper JL, Eaton J, Herrman H, Herzallah MM, Huang Y, Jordans MJD, Kleinman A, Medina-Mora ME, Morgan E, Niaz U, Omigbodun O, Prince M, Rahman A, Saraceno B, Sarkar BK, de Silva M, Singh I, Stein DJ, Sunkel C and Unützer J (2018) The Lancet Commission on global mental health and sustainable development. *The Lancet* 392, 1553–1598.

Pedersen GA, Lakshmin P, Schafer A, Watts S, Carswell K, Willhoite A, Ottman K, van't Hof E and Kohrt BA (2020) Common factors in psychological treatments delivered by non-specialists in low- and middle-income countries: manual review of competencies. *Journal of Behavioral and Cognitive Therapy* 30, 165–186.

Rønnestad MH, Orlinsky DE, Schröder TA, Skovholt TM and Willutzki U (2019) The professional development of counsellors and psychotherapists: implications of empirical studies for supervision, training and practice. *Counselling and Psychotherapy Research* 19, 214–230.

Singla DR, Kohrt BA, Murray LK, Anand A, Chorpita BF and Patel V (2017) Psychological treatments for the world: lessons from low- and middle-income countries. *Annual Review of Clinical Psychology* 13, 149–181.

Sørbye Ø, Dahl H-SJ, Eells TD, Amlo S, Hersoug AG, Haukvik UK, Hartberg CB, Høglend PA and Ulberg R (2019) Psychodynamic case formulations without technical language: a reliability study. *BMC Psychology* 7, 67.