

## NOTES FOR AUTHORS

Papers submitted for publication in the *Bulletin of Entomological Research* should be sent, in duplicate, to the Director, Commonwealth Institute of Entomology, 56 Queen's Gate, London SW7 5JR. They must deal with original research concerning insects, mites or ticks of economic importance in the agricultural, medical or veterinary fields; they must be of more than purely local interest, and taxonomic papers will not be accepted unless the organisms dealt with are of definite importance in the field of applied entomology. Review articles may also be submitted, but will normally be by invitation. Papers must not have been published or accepted for publication elsewhere. They will normally be published in order of their receipt in definitive form.

For general guidance on presentation and style, including the preparation of illustrations, contributors are urged to consult *Writing Scientific Papers in English. An ELSE-Ciba Foundation Guide for Authors* (by Maeve O'Connor & E. Peter Woodford), 116 pp., Amsterdam, Associated Scientific Publishers, 1975, ISBN 90-219-4035-3.

### Text

Papers should be typewritten (double spaced, with ample margins and on one side of the paper only) and a current number of the *Bulletin* should be consulted for details of style and lay-out. Titles should be as brief as is consistent with clarity. Abbreviations should conform to *British Standard 1991, Part 1, 1967*, and are alike in singular and plural; stops will be used only where ambiguity might occur, e.g. 'in.' not 'in' for inch, but 'in<sup>2</sup>' for square inch. Metric units are preferred; if non-metric units are used, metric equivalents should be given at least once in each article.

### Tables

Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must on no account be of a size or form that will not conveniently fit into the page of the *Bulletin*.

### Illustrations

Line drawings and graphs must be in jet-black indian ink, preferably on bristol board or tracing cloth. Graphs may be drawn on paper ruled in pale blue or light grey; paper having green, red, yellow or brown lines should not be used. Any stippling or hatching, as well as any numerals or lettering, on the text-figures must be completed, by the author, in a form suitable for reproduction; this work is *not* undertaken by the printer. Text-figures, when reproduced, cannot exceed 12.7 cm in width; if, therefore, the original drawings are larger, the lettering on them must be in proportion, so that it is still legible when reduced; special attention should be paid to the relative scales of lettering and line thicknesses in relation to the figure as a whole. Photographs for reproduction as figures are admissible when they are a real contribution to the text. Captions to figures should be typed on a separate sheet.

### Abstract

Each paper must contain a carefully prepared and accurate informative abstract that shall be complete in itself and intelligible without reference to the text or figures. It should not exceed 250 words, and should be included at the beginning immediately after the title, author and author's affiliation.

### References

References must be based on the name and year system and conform to the present usages of the *Bulletin*. Titles of journals should be abbreviated as in the 4th edition of *The World List of Scientific Periodicals*. References to unpublished information should normally be included only in the text—(A. B. Jones, pers. comm.) or (A. B. Jones. Control of sandflies by insecticides. Unpublished report. Ross Institute, London, 1950).

### Offprints

Free copies of offprints are allowed as follows: One author—20 copies; two or more authors—30 copies. Further copies may be obtained on payment, and the number required should be specified when the paper is submitted.

## BULLETIN OF ENTOMOLOGICAL RESEARCH

(Word List Abbreviation: *Bull. ent. Res.*)

ISSN 0007-4853

Review article	PAGE
TYNDALE-BISCOE, M. Age-grading methods in adult insects: a review .....	341
 <b>Original articles</b>	
PADGHAM, D. E., COOK, A. G & HUTCHISON, D. Rubidium marking of the rice pests <i>Nilaparvata lugens</i> (Stål) and <i>Sogatella furcifera</i> (Horvath) (Hemiptera: Delphacidae) for field dispersal studies .....	379
DEWAR, A. M. & CARTER, N. Decision trees to assess the risk of cereal aphid (Hemiptera: Aphididae) outbreaks in summer in England .....	387
HAYAT, M. <i>Encarsia bennetti</i> sp. n. (Hymenoptera: Aphelinidae) from India, associated with the citrus black fly, <i>Aleurocanthus woglumi</i> Ashby (Homoptera: Aleyrodidae) .....	399
ROGERS, D. J., RANDOLPH, S. E. & KUZOE, F. A. S. Local variation in the population dynamics of <i>Glossina palpalis palpalis</i> (Robineau-Desvoidy) (Diptera: Glossinidae). I. Natural population regulation .....	403
RANDOLPH, S. E., ROGERS, D. J. & KUZOE, F. A. S. Local variation in the population dynamics of <i>Glossina palpalis palpalis</i> (Robineau-Desvoidy) (Diptera: Glossinidae). II. The effect of insecticidal spray programmes .....	425
ROOM, P. M., PERRY, C. H. & BAILEY, P. T. A population study of the coconut pest <i>Segestidea uniformis</i> (Willemse) (Orthoptera: Tettigoniidae) on an equatorial island .....	439
TYNDALE-BISCOE, M. Adaptive significance of brood care of <i>Copris diversus</i> Waterhouse (Coleoptera: Scarabaeidae) .....	453
MATTHIESSEN, J. N., HAYLES, L. & PALMER, M. J. An assessment of some methods for the bioassay of changes in cattle dung as insect food, using the bush fly, <i>Musca vetustissima</i> Walker (Diptera: Muscidae) .....	463
WILSON, M. R. <i>Numicia gauhati</i> sp. n. (Homoptera: Fulgoroidea: Tropiduchidae) from rice in India, with taxonomic notes on other Indian <i>Numicia</i> species ..	469
DOLLING, W. R. Pentatomid bugs (Hemiptera) that transmit a flagellate disease of cultivated palms in South America .....	473
LENZ, M., BARRETT, R. A. & WILLIAMS, E. R. Implications for comparability of laboratory experiments revealed in studies on the effects of population density on vigour in <i>Coptotermes lacteus</i> (Froggatt) and <i>Nasutitermes exitiosus</i> (Hill) (Isoptera: Rhinotermitidae & Termitidae) .....	477
NASR, EL-SAYED A., TUCKER, M. R. & CAMPION, D. G. Distribution of moths of the Egyptian cotton leafworm, <i>Spodoptera littoralis</i> (Boisduval) (Lepidoptera: Noctuidae), in the Nile Delta interpreted from catches in a pheromone trap network in relation to meteorological factors .....	487
WATSON, J. A. L., DAI, Z. R. & BARRETT, R. A. Comparative studies on the resistance of materials to species of <i>Coptotermes</i> (Isoptera: Rhinotermitidae). I. The susceptibilities of various plastics to <i>C. acinaciformis</i> (Froggatt) from Australia and <i>C. formosanus</i> Shiraki from China .....	495
ROOM, P. M., FORNO, I. W. & TAYLOR, M. F. J. Establishment in Australia of two insects for biological control of the floating weed <i>Salvinia molesta</i> .....	505
KIDD, N. A. C. & CLEAVER, A. M. The relationship between pre-flight reproduction and migratory urge in alatae of <i>Aphis fabae</i> Scopoli (Hemiptera: Aphididae) ..	517
NOYES, J. S. A new genus and species of encyrtid (Hymenoptera: Chalcidoidea) parasitic on the cassava mealybug, <i>Phenacoccus manihoti</i> Matile-Ferrero (Hemiptera: Pseudococcidae) .....	529
WILLIAMS, D. J. Two new species of <i>Eurycoccus</i> (Hemiptera: Pseudococcidae) on <i>Tamarix</i> from Pakistan .....	535
SMITH, E. S. C. Studies on <i>Amblypelta theobromae</i> Brown (Heteroptera: Coreidae) in Papua New Guinea. I. Descriptions of the immature and adult stages. ....	541

© Commonwealth Agricultural Bureaux, 1984

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.