




Dietary fluoride intake over the course of pregnancy in Mexican women

Gina A Castiblanco-Rubio^{1,*} , Teresa V Muñoz-Rocha², Alejandra Cantoral², Martha M Téllez-Rojo², Adrienne S Ettinger^{3,4}, Adriana Mercado-García², Karen E Peterson^{3,5}, Howard Hu⁶ and Esperanza A Martínez-Mier¹

¹Department of Cariology, Operative Dentistry and Dental Public Health, Indiana University, School of Dentistry, 415 Lansing St, Indianapolis, IN 46202, USA: ²Center for Nutrition and Health Research, Instituto Nacional de Salud Pública de México (INSP), Cuernavaca, México: ³Department of Nutritional Sciences, University of Michigan, School of Public Health, Ann Arbor, MI, USA: ⁴Rutgers Biomedical and Health Sciences, Rutgers University, New Brunswick, NJ, USA: ⁵Department of Environmental Health Sciences, University of Michigan, School of Public Health, Ann Arbor, MI, USA: ⁶University of Washington, School of Public Health, Seattle, WA, USA

Submitted 4 August 2020: Final revision received 2 February 2021: Accepted 15 February 2021: First published online 19 February 2021

Abstract

Objective: To estimate dietary fluoride intake (F) over the course of pregnancy and the overall adjusted difference in dietary F intake by pregnancy stages and levels of compliance with dietary recommendations.

Design: Secondary data analysis from a longitudinal pregnancy cohort study in a population exposed to fluoridated salt. Women were followed during the early, middle and late stages of their pregnancy (n 568). The dietary intake of recommended prenatal nutrients according to Mexican dietary guidelines and F intake (mg/d) was estimated with a validated FFQ. Data were summarised with descriptive statistics. Levels of F intake were compared with the USA's Institute of Medicine adequate intake (AI) of 3 mg/d for pregnancy. Adjusted differences in F intake by pregnancy stages and levels of compliance with recommendations were estimated using random effects models.

Setting: Mexico City.

Participants: Women participating in the Early Life Exposures in Mexico to ENvironmental Toxicants (ELEMENT) project, from 2001 to 2003.

Results: Median dietary F intake throughout pregnancy ranged from 0.64 (interquartile range (IQR) 0.38) in the early to 0.70 (IQR 0.42) in the middle, and 0.72 (IQR 0.44) mg/d in the late stage (0.01 mg F/kg per d). Corresponding adjusted intakes of F were 0.72 (95% CI 0.70, 0.74), 0.76 (95% CI 0.74, 0.77) and 0.80 (95% CI 0.78, 0.82) mg/d. Women who were moderately and highly compliant with Mexican dietary recommendations ingested, on average, 0.04 and 0.14 mg F/d more than non-compliant women ($P < 0.005$).

Conclusions: Dietary F intake was below current AI, was greater with the progression of pregnancy and in women who were moderately and highly compliant with dietary recommendations.

Keywords
Fluoride
Dietary exposure
Prenatal exposure
Pregnancy

Fluoride (F) is present in small amounts in the soil, water, plants and animals; therefore, it is naturally present as a trace element in the diet⁽¹⁾. An extensive body of evidence has proven a clear reduction in dental caries prevalence in communities exposed to F, and it has been added to the diet via water or salt for human consumption as part of community fluoridation programmes to prevent dental caries⁽²⁾. Like other nutrients and trace elements in the diet,

F has both beneficial and detrimental effects: with low exposures, it prevents and controls dental caries, while higher exposures can lead to hard-tissue changes such as dental fluorosis⁽³⁾. Given that the risk for developing dental fluorosis is present only during critical periods of tooth development, dietary F intake has been extensively and traditionally monitored in children⁽⁴⁾, but rarely in other age groups. There is, however, emerging evidence on

*Corresponding author: Email ginacast@iu.edu

© The Author(s), 2021. Published by Cambridge University Press on behalf of The Nutrition Society



potential adverse effects of prenatal F exposure⁽⁵⁾. Associations between F concentration in urine during pregnancy and poor neurodevelopmental outcomes in the offspring have been reported, not only in populations with endemic fluorosis and exposed to high levels of F⁽⁶⁾ but also in populations exposed to low levels, such as the ones considered optimal for community water and salt fluoridation programmes^(7–9). This newly emerging evidence on the potential side effects of prenatal F exposure suggests monitoring of F intake in other susceptible groups, such as pregnant women, may be warranted.

In 1997 and based on data collected in children and non-pregnant adults, the USA's Institute of Medicine recommended an adequate intake (AI) for F of 3 mg/d (0.05 mg/kg per d). Using a pre-pregnancy body weight for women >19 years of ~61 kg as a reference, the recommendation for both pregnant and non-pregnant women was also set as 3 mg F/d⁽¹⁰⁾. Given the existing knowledge gap on F intake in the context of the dietary and physiological changes of pregnancy, there is a need for observational studies of F intake in populations of pregnant women. For instance, pregnant women are encouraged to increase the dietary intake of foods and supplements containing nutrients that are beneficial for maternal and fetal health, such as Ca, Fe and folate⁽¹⁰⁾. In contrast, the dietary intake of F is not particularly encouraged or discouraged during pregnancy and has no reported benefits for fetal health. In a previous investigation on the concentration of F in foods and beverages available in Mexico City, we found small amounts of F in Mexican dietary staples (cereals, legumes and animal products)⁽¹¹⁾. These foods are rich sources of folate, Ca, Fe and protein⁽¹²⁾ and their frequent consumption may increase the total daily dietary intake of F. We hypothesised that Mexican pregnant women, attempting to meet dietary recommendations, increase their dietary intake of F. Understanding how F intake changes over the course of pregnancy and, whether meeting the requirements of key beneficial nutrients also increases the dietary intake of F, can serve the purpose of informing future dietary recommendations for pregnancy.

The objectives of the current study were to estimate dietary F intake over the course of pregnancy and the overall adjusted difference of dietary F intake by pregnancy stage and levels of compliance with Mexican dietary recommendations.

Methods

Study sample and setting

The study population was pregnant women participating in the Early Life Exposures in Mexico to ENvironmental Toxicants (ELEMENT) project, which has been described previously⁽¹³⁾. ELEMENT comprises three mother–child pregnancy and birth cohorts, initiated in the 1990s to study early life exposures and health outcomes in Mexico City. In the current analysis, we included pregnant women from Cohort 3, who were recruited

between 2001 and 2003 (*n* 670). Pregnant women attending three clinics of the Mexican Institute of Social Security (IMSS) were invited to participate, but only those with gestational age <14 weeks, a healthy singleton pregnancy, no history of systemic diseases (hypertension, diabetes) and intention to stay in Mexico City who agreed to participate through informed consent were included and followed thereafter. Socio-demographic questionnaires included several questions on variables such as age, educational attainment, parity, marital status and self-reported weight before pregnancy. Details and demographics of the entire project can be found in the ELEMENT project's profile publication⁽¹³⁾. The project has a dedicated research facility next to Mexico City's ABC Medical Center and participating women were invited to attend the facility at the early, middle and late stages of their pregnancy⁽¹³⁾. Cohort 3 was originally designed as a double-blind randomised clinical trial (RCT) to examine the effects of Ca supplementation on blood lead levels during pregnancy and up to 1 year post-partum⁽¹⁴⁾. Women were randomised to receive either the Ca supplement (1200 mg Ca/d) or placebo. During each pregnancy visit, each woman was interviewed by a social worker, who performed anthropometry (weight and height) using calibrated instruments, applied a general demographic questionnaire and a FFQ. Only women from Cohort 3 with FFQ and all variables of interest available in the database were included for the analyses (*n* 568). All participants who were included had data available for at least two pregnancy stages, and 511 had data for all three.

Measurement of fluoride in foods, water and other beverages

Mexico City has naturally occurring water fluoride levels <0.7 ppm⁽¹⁵⁾ and does not have community water fluoridation as used in other countries. Instead, since 1981 a salt fluoridation programme was implemented as a method to deliver F for caries prevention (250 ppm of F/kg) in regions with water fluoride levels <0.7 ppm^(16,17). Therefore, the main sources of dietary F intake in Mexico City are foods with intrinsic F content and those containing fluoridated salt added either during the cooking process and/or at the table before consumption. A database was developed specifically for the ELEMENT project by analysing the F content of typical foods and beverages in the Mexican diet⁽¹¹⁾. Details on the methodology for the collection of food and beverages and the analysis of F can be found in the publication by Cantoral *et al.*⁽¹¹⁾. Briefly, fruits and vegetables were bought from three different major markets in Mexico City. Meat products, processed foods, juices, beverages and industrialised foods were bought from four large supermarket chains. Natural juices were purchased from street vendors, flavoured waters (a traditional Mexican beverage prepared with water, fruit and sugar) were bought from ice cream parlors and dairy was purchased in creameries. To measure the concentration of F in water drunk by participating women, a trained research assistant visited the household of study



participants of the ELEMENT project (n 552) and collected water samples (~ 5 ml). To standardise the concentration of F in the water and salt in foods that are consumed cooked (meats, rice, pasta, legumes), these were boiled using water containing negligible amounts of F (< 0.01 mg/l) and fluoridated salt (250 ppm of F/kg), following standardised recipes from the National Health and Nutrition Survey. Traditional foods (e.g., tamales and maize-based foods) were purchased cooked from street vendors. Fluoride analyses were conducted at the Fluoride Research Laboratory at the Oral Health Research Institute, Indiana University School of Dentistry using a modification of the hexamethyldisiloxane method⁽¹⁸⁾ as modified by Martinez-Mier *et al.*⁽¹⁹⁾. Participants reported the intake of mainly bottled and tap water, which had mean F levels \pm SD of 0.16 ± 0.13 ppm and 0.14 ± 0.09 ppm, respectively. These two average values were the ones included in the database for FFQ-derived estimations of F intake from water.

Estimation of dietary intake of fluoride, macro- and micronutrients

Dietary F intake (mg F/d) was assessed through a semiquantitative questionnaire consisting of 104 items, adapted from the Willett semi-quantitative FFQ⁽²⁰⁾ to include foods and beverages commonly consumed in the 1983 Dietary Survey of the Mexican National Institute of Nutrition⁽²¹⁾ and validated to estimate dietary intake over the previous month in Mexican women of child-bearing age (15–44 years)⁽²²⁾. The FFQ was applied by a trained social worker at each study visit once per trimester, using visual and measuring aids (spoons, cups) for the identification of foods and portion sizes. Estimates of dietary intake of macro- and micronutrients (in μ g or mg/d) were calculated through software developed at the National Institute of Public Health (INSP) using methods for the analysis of dietary data from the Mexican National Health and Nutrition Survey⁽²³⁾. To generate estimates of daily dietary intake, the software utilises the following data: (1) the average content of nutrients and trace elements (including F) of each food/beverage item reported the INSP-compiled nutrient composition database⁽²⁴⁾ and the ELEMENT F database, and (2) the reported frequencies and portion sizes of foods and beverages from the FFQ. Further details on the development, validation and calculations derived from the FFQ are available in the publication by Hernandez-Avila *et al.*⁽²²⁾. Although the FFQ did not include quantitative estimations of table salt intake, it did include a dichotomous question on whether table salt was added to foods right before eating them (yes/no), which we used for this secondary data analysis. Given that this cohort was originally designed as a Ca supplementation RCT and women may have chosen to ingest other dietary supplements as well, a variable with group assignment to the RCT (placebo/supplemented) and another one specifying intake of other dietary supplements (yes/no) was available to control for potential confounding.

Assessment of compliance with dietary recommendations

Dietary F-intake in Mexican women is neither contraindicated nor encouraged in Mexican pregnancy dietary recommendations. The Mexican Official Norms regulating health services for pregnant women, which were current for the cohort, highly encouraged an increase in the dietary intake of key nutrients⁽¹²⁾ in order to meet the AI or estimated average requirement (EAR) as recommended by the dietary reference intakes by the USA Institute of Medicine⁽¹⁰⁾. Recommendations on dietary intake during pregnancy were made according to pre-pregnancy BMI and pregnancy stage⁽²⁵⁾. The four key nutrients encouraged in the dietary recommendations were Ca (AI: 1000 mg/d), Fe (EAR: 22 mg/d), folate (EAR: 520 μ g/d) and protein (EAR: 0.88 g/kg per d). Intake was recommended from various dietary sources for all nutrients⁽²⁶⁾; in the case of Ca, folate and Fe, the recommendation included intake from both diet and supplements. The dietary intake of each nutrient was estimated only from the intake reported in the FFQ. Since one of the objectives of the current study was focused on the relationship between compliance with nutrients from dietary sources and dietary F intake, supplement sources were excluded from the calculation.

Women were classified as compliant with an individual nutrient if their estimated daily dietary intake was equal to or above the AI or EAR for that particular nutrient during pregnancy. Overall compliance at each stage of pregnancy was categorised according to individual-nutrient compliance, as follows: non-compliance if non-compliant with all the nutrients (0/4); moderate compliance if compliant with one or two nutrients (1/4 or 2/4) and high compliance if compliant with three or all the four nutrients (3/4 or 4/4).

Covariates

The selection of covariates was based on our knowledge of factors that may influence dietary F intake and bivariate analyses. These included pregnancy stage, compliance with dietary recommendations, group allocation in the Ca supplementation RCT, intake of other supplements, addition of salt at the table, total daily energy intake, pre-pregnancy BMI and educational attainment.

Data analyses

Study participants were stratified by pregnancy stage. Differences in key dietary variables across pregnancy stages were assessed with Friedman test (for continuous, non-normally distributed repeated measures variables) or χ^2 tests (for categorical variables). Differences in dietary F intake between the RCT allocation groups and according to table salt use were tested with the non-parametric Mann–Whitney test. To estimate the overall adjusted difference in dietary F intake by pregnancy stage (early, middle, late) and levels of compliance with Mexican dietary recommendations (none, moderate, high), we fitted multivariate



regression models for panel data using the `xreg` command in STATA. Panel data models were chosen because they examine both individual- and time-specific effects to deal with unobserved heterogeneity or individual effects. Regression diagnostics revealed significant differences in the individual- and time-specific variance components (panel effect; Breusch–Pagan LM test $P < 0.05$) and no correlation between individual effects and the regressors (Hausman test, $P > 0.05$); therefore, one-way random effects GLS regressions provided the best fit. After the model's estimates were obtained, adjusted predictions for dietary F intake were calculated for each pregnancy stage using the `margins` command. The association between dietary F intake and individual intake of nutrients (Ca, Fe, folate, protein) was assessed following the same approach. All analyses were conducted with STATA v16.0 (StataCorp.).

Results

Characteristics of the sample and dietary fluoride intake by pregnancy stage

Table 1 summarises and compares the characteristics of women included in the analytical sample ($n = 568$) and those who were excluded ($n = 102$). Overall, the median age of the study participants at the time of recruitment was 26.4 years and the median gestational age for women who completed FFQ at the early, middle or late stages of pregnancy was 13.6, 25.4 and 34.3 weeks, respectively. Median weight ranged between 60.5 and 69.4 kg during the three pregnancy stages and most women (57.7%) had a pre-pregnancy BMI categorised as normal according to the guidelines that were current at the time of data collection⁽¹⁴⁾. Most women (64.3%) had been pregnant more than once; and their highest educational level was secondary school (67.8%). Finally, in the analytical sample, 289 women were allocated to the treatment group of the Ca supplementation RCT, whereas 279 were allocated to the placebo group. The characteristics of women who were excluded were not significantly different from those who were included.

Detailed descriptive statistics of dietary F intake by pregnancy stages and for all available observations are provided in Table 2. This sample of pregnant women living in Mexico City had a median dietary F intake of 0.69 mg/d, ranging between a minimum of 0.11 and a maximum of 3.73 mg/d (for a total of 1649 observations at all pregnancy stages).

Bivariate statistics for key dietary variables by pregnancy stages

Table 3 summarises bivariate statistics of key dietary variables by pregnancy stages. We found variability in both dietary F intake and total energy intake over the course of pregnancy. Women had a median dietary F intake that ranged from 0.64

(IQR 0.38) in the early stage to 0.72 (0.44) mg F/d in the late stage, and the tendency to increase was statistically significant. Considering that the median weight across pregnancy was 65 kg, this range of median F intake throughout pregnancy corresponds to 0.01 mg/kg per d. Estimates of dietary F intake between those who reported adding salt at the table to their meals and those who did not were not significantly different (0.67 mg/d interquartile range 0.47 *v.* 0.70 mg/d, interquartile range 0.40, respectively) (data not shown in Table 3). In contrast to dietary F, median total energy intake significantly decreased during the middle stage of pregnancy. There were also variations in overall compliance and individual-nutrient compliance across pregnancy stages. Compliance with both Ca and Fe was associated with pregnancy stage. For Ca, compliance rose from 51.2% in the early, 58.6% in the middle and 62.0% in the late stages of pregnancy ($P < 0.001$), whereas for Fe it went from 6.0% in the early, 7.1% in the middle and, 5.5% in the late stage – although for the latter, no statistically significant association was found. In contrast, compliance with protein was associated ($P < 0.001$) with pregnancy stage (68.7, 62.6 and 56.8% for the early, middle and late stages of pregnancy, respectively). Compliance with folate intake from dietary sources experienced a slight increase towards the middle stage, followed by a decrease towards the end of gestation (5.8, 6.3 and 3.8% for the early, middle and late stages of pregnancy, respectively), although this change was not associated with pregnancy stage ($P = 0.151$). Over the course of pregnancy, women were mostly moderately compliant with dietary recommendations ($P = 0.412$). Only a slight increase in the proportion of women who were non-compliant towards the end of pregnancy was observed (not statistically significant). Towards the end of pregnancy, the proportion of women reporting adding table salt to meals was lower (30.3% in early *v.* 22.6% in late-pregnancy; $P = 0.012$), whereas the proportion of women reporting the use of dietary supplements was higher (23.4% in early *v.* 37.9% in late pregnancy; $P < 0.001$) (Table 3). Median F intake from dietary sources for women allocated to the Ca supplementation group in the RCT was significantly higher than that of women allocated to the placebo group (overall median 0.71 *v.* 0.67, $P = 0.03$; data not shown in Table 3); therefore, we also included RCT allocation group as a covariate in the models.

Adjusted dietary fluoride intake by pregnancy stages and compliance with dietary recommendations

After adjustment for covariates, the association between dietary F intake and pregnancy stages was significant ($P < 0.001$, Table 4). Compared with the early stage, women in the middle and late stages of their pregnancy ingested on average, 0.04 and 0.08 more mg F/d, respectively. The adjusted predictions of dietary F intake from foods and beverages during pregnancy therefore increase from 0.72 (95% CI 0.70, 0.74) in the early stage, 0.76 (95%

Table 1 Characteristics of women with complete data for all variables of interest that were included in the analytical sample (included) and women who were excluded because they had incomplete data‡

	All*	Included			Excluded			P
		n	Median	IQR	n	Median	IQR	
Age at study recruitment (years)	670	568	26.4	8.0	102	27.1	8.1	0.840
Gestational age (weeks)								
Early pregnancy	635	568	13.6	2.3	67	13.4	2.0	0.775
Middle pregnancy	557	554	25.4	2.3	3	24.7	2.3	0.503
Late pregnancy	528	527	34.3	2.5	1	33.7	0.0	0.726
Weight (kg)								
Early pregnancy	668	568	60.5	13.5	100	61.75	15.5	0.798
Middle pregnancy	555	554	65.5	13.1	1	77	0.0	0.462
Late pregnancy	528	527	69.4	13.5	1	58.2	0.0	0.163

	All*	Included		Excluded		P		
		n	Freq	%	n		Freq	%
Pre-pregnancy BMI								
Underweight	670	568	35	6.2	102	15	14.7	0.082
Normal			328	57.7		50	49.0	
Overweight			102	18.0		20	19.7	
Obese			103	18.1		17	16.6	
Educational attainment								
None	670	568	1	0.2	102	0	0.0	0.707
Elementary			129	22.7		21	20.6	
Middle school			385	67.8		68	66.7	
High school			53	9.3		13	12.7	
Marital status								
Married	670	568	387	68.1	102	73	71.6	0.451
Common-law marriage			111	19.5		22	21.6	
Single			69	12.2		7	6.8	
Divorced			1	0.2		0	0.0	
First pregnancy, yes	670	568	203	35.7	102	34	32.1	0.640
Assigned to Ca supplement, yes†	670	568	289	50.8	102	45	44.1	0.208

*Total number of women with data available for each variable.

†Number of women assigned to the Ca supplementation group in the randomised clinical trial.

‡Comparisons between included and excluded women were performed with Mann-Whitney tests for continuous variables and χ^2 tests for categorical variables.

Table 2 Dietary fluoride intake (mg/d) by pregnancy stages. The 'all stages' category represents data for all observations available during pregnancy

Pregnancy stage	n	Mean	SD	Min	10 %	25 %	50 %	75 %	90 %	Max
Early	568	0.72	0.35	0.11	0.35	0.49	0.64	0.87	1.18	2.05
Middle	554	0.76	0.36	0.14	0.38	0.52	0.70	0.94	1.20	3.73
Late	527	0.79	0.39	0.11	0.40	0.52	0.72	0.96	1.24	3.08
All stages	1649	0.76	0.37	0.11	0.38	0.51	0.69	0.93	1.21	3.73

CI 0.74, 0.77) in the middle stage, to 0.80 (95 % CI 0.78, 0.82) in the late stage (Fig. 1). Women who reported intake of dietary supplements other than the Ca supplement provided for the RCT supplementation group ingested, on average, 0.03 less mg F/d, compared with women who did not take other supplements ($P=0.027$). Furthermore, compared with non-compliant women, those who were moderately and highly compliant with dietary recommendations ingested on average, 0.04 and 0.14 more mg F/d, respectively (Table 4). The covariates included in the model were total energy intake, allocation group in the Ca supplementation RCT, intake of other supplements and pre-pregnancy BMI.

In order to understand which nutrients were associated with changes in dietary F intake, we also assessed the association between compliance with individual nutrients and dietary F intake. Compared with women who did not meet Ca and Fe recommendations, those who were compliant with Ca and Fe recommendations ingested on average, 0.12 and 0.18 more mg F/d, respectively ($P<0.001$). In contrast, women who were compliant with folate recommendations ingested on average, 0.12 less mg F/d, compared with those who did not meet recommendations ($P=0.009$) (Table 5). Compliance with protein intake recommendations was not associated with dietary F intake.

Table 3 Bivariate statistics for key dietary variables by pregnancy stages

	Pregnancy stage						<i>P</i> *
	Early (<i>n</i> 568)		Middle (<i>n</i> 554)		Late (<i>n</i> 527)		
	Median	IQR	Median	IQR	Median	IQR	
Fluoride intake (unadjusted, mg/d)	0.64	0.38	0.70	0.42	0.72	0.44	<0.001
Total energy intake (kcal)	1811.5	802.5	1785.7	828.2	1802.2	706.0	<0.001
	%		%		%		<i>P</i> -value†
Compliance with individual nutrients (% above AI or EAR)							
Ca (AI: 1000 mg/d)	51.2		58.6		62.0		<0.001
Fe (EAR: 22 mg/d)	6.0		7.1		5.5		0.571
Folate (EAR: 520 µg/d)	5.8		6.3		3.8		0.151
Protein (EAR: 0.88 g/kg/d)	68.7		62.6		56.8		<0.001
Compliance‡							
None	25.4		27.1		30.4		0.412
Moderate	66.9		64.7		62.7		
High	7.8		8.1		6.8		
Adds salt at the table, yes	30.3		28.6		22.6		0.012
Use of other supplements, yes	23.4		32.0		38.0		<0.001

IQR, interquartile range.

*Friedman test (*n* 511).

† χ^2 test.

‡None: non-compliant with all of the nutrients (0/4); moderate: compliant with one or two nutrients (1/4 or 2/4); and high: compliant with three or all the four nutrients (3/4 or 4/4).

Table 4 Adjusted associations between daily fluoride (mg/d), pregnancy stage and compliance with dietary recommendations*

<i>n</i> 568	β^*	SE	95% CI		<i>P</i>
Pregnancy stage					
Early	Ref	Ref	Ref	Ref	Ref
Middle	0.04	0.01	0.01	0.06	<0.001
Late	0.08	0.01	0.06	0.10	<0.001
Compliance‡					
None	Ref	Ref	Ref	Ref	Ref
Moderate	0.04	0.02	0.01	0.08	0.004
High	0.14	0.03	0.08	0.20	<0.001
Intake of other supplements‡					
Yes	-0.03	0.01	-0.05	0.00	0.027
Adds salt at the table					
Yes	-0.01	0.01	-0.04	0.01	0.289

*Estimates from a one-way random effects GLS regression model adjusted for total energy intake, allocation group in the Ca supplementation RCT, intake of other supplements and pre-pregnancy BMI.

‡None: non-compliant with all of the nutrients (0/4); moderate: compliant with one or two nutrients (1/4 or 2/4) and high: compliant with three or all the four nutrients (3/4 or 4/4).

§Supplements other than the one provided for the Ca supplementation RCT.




Fig. 1 Estimated fluoride intake (95 % CI) by pregnancy stage*. *Estimates from a one-way random effects GLS regression model adjusted for total energy intake, allocation group in the Ca supplementation RCT, intake of other supplements and pre-pregnancy BMI

Discussion

Median dietary F intake in this sample of pregnant women living in Mexico City was 0.69 (min – max: 0.11–3.73) mg/d or 0.01 mg/kg per d. To date, only two other studies have reported dietary F intake during pregnancy in large samples of pregnant women. In Canada, F intake from beverages was assessed in 162 pregnant women living in communities with access to fluoridated water, with a reported intake of (mean ± SD) 0.93 ± 0.43 mg/d⁽⁹⁾. Another study, conducted in Spain in 575 pregnant women also living in a community with access to fluoridated water, reported a median F intake from beverages of 0.02 mg/kg per d (min – max: 0.005–0.043)⁽²⁷⁾. While the median

dietary F intake reported in our study is lower than in the other study populations, there are differences that limit the ability to make direct comparisons between these three studies. First, the study samples were drawn from populations exposed to different vehicles for community fluoridation. In Mexico City, community-wide fluoridation of salt is used, while in the Canadian and Spanish studies, women had access to fluoridated drinking water. Second, the studies from Canada and Spain only report F intake from beverages, while we report F intake from both foods and beverages, including bottled and tap water. Nonetheless, all three study populations found dietary F intakes in pregnancy below the USA-Institute of Medicine’s AI recommendation from all sources.

Table 5 Adjusted associations between daily fluoride intake and compliance with individual key nutrients*

<i>n</i> 568	β^*	SE	95 % CI		<i>P</i>
Ca					
Non-compliant	Ref	Ref	Ref	Ref	Ref
Compliant	0.12	0.01	0.09	0.14	<0.001
Fe					
Non-compliant	Ref	Ref	Ref	Ref	Ref
Compliant	0.18	0.27	0.13	0.24	<0.001
Folate					
Non-compliant	Ref	Ref	Ref	Ref	Ref
Compliant	-0.12	0.03	-0.18	-0.064	<0.001
Protein					
Non-compliant	Ref	Ref	Ref	Ref	Ref
Compliant	-0.02	0.02	-0.05	0.01	0.247

*Estimates from a one-way random effects GLS regression model adjusted for pregnancy stage, allocation group in the Ca supplementation RCT, intake of other supplements, total energy intake, addition of salt at the table (yes/no), pre-pregnancy BMI and educational attainment.

Our report has several strengths in the area of F exposure assessment, including repeated measures reported over the course of pregnancy, use of a validated instrument of dietary assessment and a F database specific to the population under study. Limitations of the current study include those inherent to FFQ-derived estimates and secondary data analyses. First, low F water (<0.01 mg/l) and fluoridated salt with standard F levels (250 ppm/kg)⁽¹⁶⁾ were used to prepare foods that are consumed cooked. There may be variation in the natural levels of fluoride in water and even among and within brands of salt⁽¹⁵⁾, with the potential to add uncertainty to the estimates. And second, we lack quantitative estimates of salt added after cooking. We believe, however, that as fluoride levels in Mexico City are low (<0.7 ppm)⁽¹⁵⁾ and levels of F in salt are regulated by the government^(16,17), the lack of a quantitative estimate of F in salt after cooking is the main source of uncertainty in the present study, which most likely underestimates the true dietary F intake. To control for potential differences that could be explained by the habit of adding salt at the table, we used a dichotomous question on the practice of adding table salt to meals after the cooking process (yes/no) that was available in the FFQ. This variable allowed us to make a rough calculation of what total F intake would look like in women who add salt to their meals. Using a quantitative report of added table salt by Mexican non-pregnant women aged 23–50 years in the State of Mexico (5.4 g of salt/d)⁽²⁸⁾ and assuming all salt gets ingested at a concentration of 250 ppm F, women in our study who reported adding table salt to their meals (median dietary F intake of 0.67 mg/d) would be ingesting about 1.31 mg F/d (0.02 mg F/kg per d) – approximately double the amount of those who reported not adding extra salt. This rough calculation, however, should be interpreted with caution, as quantitative estimations of salt intake after cooking were not made in the sample under study. We recommend for future studies in populations of pregnant women exposed to fluoridated salt, the inclusion of

quantitative measures of table salt intake. Furthermore, the median dietary intake found for this population (0.69 mg/d) constitutes the contribution of only intrinsic F in foods and F from salt added to foods during the cooking process. To provide a broader perspective of fluoride exposure, future research should also consider the contribution of other sources of F, such as occupational exposures and the unintentional intake of fluoridated oral hygiene products.

Dietary F intake in this sample of pregnant women increased with the progression of pregnancy, suggesting that dietary F intake does change during gestation – as would be expected given increased consumption by pregnant women to meet the nutrient demands of the growing fetus. We were interested in testing whether the increase observed in F intake throughout pregnancy remained after controlling for covariates that explain dietary F intake such as total energy intake, the practice of addition of salt at the table and compliance with dietary recommendations. We chose these variables because most nutrients, including F, have a positive linear relationship with total energy intake⁽²⁹⁾; pregnant women increase their total energy intake towards the end of gestation responding to an increased RMR⁽³⁰⁾, and there is tendency to change dietary behaviours with the progression of pregnancy⁽³¹⁾. In fact, controlling for total energy intake, women ingested more F in the middle and late stage of pregnancy compared with the early stage, plus 0.14 mg F/d (~0.2 mg/d total) if they were compliant with pregnancy dietary recommendations. The biological significance of an increase of ~0.2 mg/d of F intake during the third trimester of pregnancy (~7% of the current recommendation) is, however, still unknown and should be considered for future research. Studies on the association between maternal dietary F intake and health outcomes in the offspring are needed to investigate the biological significance of the current levels of intake. Only one study has reported that a 1-mg increase in maternal F intake from beverages was associated with a 3.7 decrease in intelligence scores among boys and girls⁽⁹⁾, which would be of public health significance for about 10% of the women in the present study (90th percentile, Table 1). Further research on the effect of lower prenatal F exposure levels on neurodevelopment – such as the median dietary intake found in the current study (0.69 mg/d), is needed to inform future dietary F intake recommendations for pregnant women.

We were interested in understanding whether Mexican women, attempting to meet dietary recommendations, increase their dietary intake of F. We found an association between women who were compliant with recommendations for both Ca and Fe and increased F intake levels (Table 5). This observation could be explained by the frequent consumption of Ca- and Fe-rich foods with low-to-moderate amounts of fluoride, which can lead to an overall increase in F intake. In Mexico, foods rich in both Ca and Fe with a moderate content of F include milk, maize-based products and legumes⁽¹¹⁾. We also found that women who reported to consume dietary supplements had lower dietary



F intakes. A plausible explanation for this negative association is that individuals who proactively take supplements tend to eat lower amounts of nutrient-dense foods⁽³²⁾. Therefore, it is possible that pregnant women in this sample who reported supplement intake relied on supplementation to meet their dietary goals instead of choosing more nutrient-dense foods. Socio-demographic factors have also been reported to influence dietary F intake⁽³³⁾; however, although we found no association, we cannot entirely tease out their effect on dietary intakes of the current study's population given that it included a relatively homogeneous group of women attending the clinics of the Social Security System in Mexico (IMSS) that serves a low-to-middle income population.

Within the limitations of the current study, we conclude that the levels of dietary F intake were below the current AI, were greater towards the end of gestation and in women who were moderately and highly compliant with Mexican dietary recommendations. Given the mounting evidence of potential adverse effects⁽⁷⁻⁹⁾ and multiple sources of exposure to F, additional assessment and monitoring of dietary intakes and exposures from other community sources, especially in vulnerable populations such as pregnant women and children, should be considered in future dietary recommendations for F intake.

Acknowledgements

Acknowledgements: The authors acknowledge the American British Cowdray (ABC) Hospital in Mexico City, Mexico, for providing research facilities for the ELEMENT project. We also acknowledge the active participation of women from the ELEMENT cohort 3 and the members of the research team involved in the ELEMENT project. **Financial support:** Indiana University's President's International Research Award (PIRA 23-140-39). NIH RO1ES021446 and NIEHS/EPA P01ES022844 RD8354360. PhD in Dental Sciences Program, Indiana University School of Dentistry. Instituto Nacional de Salud Pública de México (INSP). **Conflict of interest:** There are no conflicts of interest. **Authorship:** G.A.C. participated in the conception and design of the study, analysis of data, statistical modeling, interpretation of results and writing of the manuscript. T.V.M.-R. participated in the management of the ELEMENT database, analysis of data, statistical modeling interpretation of data and, writing of the manuscript. A.C., A.M. and M.M. T.-R. participated in the conception and design of the study, interpretation of results and writing of the manuscript. H. H. was a co-founder of the ELEMENT cohort and PI of RO1ES021446, and Karen Peterson is PI of NIEHS/EPA P01ES022844/RD8354360; both provided oversight of original dietary data collection and participated in interpretation of data and preparation of the manuscript. M.M.T.-R and A.M.-G. supervised and coordinated field data collection. A.M.-G.

and A.E. participated in the design and implementation of the original cohort study and participated in interpretation of data and preparation of the manuscript. **Ethics of human subject participation:** The current study was conducted according to the guidelines laid down in the Declaration of Helsinki and the ethics committee of the INSP, Indiana University, Harvard University and the University of Michigan approved all procedures involving research study participants. Written informed consent was obtained from all study participants.

References

- Smolin LA & Grosvenor MB (2010) The trace elements: fluoride. In *Nutrition: Science and Applications*, 2nd ed., pp. 515–518. New Jersey: John Wiley and Sons, Inc.
- Sampaio FC & Levy SM (2011) Systemic fluoride. In *Fluoride and the Oral Environment. Monographs Oral Science*, pp. 133–145 [MAR Buzalaf, editor]. Basel: Karger.
- DenBesten P & Li W (2011) Chronic fluoride toxicity: dental fluorosis. In *Fluoride and the Oral Environment. Monographs Oral Science*, pp. 81–96 [MAR Buzalaf, editor]. Basel: Karger.
- Buzalaf MAR & Levy SM (2011) Fluoride intake of children: considerations for dental caries and dental fluorosis. In *Fluoride and the Oral Environment. Monographs Oral Science*, pp. 1–19 [MAR Buzalaf, editor]. Basel: Karger.
- Grandjean P (2019) Developmental fluoride neurotoxicity: an updated review. *Environ Health* **18**, 110.
- Jiménez LV, Guzmán ODL, Flores MC *et al.* (2017) In utero exposure to fluoride and cognitive development delay in infants. *Neurotoxicology* **59**, 65–70.
- Bashash M, Thomas D, Hu H *et al.* (2017) Prenatal fluoride exposure and cognitive outcomes in children at 4 and 6–12 years of age in Mexico. *Environ Health Perspect*. Published online: 19 September 2017. doi: 10.1289/EHP655.
- Bashash M, Marchand M, Hu H *et al.* (2018) Prenatal fluoride exposure and attention deficit hyperactivity disorder (ADHD) symptoms in children at 6–12 years of age in Mexico City. *Environ Int*. Published online: 10 October 2018. doi: 10.1016/j.envint.2018.09.017.
- Green R, Lanphear B, Hornung R, *et al.* (2019) Association between maternal fluoride exposure during pregnancy and IQ scores in offspring in Canada. *JAMA Pediatr*. Published online: 19 August 2019. doi: 10.1001/jamapediatrics.2019.1729.
- Institute of Medicine (1997) *Fluoride. Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride*. Washington, DC: The National Academies Press.
- Cantoral A, Luna-Villa LC, Mantilla-Rodríguez AA *et al.* (2019) Fluoride content in foods and beverages from Mexico city markets and supermarkets. *Food Nutr Bull*. Published online: 25 July 2019. doi: 10.1177/0379572119858486.
- Roselló-Soberón M & Casanueva E (2005) Orientación alimentaria en el control prenatal: una herramienta educativa [Nutritional orientation in prenatal visits: an educational tool]. *Perinatol y Reprod Humana* **19**, 161–167.
- Peng W, Tamayo-Ortiz M, Tang L *et al.* (2019) Early life exposure in Mexico to environmental toxicants (ELEMENT) project. *BMJ Open*. Published online 26 August 2019. doi: 10.1136/bmjopen-2019-030427.
- Ettinger AS, Lamadrid-Figueroa H, Mercado-García A *et al.* (2014) Effect of calcium supplementation on bone resorption



- in pregnancy and the early postpartum: a randomized controlled trial in Mexican women. *Nutr J* **13**, 116. doi: 10.1186/1475-2891-13-116.
15. Martínez-Mier EA, Soto-Rojas AE, Buckley CM *et al.* (2005) Fluoride concentration of bottled water, tap water, and fluoridated salt from two communities in Mexico. *Int Dent J* **55**, 93–99.
 16. Secretaría de Salud & Estados Unidos Mexicanos (1981) Norma oficial Mexicana NOM-040-SSA-1981: reglamento de yodación y fluoruración de la sal [Official Mexican Norm NOM-040-SSA-1981: regulation of salt iodization and fluoridation]. *Diario Oficial de la Federación* **1**, 20–22.
 17. Secretaría de Salud & Estados Unidos Mexicanos (1996) Listado mediante el cual se dan a conocer las áreas por entidad federativa donde no debe comercializarse sal yodada fluorada, por tener el agua de consumo humano una concentración de flúor natural mayor de 0.7 mg/l [List of areas within the Federation where iodized and fluoridated salt should not be distributed, due to natural fluoride water levels being higher than 0.7 mg/l]. http://dof.gob.mx/nota_detalle.php?codigo=4879673&fecha=11/04/1996 (accessed January 2021).
 18. Taves D (1968) Separation of fluoride by rapid diffusion using hexamethyldisiloxane. *Talanta* **15**, 969–974.
 19. Martínez-Mier EA, Cury JA, Heilman JR *et al.* (2011) Development of gold standard ion-selective electrode-based methods for fluoride analysis. *Caries Res* **45**, 3–12.
 20. Willett WC, Sampson L, Stampfer MJ *et al.* (1985) Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol* **122**, 51–65.
 21. Hernández M, Aguirre J, Serrano L *et al.* (1983) *Alimentación de Obreros y Sus Familias, Publicación L-61 [Nutrition of Workers and Their Families, Publication L-61]*. Mexico: División de Nutrición de Comunidad, Instituto Nacional de la Nutrición Salvador Zubirán.
 22. Hernández-Avila M, Romieu I, Parra S *et al.* (1998) Validity and reproducibility of a food frequency questionnaire to assess dietary intake of women living in Mexico City. *Salud Publica Mex* **40**, 133–140.
 23. Rodríguez-Ramírez S, Mundo-Rosas V, Jiménez-Aguilar A *et al.* (2006) Methodology for the analysis of dietary data from the Mexican national health and nutrition survey. *Salud Publica Mex* **51**, S523–S529.
 24. National Institute of Public Health (2002) *The Compiled México-INSP Food Composition Data Bank*. México, DF: National Institute of Public Health.
 25. Norma Oficial Mexicana NOM-007-SSA2–1993 (1995) Atención de la mujer durante el embarazo, parto y puerperio y del recién nacido. Criterios y procedimientos para la prestación del servicio [Healthcare of women during pregnancy, labor, postpartum and the newborn. Criteria and procedures for the delivery of healthcare services]. <http://www.salud.gob.mx/unidades/cdi/nom/m040ssa193.html#:~:text=1.1%20Esta%20Norma%20Oficial%20Mexicana,sal%20yodada%20para%20consumo%20animal> (accessed January 2021).
 26. Secretaría de Salud & Estados Unidos Mexicanos (2006) Promoción y educación para la salud en materia alimentaria. Criterios para brindar orientación [Maternal nutritional health education and promotion. Criteria to provide guidance]. *Diario Oficial de la Federación* **1**, 32–49.
 27. Jiménez-Zabala A, Santa-Marina L, Otazua M *et al.* (2017) Ingesta de flúor a través del consumo de agua de abastecimiento público en la cohorte INMA-Gipuzkoa [Fluoride intake through consumption of water from municipal network in the INMA-Gipuzkoa cohort]. *Gac Sanit*. Published online: 22 May 2017. doi: 10.1016/j.gaceta.2017.02.008.
 28. Martínez-Salgado H, Tovar-Zamora E, Chavez-Villasana A *et al.* (1993) Consumo familiar e individual de sal de mesa en el Estado de México [Individual and household consumption of table salt in the State of Mexico]. *Salud Publica Mex* **35**, 630–636.
 29. Willett WC (2013) Implications of total energy intake for epidemiologic analyses. In *Nutritional Epidemiology*, 3rd ed., pp. 260–286 [A Hofman, M Marmot, J Samet *et al.*, editors]. New York: Oxford University Press.
 30. Most J, Dervis S, Haman F *et al.* (2019) Energy intake requirements in pregnancy. *Nutrients*. Published online: 6 August 2019. doi: 10.3390/nu11081812.
 31. Saunders CM, Rehbinder EM, Carlsen KCL *et al.* (2019) Food and nutrient intake and adherence to dietary recommendations during pregnancy: a Nordic mother–child population-based cohort. *Food Nutr Res*. Published online: 20 December 2019. doi: 10.29219/fnr.v63.3676.
 32. Anders S & Schroeter C (2017) The impact of nutritional supplement intake on diet behavior and obesity outcomes. *PLoS One*. Published online: 9 October 2017. doi: 10.1371/journal.pone.0185258.
 33. Franco ÁM, Martignon S, Saldarriaga A *et al.* (2005) Total fluoride intake in children aged 22–35 months in four Colombian cities. *Community Dent Oral Epidemiol* **33**, 1–8.