



47th Annual Scientific Meeting of the Nutrition Society of Australia and Nutrition Society of New Zealand, 28 November – 1 December 2023, Nutrition & Wellbeing in Oceania

Ultra-Processed Food Consumption in the Central Division of Fiji

A. Palu¹, J. Santos¹, M. Shahid¹, D. Coyle¹, G. Waqa², A. Moala², C. Bell³ and B.L. McKenzie¹

¹The George Institute for Global Health, University of New South Wales, Sydney, NSW 2042, Australia

²Pacific Research Centre for the Prevention of Obesity and Non-communicable Diseases, Fiji National University, Tamavua Campus, P.O Box 737, Suva, Fiji

³Institute for Health Transformation, Deakin University, Geelong, 3220 Australia

Availability of ultra-processed foods is likely to be high in the Pacific⁽¹⁾ however, information on consumption is limited. This study aimed to assess consumption levels and dietary sources of ultra-processed foods (UPFs) in a population of adults in the Central Division of Fiji. A random sample of 700 adults was selected from two statistical enumeration areas (one semi-urban, one rural) in Fiji. Participant characteristics were collected, along with a three-pass 24-hour diet recall. Foods consumed were coded based on level of processing, in alignment with the NOVA categorisation system (1 = unprocessed, 2 = minimally processed,3 = processed and 4 = ultra-processed). UPF contribution to total energy, salt, fat, and sugar intake were estimated. Main sources of UPFs were then estimated by food group. 534 adults participated (76% response rate, 50% female). Preliminary results suggest that UPFs contributed 21.5% (%95 CI, 19.5% to 23.4%) of total energy intake. Further, UPFs contributed to 22.8% (%95CI 20.5% to 25.1%) of total salt intake, 24.0% (%95 CI, 21.4% to 26.6%) of fat intake and 18.6% (%95 CI, 16.5% to 20.7%) of sugar intake. UPFs contributed over 20% of total energy intake in this sample of Fijian adults and over 20% of salt, fat, and sugar. Messages and interventions that encourage consumption of minimally processed foods while reducing consumption of UPFs are likely needed to improve the healthiness of diets.

Keywords: ultra-processed foods, Fiji, small island developing states, non-communicable diseases

Ethics Declaration

Yes

Financial Support

This research was funded by the National Health and Medical Research Council as part of the Global Alliance for Chronic Diseases program on Scaling up Policy to reduce hypertension and diabetes (APP1169322). BM is supported by a National Heart Foundation of Australia Postdoctoral Fellowship (APP106651). AP is supported by a UNSW Research Training Program (RTP) PhD Scholarship.

Reference

1. Sievert K, Lawrence M, Naika A & Baker P (2019). Nutrients 1(1328).