

CHARACTERIZATION OF CONTINUOUS HOMOMORPHISMS ON ENTIRE SLICE MONOGENIC FUNCTIONS

STEFANO PINTON AND PETER SCHLOSSER

(SP) *Politecnico di Milano, Dipartimento di Matematica, Milano, Italy*

Corresponding author: Stefano Pinton, email: stefano.pinton@polimi.it

(Received 17 June 2023)

Abstract This paper is inspired by a class of infinite order differential operators arising in quantum mechanics. They turned out to be an important tool in the investigation of evolution of superoscillations with respect to quantum fields equations. Infinite order differential operators act naturally on spaces of holomorphic functions or on hyperfunctions. Recently, infinite order differential operators have been considered and characterized on the spaces of entire monogenic functions, i.e. functions that are in the kernel of the Dirac operators. The focus of this paper is the characterization of infinite order differential operators that act continuously on a different class of hyperholomorphic functions, called slice hyperholomorphic functions with values in a Clifford algebra or also slice monogenic functions. This function theory has a very reach associated spectral theory and both the function theory and the operator theory in this setting are subjected to intensive investigations. Here we introduce the concept of proximate order and establish some fundamental properties of entire slice monogenic functions that are crucial for the characterization of infinite order differential operators acting on entire slice monogenic functions.

Keywords: infinite order differential operators; proximate order for slice monogenic functions; continuous homomorphisms

AMS classification: 32A15; 32A10; 47B38

1. Introduction

Extensive research has been conducted on infinite order differential operators for a considerable period of time. In recent years, their significance in examining the evolution of superoscillations as initial data for the Schrödinger equation has emerged as a fundamental area of study. Superoscillatory functions arise in various fields of science and technology. In quantum mechanics, they are the result of Aharonov's weak values, as outlined in [1, 7]. Investigating their time evolution as initial data for quantum field equations represents a crucial problem in the domain of quantum mechanics. Further details can be found in the monograph [5], as well as in [2–4, 6, 13, 14, 18, 19, 37].

Moreover, we point out that superoscillations have been studied not only in quantum mechanics but also in various fields including optics, antenna theory and signal processing. In particular, it is believed that they can be used to improve the resolution of imaging



and spectral analysis techniques, as they allow for obtaining detailed information about structures at smaller scales than those allowed by conventional waves. Superoscillations are still an active area of study, and there are still many open questions about their nature and potential applications.

Investigating the evolution of superoscillatory functions under the time dependent Schrödinger equation presents highly intricate problems. A natural functional analytic framework for this purpose is the space of entire functions with specific growth conditions. In fact, addressing the Cauchy problem for the Schrödinger equation with superoscillatory initial data, we are naturally led to study infinite order differential operators.

In order to explain how infinite order differential operators appear in quantum mechanics, it is enough to consider the simple case of the free particle for Schrödinger equation with initial datum that is a superoscillatory function. Precisely, we consider the Cauchy problem

$$\begin{cases} i \frac{\partial \psi(x,t)}{\partial t} &= -\frac{\partial^2 \psi(x,t)}{\partial x^2} \\ \psi(x,0) &= F_n(x,a) := \sum_{k=0}^n \binom{n}{k} \left(\frac{1+a}{2}\right)^{n-k} \left(\frac{1-a}{2}\right)^k e^{i(1-2k/n)x} \end{cases} \tag{1.1}$$

for $a \in \mathbb{R}$, $a > 1$, $x \in \mathbb{R}$ where $F_n(x,a)$ is a superoscillatory function outcome of Aharonov’s weak values. In [3], it has been shown that the solution $\psi_n(x,t)$ can be written as:

$$\psi_n(x,t) = \sum_{m=0}^{\infty} \frac{(it)^m}{m!} \frac{d^{2m}}{dx^{2m}} F_n(x,a)$$

for every $x \in \mathbb{R}$ and $t \in \mathbb{R}$. To study the limit, as $n \rightarrow \infty$ for $\psi_n(x,t)$ entails the investigation of persistence of superoscillation during the time evolution. The crucial fact is to show the continuity of the operator

$$U \left(\frac{d}{dx} \right) := \sum_{m=0}^{\infty} \frac{(it)^m}{m!} \frac{d^{2m}}{dx^{2m}}$$

on a class of functions that contains $F_n(x,a)$. In order to study the continuity properly, we recall that we have to extend both the operator $U \left(\frac{d}{dx} \right)$ and the function $F_n(x,a)$ to the complex setting replacing the real variable x by the complex variable z . The natural settings for this problem are the spaces \mathcal{A}_p , for $p \geq 1$, i.e. the space of entire functions with either order lower than p or order equal to p and finite type, i.e. it consists of functions f for which there exist constants $B, C > 0$ such that

$$|f(z)| \leq C e^{B|z|^p}. \tag{1.2}$$

The convergence in these spaces is defined as follows. Let $(f_n)_{n \in \mathbb{N}}$, $f_0 \in \mathcal{A}_p$. Then $f_n \rightarrow f_0$ in \mathcal{A}_p if there exists some $B > 0$ such that

$$\lim_{n \rightarrow \infty} \sup_{z \in \mathbb{C}} \left| (f_n(z) - f_0(z)) e^{-B|z|^p} \right| = 0. \tag{1.3}$$

Proving the continuity of the infinite order differential operator $U\left(\frac{d}{dz}\right)$ on \mathcal{A}_1 implies that

$$\lim_{n \rightarrow \infty} U\left(\frac{d}{dz}\right) F_n(z, a) = U\left(\frac{d}{dz}\right) \lim_{n \rightarrow \infty} F_n(z, a).$$

Thus, we have $\lim_{n \rightarrow \infty} \psi_n(z, t) = e^{iaz - ia^2t}$ in \mathcal{A}_1 . Taking the restriction to the real axis for the complex variable z , we have $\lim_{n \rightarrow \infty} \psi_n(x, t) = e^{iax - ia^2t}$ uniformly on the compact sets of \mathbb{R} for the space variable.

More generally when we investigate the Schrödinger evolution of superoscillatory functions under different potentials V , we reduce the problem to the identification of infinite order differential operators of the type

$$\mathcal{U}\left(t, z, \frac{d}{dz}\right) := \sum_{m=0}^{\infty} u_m(t, z; V) \frac{d^m}{dz^m}$$

where z is a complex variable, and where the coefficients $u_m(t, z; V)$ depend on the potential V and also on time t . The identification and the characterization of these type of operators are crucial in quantum mechanics and from the mathematical point of view it has given a new impulse to the theory of infinite order differential operators acting on holomorphic functions, see [9, 12].

The function theory of the spaces \mathcal{A}_p from above can also be treated in a much more general framework where the constant order p is replaced by some function $\varrho(|z|)$ satisfying certain properties. In the complex setting, these spaces of proximate order are introduced in [39]. It was then generalized to several complex variables in [33]. The differential operator representation of continuous homomorphisms between the spaces of entire functions of given proximate order was only recently proven by T. Aoki and co-authors in [10, 11]. In this paper, we will generalize these results for slice monogenic functions acting on a Clifford algebra.

In this regard, there are two main theories of hyperholomorphic functions the monogenic and the slice monogenic function theory. It is important to mention that for these two classes of hyperholomorphic functions we need to define different types of infinite order differential operators involving suitable products of functions that preserve the required notion of hyperholomorphicity. The investigation of infinite order differential operators started in [8] and the characterization of continuous homomorphisms for entire monogenic functions of given proximate order is studied in [27]. In the monogenic case, entire monogenic functions have been considered by different authors in [28–31, 36, 38], while entire slice hyperholomorphic functions are considered in [17].

In order to give the perspective of the two classes of hyperholomorphic functions, referring to the notation introduced in the next section, we denote by \mathbb{R}_n the Clifford algebra with imaginary units $e_j, j = 1, \dots, n$. We recall that (left) monogenic functions, see [16, 35], are those functions $f : U \rightarrow \mathbb{R}_n$ continuously differentiable on an open subset $U \subseteq \mathbb{R}^{n+1}$ such that

$$\mathcal{D}f(x) = 0,$$

where \mathcal{D} is the Dirac operator defined by $\mathcal{D} = \partial_{x_0} + \sum_{j=1}^n e_j \partial_{x_j}$. For the case of slice monogenic functions, there is not a unique way to define them, but there are different ways to introduce them and the various definitions are not totally equivalent according to the properties on the domains on which they are defined. Precisely, we have the following possible definitions:

- (i) Via the Fueter–Sce–Qian mapping theorem, see Definition 2.1.
- (ii) Slice by slice, see [20].
- (iii) As functions in the kernel of the global operator introduced in [26].
- (iv) Via the Radon and dual Radon transform, see [25].

The most natural definition for applications to quaternionic and Clifford operators theory, see [21–23], is the definition appearing in Fueter–Sce–Qian mapping theorem which is well summarized in terms of extensions from holomorphic functions to hyperholomorphic ones. Let $\mathcal{O}(D)$ be the set of holomorphic functions on $D \subseteq \mathbb{C}$ and let $\Omega_D \subseteq \mathbb{R}^{n+1}$ be the rotation of D around the real axis. The first Fueter–Sce–Qian map T_{FS1} applied to $\mathcal{O}(D)$ generates the set $\mathcal{SH}(\Omega_D)$ of slice monogenic functions on Ω_D (which turn out to be intrinsic) and the second Fueter–Sce–Qian map T_{FS2} applied to $\mathcal{SH}(\Omega_D)$ generates axially monogenic functions on Ω_D . We denote this second class of functions by $\mathcal{M}(\Omega_D)$. The extension procedure is illustrated in the diagram:

$$\mathcal{O}(D) \xrightarrow{T_{FS1}} \mathcal{SH}(\Omega_D) \xrightarrow{T_{FS2}=\Delta^{(n-1)/2}} \mathcal{M}(\Omega_D),$$

where $T_{FS2} = \Delta^{(n-1)/2}$ and Δ is the Laplace operator in dimension $n + 1$. For more details on this important extension procedure from complex to hypercomplex analysis see the book [24], the references therein and the paper [32] for related topics.

The plan of the paper. In §2, we have some preliminary results on slice monogenic functions. In §3, we prove new results on some properties of slice monogenic functions of proximate order. Section 4 is the heart of this paper and contains the characterization representation of continuous homomorphisms on spaces of entire slice monogenic functions.

2. Preliminary results on slice monogenic functions

In this section, we recall basic results on slice monogenic functions (see Chapter 2 in [17]) and also introduce the notion of proximate order functions (see the book [33]). We recall that \mathbb{R}_n is the real Clifford algebra over n imaginary units e_1, \dots, e_n , satisfying the commutation relation

$$e_i e_j = -e_j e_i \quad \text{and} \quad e_i^2 = -1, \quad i \neq j \in \{1, \dots, n\}.$$

Any Clifford number $a \in \mathbb{R}_n$ can uniquely be written as

$$a = a_0 + a_1 e_1 + \dots + a_n e_n + \dots + a_{12\dots n} e_1 e_2 \dots e_n = \sum_A a_A e_A,$$

where the last sum is over all ordered subsets $A = \{i_1, \dots, i_r\} \subseteq \{1, \dots, n\}$ and the basis elements are $e_A := e_{i_1} \dots e_{i_r}$. Note that for $A = \emptyset$ we set $e_\emptyset := 1$. The *Euclidean norm* of a Clifford number $a \in \mathbb{R}_n$ is

$$|a|^2 := \sum_A |a_A|^2.$$

Any element of the form

$$x = x_0 + \underline{x} = x_0 + \sum_{\ell=1}^n x_\ell e_\ell$$

will be called *paravector* and we denote by \mathbb{R}^{n+1} the set of all paravectors. Note that no ambiguity arises since we can identify any $(n + 1)$ -vector (x_0, x_1, \dots, x_n) of real numbers with $x_0 + e_1x_1 + \dots + e_nx_n$. The *conjugate* of a paravector x is given by

$$\bar{x} := x_0 - \underline{x} = x_0 - \sum_{\ell=1}^n x_\ell e_\ell.$$

Recall that \mathbb{S} is the sphere

$$\mathbb{S} = \{\underline{x} = e_1x_1 + \dots + e_nx_n \mid x_1^2 + \dots + x_n^2 = 1\},$$

where every $j \in \mathbb{S}$ satisfies $j^2 = -1$. Given an element $x = x_0 + \underline{x} \in \mathbb{R}^{n+1}$ let us define

$$[x] := \{y \in \mathbb{R}^{n+1} : y = x_0 + j|\underline{x}|, j \in \mathbb{S}\},$$

as the $(n - 1)$ -dimensional sphere in \mathbb{R}^{n+1} centred in $x_0 \in \mathbb{R}$ and with radius $|\underline{x}|$. For any $j \in \mathbb{S}$, the vector space

$$\mathbb{C}_j := \{u + jv : u, v \in \mathbb{R}\} = \mathbb{R} + j\mathbb{R}$$

is an isomorphic copy of the complex numbers, passing through the real line in the direction of the imaginary unit j . Finally, a subset $U \subseteq \mathbb{R}^{n+1}$ is called *axially symmetric*, if for every $x \in U$ the whole $(n - 1)$ -sphere $[x] \subseteq U$ is contained in U .

Next we define the set of slice monogenic functions (also called slice hyperholomorphic functions), acting on paravectors \mathbb{R}^{n+1} and having values in the full Clifford algebra \mathbb{R}_n .

Definition 2.1 (Slice monogenic functions). *Let $U \subseteq \mathbb{R}^{n+1}$ be an axially symmetric open set and let $\mathcal{U} = \{(u, v) \in \mathbb{R}^2 : u + \mathbb{S}v \subseteq U\}$. A function $f : \mathcal{U} \rightarrow \mathbb{R}_n$ is called *left slice monogenic*, if it is of the form*

$$f(x) = f_0(u, v) + jf_1(u, v) \quad x = u + jv \in U, \tag{2.1}$$

where the two functions $f_0, f_1 : \mathcal{U} \rightarrow \mathbb{R}_n$ satisfy the compatibility conditions

$$f_0(u, -v) = f_0(u, v) \quad \text{and} \quad f_1(u, -v) = -f_1(u, v), \tag{2.2}$$

as well as the Cauchy–Riemann equations

$$\frac{\partial}{\partial u} f_0(u, v) = \frac{\partial}{\partial v} f_1(u, v) \quad \text{and} \quad \frac{\partial}{\partial v} f_0(u, v) = -\frac{\partial}{\partial u} f_1(u, v). \tag{2.3}$$

The set of left slice monogenic functions will be denoted by $SM_L(U)$.

Remark 2.2. Analogously, one can also define right slice monogenic functions by simply replacing the decomposition (2.1) by

$$f(s) = f_0(u, v) + f_1(u, v)j \quad x = u + jv \in U.$$

However, for the rest of the paper, we will only consider left slice monogenic functions, although all the results can also be written for right slice monogenic functions.

Definition 2.3. Let $f \in SM_L(U)$. Then for any $x \in U$ we define the left slice derivative as

$$\partial_S f(x) := \lim_{p \rightarrow x, p \in \mathbb{C}_j} (p - x)^{-1} (f(p) - f(x)), \tag{2.4}$$

where for non-real x the imaginary unit $j \in \mathbb{S}$ is chosen such that $x \in \mathbb{C}_j$ and for real x one may choose $j \in \mathbb{S}$ arbitrary.

We remark that $\partial_S f(x)$ is uniquely defined and independent of the choice of $j \in \mathbb{S}$, even if x is real. Moreover, we note that the slice derivative of f coincides with

$$\partial_S f(x) = \partial_{x_0} f(x) = f'_j(x), \tag{2.5}$$

where $\partial_{x_0} f$ is the partial derivative with respect to x_0 and f'_j is the complex derivative of the restriction $f_j = f|_{\mathbb{C}_j}$.

Theorem 2.4. For $a \in \mathbb{R}$, $r > 0$ let $B_r(a) = \{x \in \mathbb{R}^{n+1} : |x - a| < r\}$. If $f \in SM_L(B_r(a))$, then

$$f(x) = \sum_{k=0}^{\infty} (x - a)^k \frac{1}{k!} \partial_{x_0}^k f(a) \quad x \in B_r(a). \tag{2.6}$$

We now recall the natural product of two functions that preserves slice monogenicity.

Definition 2.5. Let $U \subseteq \mathbb{R}^{n+1}$ be open and axially symmetric. Then for any $f, g \in SM_L(U)$ define their star product

$$f \star_L g := f_0 g_0 - f_1 g_1 + j(f_1 g_0 + f_0 g_1),$$

with the functions f_0, f_1, g_0, g_1 from the decomposition (2.1).

Lemma 2.6. *Let $f(x) = \sum_{k=0}^{\infty} x^k a_k$ and $g(x) = \sum_{k=0}^{\infty} x^k b_k$ be two left slice monogenic power series. Then their star product is given by*

$$(f \star_L g)(x) = \sum_{\ell=0}^{\infty} x^{\ell} \sum_{k=0}^{\ell} a_k b_{\ell-k}. \tag{2.7}$$

For $x, s \in \mathbb{R}^{n+1}$ with $x \notin [s]$, the Cauchy kernel for left slice monogenic functions is given by

$$S_L^{-1}(s, x) := -(x^2 - 2s_0x + |s|^2)^{-1}(x - \bar{s}) = (s - \bar{x})(s^2 - 2x_0s + |x|^2)^{-1}. \tag{2.8}$$

With this kernel, there holds the following Clifford algebra version of the Cauchy integral formula.

Theorem 2.7 (The Cauchy formula). *Let $U \subset \mathbb{R}^{n+1}$ be axially symmetric, open, bounded and the boundary $\partial(U \cap \mathbb{C}_j)$ be a finite union of continuously differentiable Jordan curves. If we set $ds_j = -j ds$ for some $j \in \mathbb{S}$, then for f which is left slice monogenic on a neighbourhood of \bar{U} , there holds*

$$f(x) = \frac{1}{2\pi} \int_{\partial(U \cap \mathbb{C}_j)} S_L^{-1}(s, x) ds_j f(s), \quad x \in U. \tag{2.9}$$

Moreover, the integral depend neither on U nor on the imaginary unit $j \in \mathbb{S}$.

After the basic facts on slice monogenic functions, we introduce the notion of proximate order functions which will then be used in § 3 to introduce function spaces of exponentially bounded entire slice monogenic functions.

Definition 2.8 (Proximate order). *A differentiable function $\varrho : [0, \infty) \rightarrow [0, \infty)$ is called proximate order function for the order $\rho > 0$, if it satisfies*

- (1) $\lim_{r \rightarrow \infty} \varrho(r) =: \rho > 0$,
- (2) $\lim_{r \rightarrow \infty} \varrho'(r)r \ln(r) = 0$.

It is also possible to define proximate order for the order $\rho = 0$, but the upcoming results do not hold in this case. Any proximate order function admits the following properties. A proof can for example be found in [33, Proposition 1.19].

Lemma 2.9 (Properties of proximate orders). *Let ϱ be a proximate order function. Then there exists some $r_0 > 0$, such that*

- (i) $r \mapsto r^{\varrho(r)}$ is strictly increasing on $[r_0, \infty)$,
- (ii) $\lim_{r \rightarrow \infty} r^{\varrho(r)} = \infty$.

Definition 2.10 (Normalized proximate order). A proximate order function ϱ is called normalized if

- (i) $r \mapsto r^{\varrho(r)}$ is strictly increasing on $(0, \infty)$,
- (ii) $\lim_{r \rightarrow 0^+} r^{\varrho(r)} = 0$.

For a normalized proximate order function, it is then clear that $r \mapsto r^{\varrho(r)}$ maps $(0, \infty)$ bijectively onto $(0, \infty)$ and we denote by

$$\varphi : (0, \infty) \rightarrow (0, \infty) \text{ the inverse function of } r \mapsto r^{\varrho(r)}. \tag{2.10}$$

Moreover, we set the numbers

$$G_0 = G_{\varrho,0} := 1 \quad \text{and} \quad G_\ell = G_{\varrho,\ell} := \frac{\varphi(\ell)^\ell}{(e\rho)^\ell/\rho}, \quad \ell \in \mathbb{N}. \tag{2.11}$$

Remark 2.11. Note that by Lemma 2.9 it is possible to construct for any given proximate order function ϱ a normalized proximate order function $\hat{\varrho}$ such that $\varrho(r) = \hat{\varrho}(r)$ for every $r \geq r_0$ large enough. One possible choice is the function

$$\hat{\varrho}(r) := \begin{cases} \varrho(r_0) - \frac{\rho}{4} \sin\left(4 \frac{\varrho'(r_0)}{\rho}(r_0 - r)\right), & r \in [0, r_0], \\ \varrho(r), & r \in [r_0, \infty). \end{cases}$$

Next, we recall some basic properties of proximate order functions that will be used in the following. First we prove two inequalities of the mapping $r \mapsto r^{\varrho(r)}$. The first one (i) can be found in [10, Lemma 2.3] and the second one (ii) in [10, Proposition 1.20].

Lemma 2.12. Let ϱ be a normalized proximate order function. Then for any $\varepsilon > 0$ there exists a constant $C_\varepsilon \geq 0$, such that

- (i) $(r + s)^{\varrho(r+s)} \leq 2^{\rho+\varepsilon} (r^{\varrho(r)} + s^{\varrho(s)}) + C_\varepsilon, \quad r, s > 0.$
- (ii) $(sr)^{\varrho(sr)} \leq (1 + \varepsilon)s^\rho r^{\varrho(r)} + C_\varepsilon, \quad r, s > 0.$

The next lemma can be found in [10, Lemma 2.4] and it is the submultiplicativity of the numbers G_ℓ in (2.11).

Lemma 2.13. The sequence $(G_\ell)_{\ell \in \mathbb{N}_0}$ from (2.11) satisfies

$$G_\ell G_k \leq G_{\ell+k}, \quad \ell, k \in \mathbb{N}_0.$$

The following Lemma treats the limit behaviour of the function φ from Definition 2.10. The upcoming results (i) and (ii) can be found in the proof of [33, Theorem 1.23], the result (iii) in [33, Proposition 1.20] and (iv) in [10, Lemma 2.6].

Lemma 2.14. *Let ϱ be a normalized proximate order function and φ from (2.10). Then for every $s > 0$ and $0 < \sigma' < \sigma$ there holds*

- (i) $\lim_{t \rightarrow \infty} \frac{t\varphi'(t)}{\varphi(t)} = \frac{1}{\varrho},$
- (ii) $\lim_{t \rightarrow \infty} \frac{\varphi(st)}{\varphi(t)} = s^{\frac{1}{\varrho}},$
- (iii) $\lim_{r \rightarrow \infty} \frac{(sr)^{\varrho(sr)}}{r^{\varrho(r)}} = s^\rho.$
- (iv) $\exists t_0 > 0 : \frac{\varphi(t)}{\varphi(t')} \leq \frac{e^{\frac{\sigma}{t'}}}{(e\sigma')^{1/\varrho}}, \quad t, t' \geq t_0.$

3. Slice monogenic functions of proximate order

Let $\varrho(r)$ be a proximate order function for a positive order $\rho > 0$ according to Definition 2.8. We will now introduce some function spaces of entire slice monogenic functions in the spirit of [10, 33], which treat the similar problem for entire functions of several complex variables. For constant proximate order functions $\varrho(r) = \rho$ and slice monogenic functions in the quaternions, we also refer to the results in [17, Chapter 5]. In particular, we derive basic properties of these function spaces where most of them are already known in the complex and the monogenic setting but which are new in the case of slice monogenic functions. Moreover, we give an alternative and more detailed proof of [33, Theorem 1.23] in Theorem 3.7.

For any $\sigma > 0$, we consider the Banach space

$$A_{\varrho, \sigma} := \left\{ f \in \mathcal{SM}_L(\mathbb{R}^{n+1}) : \|f\|_{\varrho, \sigma} := \sup_{x \in \mathbb{R}^{n+1}} |f(x)| \exp(-\sigma|x|^{\varrho(|x|)}) < \infty \right\}$$

with the norm $\|\cdot\|_{\varrho, \sigma}$.

Remark 3.1. We observe that for any proximate function $\varrho(r)$ and any normalization function $\hat{\varrho}(r)$ according to Remark 2.11, the spaces $A_{\varrho, \sigma}$ and $A_{\hat{\varrho}, \sigma}$ coincide with equivalent norms, see [10, Page 8].

Lemma 3.2. *If $\sigma_2 > \sigma_1 > 0$, then the inclusion map $A_{\varrho, \sigma_1} \hookrightarrow A_{\varrho, \sigma_2}$ is compact.*

Proof. We will show that $B := \{f \in A_{\varrho, \sigma_1} : \|f\|_{\varrho, \sigma_1} \leq 1\}$ is relatively compact in A_{ϱ, σ_2} , i.e. any sequence $(f_k)_{k \in \mathbb{N}} \subset B$ has an accumulation point with respect to the norm of A_{ϱ, σ_2} .

First we will prove that the sequence $(f_k)_{k \in \mathbb{N}} \subset B$ admits a subsequence which converges in the uniform convergence topology of \mathbb{R}^{n+1} . By the Arzelá–Ascoli theorem, together with some standard diagonal sequence argument, it is sufficient to prove that $(f_k)_{k \in \mathbb{N}}$ is equicontinuous and uniformly bounded on any compact convex subset $K \subseteq \mathbb{R}^{n+1}$. Let us now fix one of the compact convex subsets K . Since $(f_k)_{k \in \mathbb{N}} \subset B$, the sequence is uniformly bounded on K . Moreover, we have

$$|f_k(x) - f_k(y)| \leq C_k|x - y|, \quad x, y \in K$$

where $C_k = \sup_{x \in K} |\nabla f_k(x)|$ and ∇ is the usual gradient. We choose r large enough in a such way that $K \subset B(0, r)$. It is sufficient to prove that there exists a constant C_K , only depending on K , such that

$$|\partial_{x_i} f_k(x)| \leq C_K, \quad x \in K, k \in \mathbb{N}, i = 0, \dots, n.$$

To prove this fact we need to differentiate the integral representation formula (2.9) for f_k . Let $x \in K$ and $j \in \mathbb{S}$ be such that there exist $u, v \in \mathbb{R}$ with $x = u + jv$. When $i = 0$, there exists a positive constant C'_K which depends only on K , such that

$$\begin{aligned} |\partial_{x_0} f_k(x)| &= \left| \frac{1}{2\pi} \int_{\partial(B(0,r) \cap \mathbb{C}_j)} \partial_{x_0} S_L^{-1}(s, x) \, ds_j f_k(s) \right| \\ &= \frac{1}{2\pi} \left| \int_{\partial(B(0,r) \cap \mathbb{C}_j)} \frac{1}{(s-x)^2} \, ds_j f_k(s) \right| \\ &\leq \frac{1}{2\pi} \int_{\partial(B(0,r) \cap \mathbb{C}_j)} \frac{1}{|x-s|^2} |ds_j| M(r, f_k) \\ &\leq r \frac{M(r, f_k)}{(r-|x|)^2} \leq C'_K, \end{aligned}$$

where in the last line we used $(f_k)_{k \in \mathbb{N}} \subset B$ and $\text{dist}(K, \partial B(0, r)) > 0$. When $i \neq 0$, using the second form for $S_L^{-1}(s, x)$ in (2.8), we observe that

$$\partial_{x_i} S_L^{-1}(s, x) = (e_i s^2 - 2x_0 e_i s + |x|^2 e_i - 2x_i s + 2x_i \bar{x})(s^2 - 2x_0 s + |x|^2)^{-2}.$$

Thus, when $x \in \mathbb{C}_j \cap K$ and $s \in \mathbb{C}_j \cap \partial B(0, r)$, there exists a positive constant C_2 , which depends only on K , such that

$$|\partial_{x_i} S_L^{-1}(s, x)| \leq \frac{C_2}{|s-x|^2 |s-\bar{x}|^2}.$$

By this inequality, there exists another positive constant C''_K , which only depends on K , such that

$$\begin{aligned} |\partial_{x_i} f_k(x)| &= \left| \frac{1}{2\pi} \int_{\partial(B(0,r) \cap \mathbb{C}_j)} \partial_{x_i} (S_L^{-1}(s, x)) \, ds_j f_k(s) \right| \\ &\leq \frac{1}{2\pi} \int_{\partial(B(0,r) \cap \mathbb{C}_j)} |\partial_{x_i} S_L^{-1}(s, x)| |ds_j| M(r, f_k) \\ &\leq r C_2 \frac{M(r, f_k)}{(r-|x|)^4} \leq C''_K. \end{aligned}$$

In particular, choosing $C_K = \max\{C'_K, C''_K\}$, for any $x, y \in K$ and for any $k \in \mathbb{N}$, we have

$$|f_k(x) - f_k(y)| \leq \sqrt{n} C_K |x - y|,$$

i.e. $(f_k)_{k \in \mathbb{N}}$ is equicontinuous on K . Applying the Arzelá–Ascoli theorem and using some standard diagonal sequence argument, there exists a subsequence which converges to $f \in \mathcal{SM}_L(\mathbb{R}^{n+1})$ in the topology of the uniform convergence on compact subsets. Without loss of generality, we call this subsequence again by $(f_k)_{k \in \mathbb{N}}$.

In the second step, we prove that the sequence $(f_k)_{k \in \mathbb{N}}$ is a Cauchy sequence in A_{ϱ, σ_2} . We fix $\delta > 0$ and choose $R > 0$ large enough such that

$$\exp((\sigma_1 - \sigma_2)|x|^{e(|x|)}) \leq \frac{\delta}{2}, \quad \text{for any } |x| \geq R.$$

Thus, since $f_k, f_\ell \in B$, we have

$$\begin{aligned} & \sup_{|x| \geq R} |f_k(x) - f_\ell(x)| \exp(-\sigma_2|x|^{e(|x|)}) \\ &= \sup_{|x| \geq R} |f_k(x) - f_\ell(x)| \exp(-\sigma_1|x|^{e(|x|)}) \exp((\sigma_1 - \sigma_2)|x|^{e(|x|)}) \\ &\leq 2 \cdot \frac{\delta}{2} = \delta. \end{aligned} \tag{3.1}$$

Moreover, by the uniform convergence of the sequence $(f_k)_{k \in \mathbb{N}}$ on the compact subset $\overline{B(0, R)}$ of \mathbb{R}^{n+1} , there exists a positive integer N such that for any $k, \ell \geq N$ we have

$$\sup_{|x| \leq R} |f_k(x) - f_\ell(x)| \exp(-\sigma_2|x|^{e(|x|)}) \leq \sup_{|x| \leq R} |f_k(x) - f_\ell(x)| \leq \delta. \tag{3.2}$$

Thus, combining (3.1) and (3.2), we have proved that the sequence $(f_k)_{k \in \mathbb{N}}$ is a Cauchy sequence in A_{ϱ, σ_2} . □

Definition 3.3. (The spaces A_ϱ and $A_{\varrho, \sigma+0}$). We define the space

$$A_\varrho := \lim_{\sigma \rightarrow 0} A_{\varrho, \sigma}$$

i.e. $A_\varrho = \cup_{\sigma > 0} A_{\varrho, \sigma}$ and we say that a sequence $(f_k)_{k \in \mathbb{N}} \subseteq A_\varrho$ converges to $f \in A_\varrho$ if there exists some $\sigma > 0$ such that $(f_k)_{k \in \mathbb{N}} \subseteq A_{\varrho, \sigma}$, $f \in A_{\varrho, \sigma}$ and $\lim_{k \rightarrow +\infty} \|f_k - f\|_{\varrho, \sigma} = 0$.

For every $\sigma \geq 0$, we also define the space

$$A_{\varrho, \sigma+0} := \lim_{\epsilon \rightarrow 0} A_{\varrho, \sigma+\epsilon},$$

i.e. $A_{\varrho, \sigma+0} := \cap_{\epsilon > 0} A_{\varrho, \sigma+\epsilon}$ and we say that a sequence $(f_k)_{k \in \mathbb{N}} \subseteq A_{\varrho, \sigma+0}$ converges to $f \in A_{\varrho, \sigma+0}$ if for any $\epsilon > 0$ we have $(f_k)_{k \in \mathbb{N}} \subseteq A_{\varrho, \sigma+\epsilon}$, $f \in A_{\varrho, \sigma+\epsilon}$ and $\lim_{k \rightarrow +\infty} \|f_k - f\|_{\varrho, \sigma+\epsilon} = 0$. When $\sigma = 0$, we denote $A_{\varrho, +0} := A_{\varrho, 0+0}$.

Note that A_ϱ is a (DFS)-space (see [15, Definition 2.2.1 and §2.6]) and $A_{\varrho, \sigma+0}$ is a (FS)-space. Both A_ϱ and $A_{\varrho, \sigma+0}$ are locally convex spaces.

Remark 3.4. Let ϱ be a proximate order function and $\hat{\varrho}$ its normalization according to Remark 2.11. Then the spaces A_ϱ and $A_{\hat{\varrho}}$ coincide and share the same locally convex topology. The same holds true for $A_{\varrho, \sigma+0}$ and $A_{\hat{\varrho}, \sigma+0}$.

Definition 3.5. Let ϱ be a proximate order function and $f \in A_{\varrho}$. Then we define the type of f with respect to ϱ as

$$\inf\{\beta > 0 : f \in A_{\varrho,\beta}\}.$$

Next we prove a preparatory lemma for Theorem 3.7, which characterizes the order.

Lemma 3.6. Let ϱ be a normalized proximate order function for the positive order ρ . Let $f(x) = \sum_{\ell=0}^{\infty} x^{\ell} a_{\ell} \in A_{\varrho}$. Moreover, suppose that $\sigma \geq 0$ which satisfies the property

$$\frac{1}{\rho} \ln(\sigma) \geq \limsup_{\ell \rightarrow \infty} \left(\frac{1}{\ell} \ln |a_{\ell}| + \ln(\varphi(\ell)) \right) - \frac{1}{\rho} - \frac{\ln(\rho)}{\rho},$$

where we interpret $\ln(0) = -\infty$ in the case $\sigma = 0$. Then, for any $\tau' > \sigma$, there exist positive constants N and C such that

$$\sup_{\ell \geq N} (\ln |a_{\ell}| + \ell \ln(r)) \leq \tau' r^{\varrho(r)} + C \quad r > 0. \tag{3.3}$$

Proof. Choose $\sigma < \sigma' < \sigma'' < \tau < \tau'$ arbitrary. Then, by assumption, there exists $N_1 \in \mathbb{N}$, such that

$$|a_{\ell}|^{\frac{1}{\ell}} \varphi(\ell) \leq (e\rho\sigma')^{\frac{1}{\rho}}, \quad \ell \geq N_1.$$

Moreover, by the limit $\lim_{\ell \rightarrow \infty} \frac{\varphi(\frac{\ell}{\rho\sigma''})}{\varphi(\ell)} = (\frac{1}{\rho\sigma''})^{\frac{1}{\rho}}$ from Lemma 2.14 (ii), there exists some $N_2 \geq N_1$ such that $\frac{\varphi(\frac{\ell}{\rho\sigma''})}{\varphi(\ell)} \leq (\frac{1}{\rho\sigma'})^{\frac{1}{\rho}}$ and hence

$$|a_{\ell}|^{\frac{1}{\ell}} \leq \frac{(e\rho\sigma')^{\frac{1}{\rho}}}{\varphi(\ell)} \leq \frac{e^{\frac{1}{\rho}}}{\varphi(\frac{\ell}{\rho\sigma''})}, \quad \ell \geq N_2. \tag{3.4}$$

Next, by Lemma 2.14 (i), there exists some $N \geq N_2$ such that

$$-1 \leq \frac{\frac{t}{\sigma''\rho} \varphi'(\frac{t}{\sigma''\rho})}{\varphi(\frac{t}{\sigma''\rho})} - \frac{1}{\rho} \leq \frac{\tau - \sigma''}{\rho\sigma''}, \quad t \geq N. \tag{3.5}$$

We will now prove that there exists a positive constant C , which does not depend on r , and $r_0 > 0$, such that

$$\sup_{\ell \geq N} (\ln |a_{\ell}| + \ell \ln(r)) \leq \tau r^{\varrho(r)} + C, \quad r > r_0. \tag{3.6}$$

Due to the estimate (3.4), we get

$$\begin{aligned} \sup_{\ell \geq N} (\ln |a_\ell| + \ell \ln(r)) &\leq \sup_{\mathbb{N} \ni \ell \geq N} \ell \left(\frac{1}{\rho} - \ln \left(\varphi \left(\frac{\ell}{\sigma'' \rho} \right) \right) + \ln(r) \right) \\ &\leq \sup_{\mathbb{R} \ni t \geq N} \underbrace{t \left(\frac{1}{\rho} - \ln \left(\varphi \left(\frac{t}{\sigma'' \rho} \right) \right) + \ln(r) \right)}_{=: \mu_r(t)}. \end{aligned} \tag{3.7}$$

Let $t_{\max}(r)$ be the supremum of the points where the function $\mu_r(\cdot)$ attains its maximum. We observe that this point exists and it is finite since $\mu_r(t)$ is continuous and converges to $-\infty$ when $t \rightarrow \infty$. First we prove that $t_{\max}(r) \rightarrow +\infty$ when $r \rightarrow +\infty$. We observe that

$$\mu'_r(t) = \frac{1}{\rho} - \ln \left(\varphi \left(\frac{t}{\sigma'' \rho} \right) \right) + \ln(r) - \frac{\frac{t}{\sigma'' \rho} \varphi' \left(\frac{t}{\sigma'' \rho} \right)}{\varphi \left(\frac{t}{\sigma'' \rho} \right)}.$$

We assume by contradiction that $t_{\max}(r)$ is bounded. Since $\mu'_r(t_{\max}(r)) \leq 0$, we have

$$\ln \left(\varphi \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right) \right) \geq \frac{1}{\rho} + \ln(r) - \frac{\frac{t_{\max}(r)}{\sigma'' \rho} \varphi' \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right)}{\varphi \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right)}. \tag{3.8}$$

The previous inequality gives a contradiction since the left hand side is bounded for any $r > 0$, instead the right hand side tends to $+\infty$ when $r \rightarrow +\infty$.

Since we just have proven that $t_{\max}(r) \xrightarrow{r \rightarrow \infty} \infty$ there exists some r_0 such that $t_{\max}(r) > N$ for any $r \geq r_0$ and hence also $\mu'_r(t_{\max}(r)) = 0$ has to be satisfied for any $r \geq r_0$. This means that the inequality (3.8) becomes an equation, i.e.

$$\ln \left(\frac{1}{r} \varphi \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right) \right) = \frac{1}{\rho} - \frac{\frac{t_{\max}(r)}{\sigma'' \rho} \varphi' \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right)}{\varphi \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right)}.$$

In view of Lemma 2.14 (i), we have

$$\lim_{r \rightarrow \infty} \ln \left(\frac{1}{r} \varphi \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right) \right) = 0.$$

Next we choose $\epsilon, \eta > 0$ such that $(1 + \eta)(1 + \epsilon)^\rho \tau \leq \tau'$. By the previous limit, we can enlarge $r_0 > 0$ such that

$$\varphi \left(\frac{t_{\max}(r)}{\sigma'' \rho} \right) \leq (1 + \epsilon)r, \quad r \geq r_0.$$

By applying the inverse function $\varphi^{-1}(r) = r^{\varrho(r)}$ to both sides of the this inequality and by Lemma 2.12 (ii) with the above chosen η , we get

$$t_{\max}(r) \leq \sigma'' \rho ((1 + \epsilon)r)^{\varrho((1 + \epsilon)r)} \leq \sigma'' \rho (1 + \eta) (1 + \epsilon)^\rho r^{\varrho(r)} + \sigma'' \rho C_\eta. \tag{3.9}$$

Moreover, rearranging the equation $\mu'_r(t_{\max}(r)) = 0$ and using the second inequality in (3.5), we have that

$$\ln\left(\varphi\left(\frac{t_{\max}(r)}{\sigma''\rho}\right)\right) = \ln(r) + \frac{1}{\rho} - \frac{\frac{t_{\max}(r)}{\sigma''\rho}\varphi'\left(\frac{t_{\max}(r)}{\sigma''\rho}\right)}{\varphi\left(\frac{t_{\max}(r)}{\sigma''\rho}\right)} \geq \ln(r) - \frac{\tau - \sigma''}{\rho\sigma''}. \tag{3.10}$$

Using the inequalities (3.9) and (3.10) in (3.7), we obtain for every $r \geq r_0$

$$\begin{aligned} \sup_{\ell \geq N} (\ln |a_\ell| + \ell \ln(r)) &= \mu_r(t_{\max}(r)) \\ &= t_{\max}(r) \left(\frac{1}{\rho} - \ln\left(\varphi\left(\frac{t_{\max}(r)}{\sigma''\rho}\right)\right) + \ln(r) \right) \\ &\leq (\sigma''\rho(1 + \eta)(1 + \epsilon)^\rho r^{\varrho(r)} + \sigma''\rho C_\eta) \left(\frac{1}{\rho} - \ln(r) + \frac{\tau - \sigma''}{\rho\sigma''} + \ln(r) \right) \\ &\leq \tau' r^{\varrho(r)} + \tau' C_\eta. \end{aligned} \tag{3.11}$$

Since $\ln(r)$ is increasing we furthermore get the estimate

$$\sup_{\ell \geq N} (\ln |a_\ell| + \ell \ln(r)) \leq \sup_{\ell \geq N} (\ln |a_\ell| + \ell \ln(r_0)) \leq \tau' r_0^{\varrho(r_0)} + \tau' C_\eta, \quad r \in (0, r_0].$$

Hence, choosing $C' = \tau' r_0^{\varrho(r_0)} + \tau' C_\eta$, we finally have

$$\sup_{\ell \geq N} (\ln |a_\ell| + \ell \ln(r)) \leq \tau' r^{\varrho(r)} + C' \quad r > 0. \quad \square$$

Next we prove the main theorem of this section, a characterization of functions in the spaces $A_{\varrho, \sigma+0}$ with respect to the order, their growth condition and their Taylor coefficients.

Theorem 3.7. *Let ϱ be a normalized proximate order function, $\sigma \geq 0$ and $f(x) = \sum_{\ell=0}^{+\infty} x^\ell a_\ell \in \mathcal{SM}_L(\mathbb{R}^{n+1})$. Then the following four statements are equivalent:*

- (1) $f \in A_{\varrho, \sigma+0}$;
- (2) $\limsup_{r \rightarrow +\infty} \frac{\sup_{|x| \leq r} \ln(|f(x)|)}{r^{\varrho(r)}} \leq \sigma$;
- (3) $\limsup_{\ell \rightarrow \infty} |a_\ell|^{\frac{1}{\ell}} \varphi(\ell) \leq (e\rho\sigma)^{\frac{1}{\rho}}$;
- (4) $\inf\{\beta > 0 : f \in A_{\varrho, \beta}\} \leq \sigma$.

Remark 3.8. Note that using the numbers $(G_\ell)_{\ell \in \mathbb{N}_0}$ from (2.11), one can alternatively write (3) as

$$\limsup_{\ell \rightarrow \infty} (|a_\ell| G_{\varrho, \ell})^{\frac{\rho}{\ell}} \leq \sigma.$$

Proof of Theorem 3.7 We start with (1) \Rightarrow (2). Since $f \in A_{\varrho, \sigma+0}$, there exists for any $\epsilon > 0$ a $D_\epsilon > 0$ such that

$$|f(x)| \leq D_\epsilon \exp((\sigma + \epsilon)|x|^{\varrho(|x|)}) \quad \text{for all } x \in \mathbb{R}^{n+1}. \tag{3.12}$$

Taking the logarithm on both sides of (3.12) we have

$$\ln(|f(x)|) \leq \ln(D_\epsilon) + (\sigma + \epsilon)|x|^{\varrho(|x|)}.$$

Let $r > 0$ be arbitrary, this implies that

$$\sup_{|x| < r} \ln(|f(x)|) \leq \ln(D_\epsilon) + (\sigma + \epsilon)r^{\varrho(r)}.$$

Since $r^{\varrho(r)} \rightarrow +\infty$ as $r \rightarrow +\infty$, we have

$$\limsup_{r \rightarrow +\infty} \frac{\sup_{|x| \leq r} \ln(|f(x)|)}{r^{\varrho(r)}} \leq \sigma + \epsilon. \tag{3.13}$$

Because inequality (3.13) holds to be true for any $\epsilon > 0$, we get (2), i.e.

$$\limsup_{r \rightarrow +\infty} \frac{\sup_{|x| \leq r} \ln(|f(x)|)}{r^{\varrho(r)}} \leq \sigma.$$

For the implication (2) \Rightarrow (4), let $\tau > \sigma$, which by the assumption (2) also means that $\tau > \limsup_{r \rightarrow +\infty} \frac{\sup_{|x| \leq r} \ln(|f(x)|)}{r^{\varrho(r)}}$. Then we can choose $r_0 > 0$ such that for any $|x| > r_0$ we have

$$|f(x)| \leq \exp(\tau|x|^{\varrho(|x|)}).$$

Thus there exists a positive constant C such that $|f(x)| \leq C \exp(\tau|x|^{\varrho(|x|)})$ for any $x \in \mathbb{R}^{n+1}$, i.e. $f \in A_{\varrho, \tau}$. Since this is true for every $\tau > \sigma$, this shows (4).

Now we prove (4) \Rightarrow (3). Assuming that (4) holds to be true, then for any $\epsilon > 0$ there exists $\beta > 0$, such that $\beta < \sigma + \epsilon$ and $f \in A_{\varrho, \beta}$. Then for any $r > 0$, we can estimate the Taylor coefficients a_ℓ of f using the Cauchy integral formula

$$|a_\ell| \leq \frac{1}{2\pi} \int_0^{2\pi} r^{-\ell-1} r |f(re^{j\theta})| d\theta \leq \frac{\|f\|_{\varrho, \beta} \exp(\beta r^{\varrho(r)})}{r^\ell}.$$

Using this inequality with the specific value $r = \varphi(\frac{\ell}{\beta\rho})$, i.e. the one $r > 0$ such that $r^{\varrho(r)} = \frac{\ell}{\beta\rho}$, gives

$$|a_\ell| \leq \frac{\|f\|_{\varrho, \beta} \exp(\frac{\ell}{\rho})}{\varphi(\frac{\ell}{\beta\rho})^\ell}.$$

Taking this inequality to the power $\frac{1}{l}$ and multiplying $\varphi(l)$, it becomes

$$\varphi(l)|a_l|^{\frac{1}{l}} \leq \frac{\|f\|_{\varrho,\beta}^{\frac{1}{l}} e^{\frac{1}{p}} \varphi(l)}{\varphi(\frac{l}{\beta\rho})}.$$

Taking now the $\limsup_{l \rightarrow \infty}$ and using the limit $\lim_{l \rightarrow \infty} \frac{\varphi(l)}{\varphi(\frac{l}{\beta\rho})} = (\beta\rho)^{\frac{1}{p}}$ from Lemma 2.14 (ii), gives

$$\limsup_{l \rightarrow +\infty} \varphi(l)|a_l|^{\frac{1}{l}} \leq (e\beta\rho)^{\frac{1}{p}}.$$

However, since $\beta \leq \sigma + \varepsilon$ and $\varepsilon > 0$ is arbitrary, there also holds

$$\limsup_{l \rightarrow +\infty} \varphi(l)|a_l|^{\frac{1}{l}} \leq (e\rho\sigma)^{\frac{1}{p}},$$

which is exactly (3).

For the last implication (3) \Rightarrow (1) let $\tau > \sigma$ and choose $\tau > \tau' > \sigma$. We want to show that $f \in A_{\varrho,\tau}$ by estimating in a suitable way $\sum_{\ell=0}^{+\infty} |a_\ell||x|^\ell$. In what follows we are going to split the previous summation in three parts. We know that

$$\limsup_{l \rightarrow +\infty} \ln \left(\varphi(l)|a_l|^{\frac{1}{l}} \right) \leq \ln \left((e\sigma\rho)^{\frac{1}{p}} \right).$$

Thus by Lemma 3.6, there exist positive constants N and C such that

$$\sup_{l \geq N} \left(\ln |a_l| + l \ln(r) \right) \leq \tau' r^{e(r)} + C \quad r > 0.$$

Moreover, let $\rho_1 > \rho$ and fix $x \in \mathbb{R}^{n+1}$ with $r := |x| \geq r_0$ for some $r_0 > 0$ large enough. Then define $m_r := \lfloor 2e\tau\rho_1 r^{\rho_1} \rfloor$. As it is shown in (3.4) in such a way that for any $l \geq m_r$ we have

$$|a_l|^{\frac{1}{l}} \leq \frac{(e\rho_1\tau)^{\frac{1}{p_1}}}{\varphi(l)} \leq \frac{2^{-1}}{\varphi(\frac{l}{2e\tau\rho_1})} \leq \frac{2^{-1}}{\varphi(r^{\rho_1})} \leq \frac{2^{-1}}{\varphi(r^{e(r)})} = \frac{2^{-1}}{r}, \quad l \geq m_r. \tag{3.14}$$

Hence we have

$$|f(x)| \leq \sum_{\ell=0}^{N-1} |a_\ell||x|^\ell + \sum_{\ell=N}^{m_r} |a_\ell||x|^\ell + \sum_{\ell=m_r+1}^{+\infty} |a_\ell||x|^\ell.$$

We want to estimate all the three terms of the previous summation. Since the number of terms in the first summation is finite and it does not depend on $|x|$, there exists a

positive constant C'' that depends only on N such that for any $r > 0$ we have

$$\sum_{\ell=0}^{N-1} |a_\ell||x|^\ell \leq C'' \exp(\tau'|x|^{\varrho(|x|)}).$$

For the second summation, using (3.3), we get

$$\sum_{\ell=N}^{m_r} |a_\ell||x|^\ell \leq m_r e^C \exp(\tau'|x|^{\varrho(|x|)}).$$

For the third equation, we use (3.14) to obtain the estimate

$$\sum_{\ell=m_r+1}^{+\infty} |a_\ell||x|^\ell \leq \sum_{\ell=0}^{\infty} 2^{-\ell} = 2.$$

Summing up the three previous inequalities, there exists a positive constant C such that

$$\begin{aligned} |f(x)| &\leq C'' \exp(\tau'|x|^{\varrho(|x|)}) + m_r C''' \exp(\tau'|x|^{\varrho(|x|)}) + 2 \\ &\leq (C'' + CC''' + 2) \exp(\tau|x|^{\varrho(|x|)}). \end{aligned}$$

Here $C > 0$ was chosen independent of r large enough such that

$$m_r \exp(-(\tau - \tau')r^{\varrho(r)}) \leq C,$$

which is possible since $m_r = \lfloor 2e\tau\rho_1 r^{\rho_1} \rfloor$ grows polynomially in r . Since this is true for every $|x| \geq r_0$, this implies $f \in A_{\varrho, \tau}$. □

The following proposition is a direct consequence of Theorem 3.7.

Proposition 3.9. *Let ϱ be a normalized proximate order function and $f(x) = \sum_{\ell=0}^{+\infty} x^\ell a_\ell \in \mathcal{SM}_L(\mathbb{R}^{n+1})$. Then the following four statements are equivalent:*

- (1) $f \in A_{\varrho}$;
- (2) $\limsup_{r \rightarrow +\infty} \frac{\sup_{|x| \leq r} \ln(|f(x)|)}{r^{\varrho(r)}} < \infty$;
- (3) $\limsup_{\ell \rightarrow \infty} |a_\ell|^{\frac{1}{\ell}} \varphi(\ell) < \infty$;
- (4) $\inf\{\beta > 0 : f \in A_{\varrho, \beta}\} < \infty$.

In § 4, we will need some estimates on the norms of monomials, which will be provided in the following lemma.

Lemma 3.10. *Let ϱ be a normalized proximate order function. Then for every $0 < \sigma' < \sigma$, there exists a constant $C(\sigma, \sigma')$ such that*

$$\|x^\ell\|_{\varrho, \sigma} \leq C(\sigma, \sigma') \frac{G_\ell}{\sigma'^{\ell/\rho}}, \quad \ell \in \mathbb{N}_0. \tag{3.15}$$

Proof. First of all, by Lemma 2.14, there exists some $t_0 \geq 0$, such that

$$\frac{\varphi(t)}{\varphi(t')} \leq \frac{e^{\sigma \frac{t}{t'}}}{(e\sigma'\rho)^{1/\rho}}, \quad t, t' \geq t_0. \tag{3.16}$$

Let now $r_0 := \varphi(t_0)$. Since $\lim_{t \rightarrow \infty} \varphi(t) = \infty$, there exists another constant $t_1 \geq 0$ such that

$$\varphi(\ell) \geq r_0(e\rho\sigma')^{\frac{1}{\rho}}, \quad \ell \geq t_1. \tag{3.17}$$

Let now $\ell \geq t_1$. Then for every $|x| \geq r_0$ we set $t := |x|^{e(|x|)}$ and get

$$\frac{|x|^\ell}{G_\ell} \exp(-\sigma|x|^{e(|x|)}) = \frac{|x|^\ell (e\rho)^{\frac{\ell}{\rho}}}{\varphi(\ell)^\ell} \exp(-\sigma|x|^{e(|x|)}) = \frac{\varphi(t)^\ell (e\rho)^{\frac{\ell}{\rho}}}{\varphi(\ell)^\ell} e^{-\sigma t}.$$

Since $t = |x|^{e(|x|)} \geq r_0^{e(r_0)} = t_0$, we are allowed to use (3.16) and estimate this equation by

$$\frac{|x|^\ell}{G_\ell} \exp(-\sigma|x|^{e(|x|)}) \leq \frac{e^{\ell\sigma \frac{t}{t'}} (e\rho)^{\frac{\ell}{\rho}}}{(e\sigma'\rho)^{\frac{\ell}{\rho}}} e^{-\sigma t} = \frac{1}{\sigma'^{\frac{\ell}{\rho}}}, \quad |x| \geq r_0, \ell \geq t_1. \tag{3.18}$$

For $|x| \leq r_0$ on the other hand, we use (3.17) to get

$$\frac{|x|^\ell}{G_\ell} \exp(-\sigma|x|^{e(|x|)}) \leq \frac{r_0^\ell}{G_\ell} = \frac{r_0^\ell (e\rho)^{\frac{\ell}{\rho}}}{\varphi(\ell)^\ell} \leq \frac{1}{\sigma'^{\frac{\ell}{\rho}}}, \quad |x| \leq r_0, \ell \geq t_1. \tag{3.19}$$

Combining now (3.18) and (3.19) gives the estimate

$$\|x^\ell\|_{\varrho, \sigma} = \sup_{x \in \mathbb{R}^{n+1}} |x|^\ell \exp(-\sigma|x|^{e(|x|)}) \leq \frac{G_\ell}{\sigma'^{\frac{\ell}{\rho}}}, \quad \ell \geq t_1.$$

Finally, choosing the constant

$$C(\sigma, \sigma') := \max \left\{ \max_{0 \leq \ell \leq t_1} \frac{\|x^\ell\|_{\varrho, \sigma} \sigma'^{\frac{\ell}{\rho}}}{G_\ell}, 1 \right\},$$

gives the stated estimate (3.15) for every $\ell \in \mathbb{N}_0$. □

Lemma 3.11. *There exists a constant k depending only on ρ for which the following statement holds: For any $\sigma > 0$, we can take $C(\sigma)$ such that for any $f \in A_{\hat{e},\sigma}$, and any $\ell \in \mathbb{N}_0$, the inequality*

$$\frac{1}{\ell!} \|\partial_{x_0}^\ell f(x)\|_{\hat{e},k\sigma} \leq C(\sigma) \|f\|_{\hat{e},\sigma} \frac{(2k\sigma)^{\ell/\rho}}{G_{\hat{e},\ell}} \tag{3.20}$$

holds.

Proof. Let $x \in \mathbb{R}^{n+1}$ and $j \in \mathbb{S}$, $u, v \in \mathbb{R}$ such that $x = u + jv$. Then the Cauchy estimates for the restriction $f|_{\mathbb{C}_j}$ give for any $s > 0$

$$\begin{aligned} |\partial_{x_0}^\ell f(x)| &\leq \frac{\ell!}{s^\ell} \max_{|\xi|=s, \xi \in \mathbb{C}_j} |f(x + \xi)| \\ &\leq \frac{\ell!}{s^\ell} \|f\|_{\hat{e},\sigma} \max_{|\xi|=s, \xi \in \mathbb{C}_j} \exp(\sigma|x + \xi|^{\ell(|x+\xi|)}) \\ &\leq \frac{\ell!}{s^\ell} \|f\|_{\hat{e},\sigma} \exp(\sigma(|x| + s)^{\ell(|x|+s)}) \\ &\leq \frac{\ell!}{s^\ell} \|f\|_{\hat{e},\sigma} e^{\sigma C_\varepsilon} \exp(2^{\rho+\varepsilon} \sigma (r^{\ell(r)} + s^{\ell(s)})), \end{aligned}$$

where in the last inequality we used Lemma 2.9 (i). Choosing now the special value $s = \varphi(\frac{\ell}{2^{\rho+\varepsilon}\sigma\rho})$, i.e. $s^{\ell(s)} = \frac{\ell}{2^{\rho+\varepsilon}\sigma\rho}$, we can further estimate

$$|\partial_{x_0}^\ell f(x)| \leq \frac{\ell!}{\varphi(\frac{\ell}{2^{\rho+\varepsilon}\sigma\rho})^\ell} \|f\|_{\hat{e},\sigma} e^{\sigma C_\varepsilon} \exp(2^{\rho+\varepsilon} \sigma r^{\ell(r)} + \frac{\ell}{\rho}), \quad \ell \in \mathbb{N}_0.$$

By Lemma 2.14 (ii), there exists some $N \in \mathbb{N}$ such that

$$\frac{\varphi(\ell)^\ell}{\varphi(\frac{\ell}{2^{\rho+\varepsilon}\sigma\rho})^\ell} \leq 2^{\frac{\ell}{\rho}} (2^{\rho+\varepsilon}\sigma\rho)^{\frac{\ell}{\rho}}, \quad \ell \geq N.$$

Hence we can choose a larger constant C_1 such that this inequality holds for all $\ell \in \mathbb{N}_0$, i.e.

$$\frac{\varphi(\ell)^\ell}{\varphi(\frac{\ell}{2^{\rho+\varepsilon}\sigma\rho})^\ell} \leq C_1 (2^{\rho+1+\varepsilon}\sigma\rho)^{\frac{\ell}{\rho}}, \quad \ell \in \mathbb{N}_0.$$

This leads to the estimate

$$\begin{aligned} |\partial_{x_0}^\ell f(x)| &\leq C_1 \frac{\ell!}{\varphi(\ell)^\ell} ((2^{\rho+1+\varepsilon}e\sigma\rho)^{\frac{1}{\rho}})^\ell \|f\|_{\hat{e},\sigma} e^{\sigma C_\varepsilon} \exp(2^{\rho+\varepsilon} \sigma r^{\ell(r)}) \\ &= C_1 e^{\sigma C_\varepsilon} \frac{\ell!}{G_\ell} (2^{\rho+1+\varepsilon}\sigma)^{\frac{\ell}{\rho}} \|f\|_{\hat{e},\sigma} \exp(2^{\rho+\varepsilon} \sigma r^{\ell(r)}), \quad \ell \in \mathbb{N}_0. \end{aligned}$$

This is exactly the stated inequality (3.20) for $\ell \geq N$. □

In view of the above stated properties, we can now prove the following crucial results.

Proposition 3.12. *For an entire left slice monogenic function $f(x)$ belonging to $A_{\varrho, \sigma+0}$ for $\sigma \geq 0$, its Taylor expansion $\sum_{\ell=0}^{\infty} x^\ell a_\ell$ converges to $f(x)$ in the space $A_{\varrho, \sigma+0}$. In particular, the set of Fueter polynomials is dense in $A_{\varrho, +0}$ and also dense in A_ϱ .*

Proof. For the former statement, it suffices to show that

$$\sum_{\ell=0}^{\infty} \|x^\ell a_\ell\|_{\varrho, \sigma+\epsilon}$$

is finite for any $\epsilon > 0$. By Lemma 3.10, there exists a positive constant C_0 such that for any $\ell \geq 0$ we have

$$\|x^\ell\|_{\varrho, \sigma+\epsilon} \leq C_0(\sigma + \epsilon/2)^{-\ell/\rho} G_{\hat{\varrho}, \ell}.$$

On the other hand, by Remark 3.8, there exists a positive constant C_1 such that for any $\ell \geq 0$ we have

$$|a_\ell| G_{\hat{\varrho}, \ell} \leq C_1(\sigma + \epsilon/4)^{\ell/\rho}.$$

Therefore, we have

$$\sum_{\ell=0}^{\infty} \|x^\ell a_\ell\|_{\varrho, \sigma+\epsilon} \leq C_0 C_1 \sum_{\ell=0}^{\infty} \left(\frac{\sigma + \epsilon/4}{\sigma + \epsilon/2}\right)^{\ell/\rho} < \infty.$$

For the latter statement in the case $f \in A_{\varrho, +0}$, it follows from the former one with $\sigma = 0$ that

$$\lim_{\ell \rightarrow +\infty} \sum_{q \leq \ell} x^q a_q = f(x)$$

in the space $A_{\varrho, +0}$. In the case $f \in A_\varrho$, there exists $\sigma > 0$ such that $f \in A_{\varrho, \sigma+0}$. Then the same convergence holds in the space $A_{\varrho, \sigma+0}$ and therefore also in the space A_ϱ . □

Lemma 3.13. *Let ϱ be a proximate order function, $\sigma, \tau > 0$, $f \in A_{\varrho, \sigma}$, $g \in A_{\varrho, \tau}$. Then $f \star_L g \in A_{\varrho, \sigma+\tau}$ and*

$$\|f \star_L g\|_{\varrho, \sigma+\tau} \leq 2^{\frac{n+4}{2}} \|f\|_{\varrho, \sigma} \|g\|_{\varrho, \tau},$$

where n is the number of imaginary units in the Clifford algebra \mathbb{R}_n .

Proof. Due to Definition 2.1, the function $f \in \mathcal{SM}(\mathbb{R}^{n+1})$ does admit the decomposition

$$f(u + jv) = f_0(u, v) + jf_1(u, v), \quad u, v \in \mathbb{R}, j \in \mathbb{S},$$

where for any arbitrary $j \in \mathbb{S}$, the functions f_0, f_1 are given by

$$f_0(u, v) = \frac{f(u + jv) + f(u - jv)}{2}$$

and

$$f_1(u, v) = j \frac{f(u - jv) - f(u + jv)}{2},$$

for any $u, v \in \mathbb{R}$. Hence, since $f \in A_{\varrho, \sigma}$, the functions f_0, f_1 admit the estimates

$$\begin{aligned} |f_0(u, v)| &\leq \frac{|f(u + jv)| + |f(u - jv)|}{2} \leq \|f\|_{\varrho, \sigma} \exp\left(\sigma(u^2 + v^2)^{\frac{1}{2}} \varrho((u^2 + v^2)^{\frac{1}{2}})\right), \\ |f_1(u, v)| &\leq \frac{|f(u - jv)| + |f(u + jv)|}{2} \leq \|f\|_{\varrho, \sigma} \exp\left(\sigma(u^2 + v^2)^{\frac{1}{2}} \varrho((u^2 + v^2)^{\frac{1}{2}})\right). \end{aligned}$$

The same obviously holds true for the decomposition $g(u + jv) = g_0(u, v) + jg_1(u, v)$, i.e.

$$\begin{aligned} |g_0(u, v)| &\leq \|g\|_{\varrho, \tau} \exp\left(\tau(u^2 + v^2)^{\frac{1}{2}} \varrho((u^2 + v^2)^{\frac{1}{2}})\right), \\ |g_1(u, v)| &\leq \|g\|_{\varrho, \tau} \exp\left(\tau(u^2 + v^2)^{\frac{1}{2}} \varrho((u^2 + v^2)^{\frac{1}{2}})\right). \end{aligned}$$

Altogether, for every $x \in \mathbb{R}^{n+1}$ with the decomposition $x = u + jv$, we then get for the star product

$$\begin{aligned} |(f \star g)(x)| &= \left| f_0(u, v)g_0(u, v) - f_1(u, v)g_1(u, v) + j(f_0(u, v)g_1(u, v) + f_1(u, v)g_0(u, v)) \right| \\ &\leq 4 \cdot 2^{\frac{n}{2}} \|f\|_{\varrho, \sigma} \|g\|_{\varrho, \tau} \exp\left(\sigma(u^2 + v^2)^{\frac{1}{2}} \varrho((u^2 + v^2)^{\frac{1}{2}})\right) \times \\ &\quad \times \exp\left(\tau(u^2 + v^2)^{\frac{1}{2}} \varrho((u^2 + v^2)^{\frac{1}{2}})\right) \\ &= 2^{\frac{n+4}{2}} \|f\|_{\varrho, \sigma} \|g\|_{\varrho, \tau} \exp\left((\sigma + \tau)|x|^{\varrho(|x|)}\right), \end{aligned}$$

where $x \in \mathbb{R}^{n+1}$ and the factor $2^{\frac{n}{2}}$ in the second lines comes from the fact that for two Clifford numbers $a, b \in \mathbb{R}_n$ the modulus of the product estimates as $|ab| \leq 2^{\frac{n}{2}}|a||b|$. □

The results of this section will be used to characterize continuous homomorphisms in terms of differential operators in the sense that will be specified in the next section.

4. Differential operators, representations of continuous homomorphisms

In this section, we study continuous homomorphism from A_{ϱ_1} and A_{ϱ_2} and those from $A_{\varrho_1,+0}$ and $A_{\varrho_2,+0}$ where $\varrho_i(r)$ ($i = 1, 2$) are two proximate orders functions for positive orders $\rho_i = \lim_{r \rightarrow \infty} \varrho_i(r) > 0$, satisfying

$$r^{\varrho_1(r)} = O(r^{\varrho_2(r)}), \quad \text{as } r \rightarrow \infty. \tag{4.1}$$

Definition 4.1. Let ϱ_i ($i = 1, 2$) be two proximate orders functions for orders $\rho_i > 0$ satisfying (4.1). We take normalization $\hat{\varrho}_1$ of ϱ_1 as in Remark 2.11 and we define $G_{\hat{\varrho}_1,q}$ as in (2.11). We denote by $\mathbf{D}_{\varrho_1 \rightarrow \varrho_2}$ the set of all formal right linear differential operator P of the form

$$P = \sum_{\ell \in \mathbb{N}_0} u_\ell \star_L \partial_{x_0}^\ell$$

where the sequence $(u_\ell(x))_{\ell \in \mathbb{N}_0} \subset A_{\varrho_2}$ satisfies

$$\forall \lambda > 0, \exists \sigma > 0, \exists C > 0, \forall \ell \in \mathbb{N}_0, \|u_\ell\|_{\varrho_2,\sigma} \leq C \frac{G_{\hat{\varrho}_1,\ell}}{\ell!} \lambda^\ell. \tag{4.2}$$

We denote by $\mathbf{D}_{\varrho_1 \rightarrow \varrho_2,0}$ the set of all formal right linear differential operator P of the form

$$P = \sum_{\ell \in \mathbb{N}_0} u_\ell \star_L \partial_{x_0}^\ell$$

where the sequence $(u_\ell(x))_{\ell \in \mathbb{N}_0} \subset A_{\varrho_2}$ satisfies

$$\forall \sigma > 0, \exists \lambda > 0, \exists C > 0, \forall \ell \in \mathbb{N}_0, \|u_\ell\|_{\varrho_2,\sigma} \leq C \frac{G_{\hat{\varrho}_1,\ell}}{\ell!} \lambda^\ell. \tag{4.3}$$

Note that in the latter case, each u_ℓ belongs to $A_{\varrho_2,+0}$.

For the following remark see also [10, Remark 4.4].

Remark 4.2. By adding $\ln(c)/\ln(r)$ for a constant $c > 0$ (with a suitable modification near $r = 0$) to a proximate order function $\varrho_2(r)$ with order ρ_2 , we get a new proximate order function $\tilde{\varrho}_2(r)$ for the same order ρ_2 satisfying

$$\tilde{\varrho}_2(r) = \varrho_2(r) + \ln(c)/\ln(r)$$

for $r > 1$, that is,

$$r^{\tilde{\varrho}_2(r)} = cr^{\varrho_2(r)},$$

eventually. Then $\|\cdot\|_{\tilde{\varrho}_2,\sigma}$ and $\|\cdot\|_{\varrho_2,c\sigma}$ become equivalent norms for $\sigma > 0$, and the spaces $A_{\tilde{\varrho}_2}$ and $A_{\tilde{\varrho}_2,+0}$ are homeomorphic to A_{ϱ_2} and $A_{\varrho_2,+0}$ respectively. By taking c

sufficiently large, we can take $\tilde{\varrho}_2$ as

$$\varrho_1(r) \leq \tilde{\varrho}_2(r), \quad \text{for } r \geq r_0$$

for a suitable r_0 , and we can choose normalizations $\hat{\varrho}_1$ of ϱ_1 and $\hat{\varrho}_2$ of $\tilde{\varrho}_2$ as

(i) $\hat{\varrho}_1(r) \leq \hat{\varrho}_2(r)$ for $r \geq 0$.

Since a proximate order function and its normalization define equivalent norms, we have

(ii) $\|\cdot\|_{\hat{\varrho}_2, \sigma}$ and $\|\cdot\|_{\varrho_2, c\sigma}$ are equivalent for any $c > 0$,

(iii) $\mathbf{D}_{\varrho_1 \rightarrow \varrho_2} = \mathbf{D}_{\hat{\varrho}_1 \rightarrow \hat{\varrho}_2}$, $\mathbf{D}_{\varrho_1 \rightarrow \varrho_2, 0} = \mathbf{D}_{\hat{\varrho}_1 \rightarrow \hat{\varrho}_2, 0}$.

Note further that Theorems 4.3 and 4.4 below are not affected by the replacement of ϱ_1 and ϱ_2 by $\hat{\varrho}_1$ and $\hat{\varrho}_2$.

Theorem 4.3. *Let ϱ_1, ϱ_2 be two proximate orders functions satisfying (4.1). Then all continuous linear operators $T : A_{\varrho_1} \rightarrow A_{\varrho_2}$ are characterized by operators of the form*

$$T := \sum_{\ell=0}^{\infty} u_{\ell} \star_L \frac{\partial^{\ell}}{\partial x_0^{\ell}}, \tag{4.4}$$

with coefficients $(u_{\ell})_{\ell \in \mathbb{N}} \subseteq \mathcal{SM}(\mathbb{R}^{n+1})$ satisfying

$$\forall \varepsilon > 0, \exists \sigma, C > 0 : \|u_{\ell}\|_{\varrho_2, \sigma} \leq C \frac{G_{\varrho_1, \ell}}{\ell!} \varepsilon^{\ell}, \tag{4.5}$$

i.e. $\mathbf{D}_{\varrho_1 \rightarrow \varrho_2}$ is the set of all continuous operators from A_{ϱ_1} to A_{ϱ_2} .

Proof. Due to Remark 4.2, we may assume that $\varrho_1(r) \leq \varrho_2(r)$ for every $r \geq 0$, which implies $A_{\varrho_1, \sigma} \subseteq A_{\varrho_2, \sigma}$ for every $\sigma > 0$ and

$$\|f\|_{\varrho_2, \sigma} \leq \|f\|_{\varrho_1, \sigma}, \quad f \in A_{\varrho_1, \sigma}.$$

Step 1. For the first implication, let T be an operator of the form (4.4) with coefficients $(u_{\ell})_{\ell \in \mathbb{N}}$ satisfying (4.5). Let $\varepsilon > 0$, then by (4.5) there exists $\sigma, C > 0$ such that

$$\|u_{\ell}\|_{A_{\varrho_2, \sigma}} \leq C \frac{G_{\varrho_1, \ell}}{\ell!} \varepsilon^{\ell}, \quad \ell \in \mathbb{N}.$$

Moreover, for $f \in A_{\varrho_1}$ one has $f \in A_{\varrho_1, \tau}$ for some $\tau > 0$ and we can estimate

$$\begin{aligned} \|Tf\|_{\varrho_2, \sigma + 2\rho_1 + 1\tau} &\leq \sum_{\ell=0}^{\infty} \|u_{\ell} \star_L \partial_{x_0}^{\ell} f\|_{\varrho_2, \sigma + 2\rho_1 + 1\tau} \\ &\leq 4 \sum_{\ell=0}^{\infty} \|u_{\ell}\|_{\varrho_2, \sigma} \|\partial_{x_0}^{\ell} f\|_{\varrho_2, 2\rho_1 + 1\tau} \end{aligned}$$

$$\begin{aligned}
 &\leq 4C \sum_{\ell=0}^{\infty} \frac{G_{\varrho_1, \ell}}{\ell!} \varepsilon^\ell \|\partial_{x_0}^\ell f\|_{\varrho_1, 2^{\rho_1+1}\tau} \\
 &\leq 4C \sum_{\ell=0}^{\infty} \frac{G_{\varrho_1, \ell}}{\ell!} \varepsilon^\ell C(\tau) \frac{\ell!(2^{\rho_1+2}\tau)^{\frac{\ell}{\rho_1}}}{G_{\varrho_1, \ell}} \|f\|_{\varrho_1, \tau} \\
 &= 4CC(\tau) \|f\|_{\varrho_1, \tau} \sum_{\ell=0}^{\infty} \left(\varepsilon(2^{\rho_1+2}\tau)^{\frac{1}{\rho_1}}\right)^\ell.
 \end{aligned}$$

Choosing $\varepsilon < (2^{\rho_1+2}\tau)^{-\frac{1}{\rho_1}}$, the right hand side is finite and we showed that T is bounded as an operator from $A_{\varrho_1, \tau}$ to $A_{\varrho_2, \sigma+2^{\rho_1+1}\tau}$. Since f was arbitrary, this also proves the continuity of T as an operator from A_{ϱ_1} to A_{ϱ_2} according to the Definition 3.3.

Step 2. For the second implication, let $T : A_{\varrho_1} \rightarrow A_{\varrho_2}$ be an everywhere defined continuous operator. Then, thanks to the theory of locally convex spaces, there exists for every $\tau > 0$ some $\tau' > 0$, such that

$$\|Tf\|_{\varrho_2, \tau'} \leq C_\tau \|f\|_{\varrho_1, \tau}, \quad f \in A_{\varrho_1, \tau}, \tag{4.6}$$

(see [34, Chapter 4, Part 1, 5, Corollary 1]). Let us now define the functions

$$u_\ell(x) := \frac{1}{\ell!} \sum_{k=0}^{\ell} \binom{\ell}{k} T(x^k) \star_L (-x)^{\ell-k}. \tag{4.7}$$

For any $\varepsilon > 0$ choose $\tau_0 := (\frac{2}{\varepsilon})^{\rho_1}$ and any $\tau > \tau_0$. Then the functions u_ℓ admit the estimate

$$\begin{aligned}
 \|u_\ell\|_{\varrho_2, \tau'+\tau} &\leq \frac{1}{\ell!} \sum_{k=0}^{\ell} \binom{\ell}{k} \|T(x^k) \star_L (-x)^{\ell-k}\|_{\varrho_2, 2^{\rho_1+1}(\tau'+\tau)} \\
 &\leq \frac{2^{\frac{\ell+4}{2}}}{\ell!} \sum_{k=0}^{\ell} \binom{\ell}{k} \|T(x^k)\|_{\varrho_2, \tau'} \|x^{\ell-k}\|_{\varrho_2, \tau} \\
 &\leq \frac{C(\tau)2^{\frac{\ell+4}{2}}}{\ell!} \sum_{k=0}^{\ell} \binom{\ell}{k} \|x^k\|_{\varrho_1, \tau} \|x^{\ell-k}\|_{\varrho_1, \tau} \\
 &\leq \frac{C(\tau)C(\tau, \tau_0)2^{\frac{\ell+4}{2}}}{\ell! \tau_0^{\frac{\ell}{\rho_1}}} \sum_{k=0}^{\ell} \binom{\ell}{k} G_k G_{\ell-k} \\
 &\leq \frac{C(\tau)C(\tau, \tau_1)2^{\frac{\ell+4}{2}} G_\ell}{n! \tau_0^{\frac{\ell}{\rho_1}}} 2^\ell = \frac{C(\tau)C(\tau, \tau_1)2^{\frac{\ell+4}{2}} G_\ell}{\ell!} \varepsilon^\ell.
 \end{aligned}$$

This estimate shows that the functions u_ℓ satisfy the assumptions (4.5). Next consider a function $f(x) = \sum_{m=0}^M x^m a_m$ be a polynomial. Then

$$\begin{aligned} \sum_{\ell=0}^{\infty} u_\ell(x) \star_L \partial_{x_0}^\ell f(x) &= \sum_{\ell=0}^{\infty} \sum_{m=0}^{\infty} u_\ell(x) \star_L \partial_{x_0}^\ell x^m a_m \\ &= \sum_{\ell=0}^{\infty} \sum_{m=\ell}^{\infty} \frac{1}{\ell!} \sum_{k=0}^{\ell} \binom{\ell}{k} \frac{m!}{(m-\ell)!} T(x^k) \star_L (-x)^{\ell-k} \star_L x^{m-\ell} a_m \\ &= \sum_{\ell=0}^{\infty} \sum_{m=\ell}^{\infty} \sum_{k=0}^{\ell} (-1)^{\ell-k} \binom{n}{k} \binom{m}{\ell} T(x^k) \star_L x^{m-k} a_m \\ &= \sum_{m=0}^{\infty} \sum_{k=0}^m \sum_{\ell=0}^{m-k} (-1)^\ell \binom{m-k}{\ell} \binom{m}{k} T(x^k) \star_L x^{m-k} a_m \\ &= \sum_{m=0}^{\infty} \sum_{k=0}^m \delta_{m-k,0} \binom{m}{k} T(x^k) \star_L x^{m-k} a_m \\ &= \sum_{m=0}^{\infty} T(x^m) a_m = T \left(\sum_{m=0}^{\infty} x^m a_m \right) = Pf(x). \end{aligned} \tag{4.8}$$

Since the polynomials are dense by Lemma 3.12 and the operator T is continuous, the same result holds for any function $f \in A_{\varrho_1}$. \square

Theorem 4.4. *Let ϱ_1, ϱ_2 be two proximate order functions satisfying (4.1). Then all continuous linear operators $T : A_{\varrho_1,+0} \rightarrow A_{\varrho_2,+0}$ are characterized by operators of the form*

$$T := \sum_{\ell=0}^{\infty} u_\ell \star_L \frac{\partial^\ell}{\partial x_0^\ell}, \tag{4.9}$$

with coefficients $(u_\ell)_{\ell \in \mathbb{N}} \subseteq \mathcal{SM}(\mathbb{R}^{n+1})$ satisfying

$$\forall \sigma > 0, \exists \varepsilon, C > 0 : \|u_\ell\|_{\varrho_2,\sigma} \leq C \frac{G_{\varrho_1,\ell}}{\ell!} \varepsilon^\ell, \tag{4.10}$$

i.e. $\mathbf{D}_{\varrho_1 \rightarrow \varrho_2,0}$ is the set of all continuous operators from $A_{\varrho_1,+0}$ to $A_{\varrho_2,+0}$.

Proof. Due to Remark 4.2, we may assume that $\varrho_1(r) \leq \varrho_2(r)$ for every $r \geq 0$, which implies $A_{\varrho_1,\sigma} \subseteq A_{\varrho_2,\sigma}$ for every $\sigma > 0$ and

$$\|f\|_{\varrho_2,\sigma} \leq \|f\|_{\varrho_1,\sigma}, \quad f \in A_{\varrho_1,\sigma}.$$

We proceed in two steps.

Step 1. For the first implication let T be an operator of the form (4.9) with coefficients $(u_\ell)_{\ell \in \mathbb{N}}$ satisfying (4.10). Let $\sigma > 0$. Then with the coefficients $\varepsilon, C > 0$ from (4.10) we choose $0 < \tau < \frac{1}{2\rho_1 + 2\varepsilon\rho_1}$. Then we obtain the estimate

$$\begin{aligned} \sum_{\ell \in \mathbb{N}_0} \|u_\ell \star_L \partial_{x_0}^\ell f\|_{\varrho_{2, \sigma + 2\rho_1 + 1}\tau} &\leq 2^{\frac{n+4}{2}} \sum_{\ell \in \mathbb{N}_0} \|u_\ell\|_{\varrho_{2, \sigma}} \|\partial_{x_0}^\ell f\|_{\varrho_{2, 2\rho_1 + 1}\tau} \\ &\leq 2^{\frac{n+4}{2}} C(\eta, \sigma, \tau) \sum_{\ell \in \mathbb{N}_0} \|u_\ell\|_{\varrho_{2, \sigma}} \|\partial_{x_0}^\ell f\|_{\varrho_{1, 2\rho_1 + 1}\tau} \\ &\leq C'(\eta, \sigma, \tau) \sum_{\ell \in \mathbb{N}_0} \frac{G_{\varrho_1, \ell}}{\ell!} \varepsilon^\ell \|f\|_{\varrho_{1, \tau}} \frac{\ell!}{G_{\rho_1, \ell}} (2^{\rho_1 + 2}\tau)^{\frac{\ell}{\rho_1}} \\ &= C'(\eta, \sigma, \tau) \|f\|_{\varrho_{1, \tau}} \sum_{\ell \in \mathbb{N}_0} \varepsilon^\ell (2^{\rho_1 + 2}\tau)^{\frac{\ell}{\rho_1}}, \end{aligned}$$

for $f \in A_{\varrho_1, \tau}$. Note that the last sum is finite due to the choice of τ . This implies that $P : A_{\varrho_1, +0} \rightarrow A_{\varrho_2, +0}$ is a continuous operator.

Step 2. For the second implication let $P : A_{\varrho_1, +0} \rightarrow A_{\varrho_2, +0}$ be a continuous operator. Then for every $\varepsilon > 0$ there exist $\tau > 0$ and $C(\varepsilon) > 0$, such that $P : A_{\varrho_1, \tau} \rightarrow A_{\varrho_2, \varepsilon}$ is continuous with

$$\|Pf\|_{\varrho_2, \varepsilon} \leq C(\varepsilon) \|f\|_{\varrho_1, \tau}, \quad f \in A_{\varrho_1, \tau}, \tag{4.11}$$

(see [34, Chapter 4, Part 1, 5, Corollary 1]). Defining the functions u_ℓ as in (4.7), they admit the estimate

$$\begin{aligned} \|u_\ell\|_{\varrho_2, \sigma} &\leq 2^{\frac{n+4}{2}} \sum_{k \leq \ell} \frac{\|x^{\ell-k}\|_{\varrho_2, \sigma/2} \|P(x^k)\|_{\varrho_2, \sigma/2}}{(\ell-k)!k!} \\ &\leq \sum_{k \leq \ell} \frac{\|x^{\ell-k}\|_{\varrho_1, \sigma/2} C(\tau_1) \|x^k\|_{\varrho_1, \tau_1}}{(\ell-k)!k!} \\ &\leq \sum_{k \leq \ell} \frac{C(\tau_0, \sigma'/2) \tau_0^{-(\ell-k)/\rho_1} G_{\varrho_1, \ell-k} C(\tau_1) C(\tau_0, \tau_1) \tau_0^{-k/\rho_1} G_{\varrho_1, k}}{(\ell-k)!k!} \tag{4.12} \\ &\leq C(\tau_1) C(\tau_0, \sigma/2) C(\tau_0, \tau_1) \sum_{k \leq \ell} \binom{\ell}{k} \frac{G_{\varrho_1, \ell}}{\ell!} \tau_0^{-\ell/\rho_1} \\ &= C(\tau_1) C(\tau_0, \sigma/2) C(\tau_0, \tau_1) \frac{G_{\varrho_1, \ell}}{\ell!} 2^\ell \tau_0^{-\ell/\rho_1}. \end{aligned}$$

Here we used Lemma 3.13 at the first inequality, (4.11) at the second with $\varepsilon = \sigma/2$, Lemma 3.10 twice with $0 < \tau_0 < \sigma/2$ and $0 < \tau_0 < \tau_1$ at the third and Lemma 2.13 at the fourth. Therefore, by defining

$$\lambda := 2\tau_0^{-1/\rho_1},$$

we have

$$\|u_\ell\|_{\mathcal{E}^{2,\sigma}} \leq C' \frac{G_{\mathcal{E}^{1,\ell}}}{\ell!} \lambda^\ell$$

for any ℓ . Since $\sigma > 0$ can be chosen arbitrarily, the functions u_ℓ satisfy all the conditions in (4.10). Moreover, the operator P admits the representation

$$P = \sum_{\ell=0}^{\infty} u_\ell \star_L \partial_{x_0}^\ell$$

in the same way as in (4.8) for polynomials. Since the polynomials are dense due to Lemma 3.12 and the operator T is continuous, the same result holds for any function $f \in A_{\mathcal{E}^{1,+0}}$. Thus $P \in \mathbf{D}_{\mathcal{E}^1 \rightarrow \mathcal{E}^{2,0}}$. \square

Acknowledgements. S. Pinton is supported by MUR grant ‘Dipartimento di Eccellenza 2023-2027’. P. Schlosser was funded by the Austrian Science Fund (FWF) under Grant No. J 4685-N and by the European Union – NextGenerationEU.

Competing interests. The authors declare none.

References

- (1) Y. Aharonov, D. Albert and L. Vaidman, How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100, *Phys. Rev. Lett.* **60** (1988), 1351–1354.
- (2) Y. Aharonov, J. Behrndt, F. Colombo and P. Schlosser, Schrödinger evolution of superoscillations with δ - and δ' -potentials, *Quantum Stud. Math. Found.* **7**(3) (2020), 293–305.
- (3) Y. Aharonov, F. Colombo, I. Sabadini, D. C. Struppa and J. Tollaksen, On the Cauchy problem for the Schrödinger equation with superoscillatory initial data, *J. Math. Pures Appl.* **99** (2013), 165–173.
- (4) Y. Aharonov, F. Colombo, I. Sabadini, D. C. Struppa and J. Tollaksen, Superoscillating sequences as solutions of generalized Schrodinger equations, *J. Math. Pures Appl.* **103** (2015), 522–534.
- (5) Y. Aharonov, F. Colombo, I. Sabadini, D. C. Struppa and J. Tollaksen, The mathematics of superoscillations, *Mem. Amer. Math. Soc.* **247**(1174) (2017), v+107.
- (6) Y. Aharonov, F. Colombo, I. Sabadini, D. C. Struppa and J. Tollaksen, Evolution of superoscillations in the Klein-Gordon field, *Milan J. Math.* **88** (2020), 171–189.
- (7) Y. Aharonov and D. Rohrlich, *Quantum Paradoxes: Quantum Theory for the Perplexed* (Wiley-VCH Verlag, Weinheim, 2005).
- (8) D. Alpay, F. Colombo, S. Pinton, I. Sabadini and D. C. Struppa, Infinite-order differential operators acting on entire hyperholomorphic functions, *J. Geom. Anal.* **31**(10) (2021), 9768–9799.
- (9) T. Aoki, F. Colombo, I. Sabadini and D. C. Struppa, Continuity theorems for a class of convolution operators and applications to superoscillations, *Ann. Mat. Pura Appl.* **197**(5) (2018), 1533–1545.

- (10) T. Aoki, R. Ishimura and Y. Okada, A differential operator representation of continuous homomorphisms between the spaces of entire functions of given proximate order, *Complex Anal. Oper. Theory* **14**(8) (2020), Paper No. 75, 22.
- (11) T. Aoki, R. Ishimura, Y. Okada, D. C. Struppa and S. Uchida, Characterization of continuous endomorphisms of the space of entire functions of a given order, *Complex Var. Elliptic Equ.* **66**(9) (2021), 1439–1450.
- (12) T. Aoki, F. Colombo, I. Sabadini and D. C. Struppa, Continuity of some operators arising in the theory of superoscillations, *Quantum Stud. Math. Found.* **5** (2018), 463–476.
- (13) J. Behrndt, F. Colombo and P. Schlosser, Evolution of Aharonov–Berry superoscillations in Dirac δ -potential, *Quantum Stud. Math. Found.* **6** (2019), 279–293.
- (14) J. Behrndt, F. Colombo, P. Schlosser and D. C. Struppa, Integral representation of superoscillations via Borel measures and their convergence, *Trans. Amer. Math. Soc.* **376**(9) (2023), 6315–6340. doi:10.1090/tran/8983
- (15) C. Berenstein and R. Gay, *Complex Analysis and Special Topics in Harmonic Analysis* (Springer, New York, 1995).
- (16) F. Brackx, R. Delanghe and F. Sommen, *Clifford Analysis*, Research Notes in Mathematics, Vol. 76 (Pitman (Advanced Publishing Program), Boston, MA, 1982), x+308 pp.
- (17) F. Colombo, I. Sabadini and D. C. Struppa, *Entire Slice Regular Functions*, SpringerBriefs in Mathematics (Springer, Cham, 2016), v+118 pp.
- (18) F. Colombo, I. Sabadini, D. C. Struppa and A. Yger, Superoscillating sequences and supershifts for families of generalized functions, *Complex Anal. Oper. Theory* **16**(3) (2022), Paper No. 34, 37.
- (19) F. Colombo, I. Sabadini, D. C. Struppa and A. Yger, Superoscillating sequences and hyperfunctions, *Publ. Res. Inst. Math. Sci.* **55** (2019), 665–688.
- (20) F. Colombo, I. Sabadini and D. C. Struppa, Slice monogenic functions, *Israel J. Math.* **171** (2009), 385–403.
- (21) F. Colombo and J. Gantner, *Quaternionic Closed Operators, Fractional Powers and Fractional Diffusion Processes*, Operator Theory: Advances and Applications, Vol. 274 (Birkhäuser/Springer, Cham, 2019), viii+322 pp.
- (22) F. Colombo, J. Gantner and D. P. Kimsey, *Spectral Theory on the S-Spectrum for Quaternionic Operators*, Volume 270 of Operator Theory: Advances and Applications (Birkhäuser/Springer, Cham, 2018).
- (23) F. Colombo, I. Sabadini and D. C. Struppa, *Noncommutative Functional Calculus. Theory and Applications of Slice Hyperholomorphic Functions*, Volume 289 of Progress in Mathematics (Birkhäuser/Springer Basel AG, Basel, 2011).
- (24) F. Colombo, I. Sabadini and D. C. Struppa, *Michele Sce's Works in Hypercomplex Analysis – A Translation with Commentaries* (Birkhäuser/Springer, Cham, 2020), vi+122 pp.
- (25) F. Colombo, R. Lavicka, I. Sabadini and V. Soucek, The Radon transform between monogenic and generalized slice monogenic functions, *Math. Ann.* **363** (2015), no. 3–4 733–752.
- (26) F. Colombo, O. J. Gonzalez-Cervantes and I. Sabadini, A nonconstant coefficients differential operator associated to slice monogenic functions, *Trans. Amer. Math. Soc.* **365**(1) (2013), 303–318.
- (27) F. Colombo, R. S. Krausshar, S. Pinton and I. Sabadini, Entire monogenic functions of given proximate order and continuous homomorphisms, *Mediterr. J. Math.* **21**(44) (2024), 28. doi:10.1007/s00009-023-02585-x
- (28) D. Constales, R. De Almeida and R. S. Krausshar, On the growth type of entire monogenic functions, *Arch. Math.* **88** (2007), 153–163.

- (29) D. Constaes, R. De Almeida and R. S. Krausshar, On the relation between the growth and the Taylor coefficients of entire solutions to the higher-dimensional Cauchy–Riemann system in \mathbb{R}^{n+1} , *J. Math. Anal. Appl.* **327** (2007), 763–775.
- (30) R. De Almeida and R. S. Krausshar, On the asymptotic growth of monogenic functions, *Zeitschrift Anal. Anw.* **24**(4) (2005), 791–813.
- (31) R. De Almeida and R. S. Krausshar, Generalized growth orders for polymonogenic functions and related inequalities, *Complex Anal. Operator Theory* **10** (2016), 233–250.
- (32) A. De Martino, K. Diki and A. Guzman Adan, On the connection between the Fueter–Sce–Qian theorem and the generalized CK-extension, *Results Math.* **78**(2) (2023), Paper No. 55, 35.
- (33) P. Lelong and L. Gruman, *Entire Functions of Several Complex Variables*, Vol. 282 (Grung. Math. Wiss., Berlin, Heidelberg, New York, Springer, 1986).
- (34) A. Grothendieck, *Topological Vector Spaces*, Notes on Mathematics and its Applications (Gordon and Breach Science Publishers, New York, London, Paris, 1973).
- (35) K. Gürlebeck, K. Habetha and W. Sprössig, *Application of Holomorphic Functions in Two and Higher Dimensions* (Birkhäuser/Springer, Cham, 2016), xv+390 pp.
- (36) S. Kumar and K. Bala, Generalized order of entire monogenic functions of slow growth, *J. Nonlinear Sci. Appl.* **5** (2012), 418–425.
- (37) P. Schlosser, Time evolution of superoscillations for the Schrödinger equation on $\mathbb{R}\setminus 0$, *Quantum Stud. Math. Found.* **9**(3) (2022), 343–366.
- (38) G. D. Srivastava and S. Kumar, On the generalized order and generalized type of entire monogenic functions, *Demonstr. Math* **46**(4) (2013), 663–677.
- (39) G. Valiron, *Lectures on the General Theory of Integral Functions* (Privat, Toulouse, 1923).