

New theoretical and empirical work is urgently needed so that threats can be tracked and mitigated.

**Keywords:** disasters; climate change; developmental psychopathology; child and adolescent mental health

### EPP0184

#### Childhood mental health and juvenile delinquency: A within-family comparison

S. Kim<sup>1\*</sup> and D.H. Kim<sup>2</sup>

<sup>1</sup>Psychiatry, Ajou university school of medicine, Suwon, Korea, Republic of and <sup>2</sup>Economics, Myongji University, Seoul, Korea, Republic of

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.620

**Introduction:** Child mental health is associated with prospective delinquent outcomes. However, this association might be confounded by genetic and other shared factors

**Objectives:** We aimed to examine the association between the behavioral symptoms of Attention-Deficit/Hyperactivity Disorder, conduct disorder, depression, and oppositional defiant disorder in childhood (age 4-12) and the prospective delinquent outcomes as measured by lifetime illicit drug use, criminal activities, and victimization prior to age 18, using the nationally representative U.S. survey that allowed us to compare siblings in the same mother.

**Methods:** Aged-adjusted subscales of ADHD, conduct disorder, depression, and ODD were obtained from the mother-reported survey responses. Within-family analyses were performed to control for family-specific unobserved factors as well as child-specific observed factors.

**Results:** Antisocial scores in childhood were strongly associated with lifetime arrest, probation, and incarceration as well as lifetime illicit drug use in adolescence. ADHD scores are associated with lifetime victimization in physical attack and rape, but not with criminal activities or illicit drug use.

**Conclusions:** Conduct disorder consistently increases lifetime illicit drug use and criminal activities independently of genetic factors and gender. ADHD is not associated with lifetime illicit drug use or criminal activities, but is associated with lifetime victimization. No significant gender differences are found although anxiety/depression symptoms are often positively associated with delinquent outcomes only among females.

**Keywords:** childhood; mental health; ADHD; delinquency

### EPP0185

#### Behaviour in childhood is associated with distinct patterns of partnering in adulthood

F. Vergunst<sup>1\*</sup>, Y. Zheng<sup>2</sup>, P. Domond<sup>1</sup>, F. Vitaro<sup>3</sup>, R. Tremblay<sup>3</sup>, D. Nagin<sup>4</sup>, J. Park<sup>5</sup> and S. Cote<sup>1</sup>

<sup>1</sup>Public Health, University of Montreal, Montréal, Canada;

<sup>2</sup>Psychology, University of Alberta, Edmonton, Canada;

<sup>3</sup>Psychoeducation, University of Montreal, Montreal, Canada;

<sup>4</sup>Statistics, Carnegie Mellon University, Pittsburgh, United States of America and <sup>5</sup>Health, Statistics Canada, Ottawa, Canada

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.621

**Introduction:** Childhood behavioral problems are highly prevalent in school-aged children and are associated with poor long-term outcomes. Yet little is known about their association with patterns of partnering in adulthood.

**Objectives:** To (1) describe patterns of partnering from age 18-35 years in a large population-based sample, and (2) examine the association between childhood behavioural problems and adult partnering patterns.

**Methods:** Behavioural ratings were prospectively obtained from teachers when children (n=2960) were aged 10-12 years – for inattention, hyperactivity, aggression-opposition, anxiety and pro-sociality – and linked to their tax return records from age 18-35 years. We used group-based trajectory modelling to identify distinct trajectories of partnering (married or cohabitating) and multinomial regression models to examine the association between childhood behaviour and trajectory group membership.

**Results:** Five distinct trajectories of partnering were identified: early-partnered (n=420, 14.4%), mid-partnered (n=620, 21.3%), late-partnered (n=570, 19.2%), early-separated (n=460, 15.5%), and delayed-or-unpartnered (n=890, 30.0%). After adjustment for sex and family background, children rated as being anxious or inattentive were more likely to remain unpartnered from age 18 to 35 years, while those rated as aggressive-oppositional or inattentive were more likely to separate and return to unpartnered status. Prosocial behaviours were consistently associated with earlier and more sustained partnership. Participants in the early-separated and delayed-or-unpartnered trajectories were also more likely to have left high school without a diploma and to have lower earnings.

**Conclusions:** Childhood behavioural problems were associated with increased likelihood of being unpartnered and of partnership dissolution, which has implications for the psychological health and wellbeing of individuals and their families.

**Keywords:** ADHD; Conduct disorder; Disruptive behaviours

### EPP0188

#### Honouring COVID-19 restrictions: A qualitative study of the virtual ASD diagnostic pathway in a UK NHS CAMHS service.

C. Ecob\*, S. Olety, R. Lancaster and A. Harris

Heathy Young Minds, Pennine Care NHS Foundation Trust, Stalybridge, United Kingdom

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.622

**Introduction:** The Multi-Agency Autism Team (MAAT) diagnose Autism Spectrum Disorder (ASD) in children and young people using a multi-stage assessment process. In March 2020, the UK went into lockdown due to the COVID-19 pandemic, affecting the MAAT's ability to continue their typical diagnostic pathway.

**Objectives:** This qualitative study aimed to assess the effectiveness and feasibility of a virtual ASD diagnostic pathway.

**Methods:** From March – September; one hundred detailed developmental history assessments were conducted over the telephone, fifteen socially-distanced BOSA (Brief Observation of Symptoms of Autism) assessments were piloted, twenty-five multi-disciplinary formulation meetings were held over a video platform, and sixty diagnosis feedback consultations were conducted via telephone or video call. Structured interviews were conducted with clinicians and service-users.