

FITTING CLASSES OF CC-GROUPS

by MARTYN R. DIXON

(Received 10th February 1987)

1. Introduction

The theory of Fitting classes is, by now, a well established part of the theory of finite soluble groups. In contrast, Fitting classes have received rather scant attention in infinite groups, although some recent work of Beidleman and Karbe [2] and Beidleman, Karbe and Tomkinson [3] suggest that one can obtain results in this direction. The paper [2], cited above, in fact generalizes earlier work of Tomkinson [9] to the class of locally soluble FC-groups. The present paper is concerned with the theory of Fitting classes in a class of groups somewhat similar to the class of FC-groups, namely the class of CC-groups, introduced by Polovickii in [6]. A group G is a CC-group if $G/C_G(x^G)$ is a Černikov group for all $x \in G$ where, as in the rest of this paper, we use the standard group theoretic notation of [7]. Recently, Alcázar and Otal [1] have shown how to generalize results of B. H. Neumann [5] to the class of CC-groups. The main purpose of the present note is to illustrate further how one can handle CC-groups, in an analogous manner to FC-groups, by using techniques similar to those used in [1] and [4].

Throughout this paper \mathfrak{Y} will denote the class of locally soluble CC-groups. By lemma 1 of [1] each \mathfrak{Y} -group therefore has a local system of normal subgroups each of which is Černikov-by-free abelian of finite rank. The subgroups forming the local system here are the normal closures of finite subsets and hence are centre-by-Černikov, since the class of CC-groups is clearly closed under taking subgroups and homomorphic images.

By a *Fitting class* of \mathfrak{Y} -groups we shall mean a subclass \mathfrak{F} of \mathfrak{Y} with the following properties:

- (i) If $G \in \mathfrak{F}$ and $H \text{ ser } G$ then $H \in \mathfrak{F}$.
- (ii) If $G = \langle H_\lambda \mid \lambda \in \Lambda \rangle \in \mathfrak{Y}$ with $H_\lambda \in \mathfrak{F}$ and $H_\lambda \text{ ser } G$ then $G \in \mathfrak{F}$ also.

(Here Λ is an index set and $H \text{ ser } G$ means H is a serial subgroup of G , our definition of serial being precisely that given in Robinson [7].)

A subgroup V of $G \in \mathfrak{Y}$ will be called an \mathfrak{F} -injector of G if $V \cap S$ is a maximal \mathfrak{F} -subgroup of S whenever $S \text{ ser } G$. The set of maximal \mathfrak{F} -subgroups of G will be denoted by $\text{Max}_{\mathfrak{F}} G$ and the set of \mathfrak{F} -injectors of G will be denoted by $\text{Inj}_{\mathfrak{F}} G$. Our main result, proven in Section 3 of this paper, may then be stated as follows:

Theorem. *If \mathfrak{F} is a Fitting class of \mathfrak{Y} -groups and if $G \in \mathfrak{Y}$ then $\text{Inj}_{\mathfrak{F}} G \neq \emptyset$. Also the elements of $\text{Inj}_{\mathfrak{F}} G$ form a unique local conjugacy class.*

(We recall that subgroups H, K of a group G are *locally conjugate* if there is a locally inner automorphism of G mapping H to K). Thus our theorem generalizes Theorems 3.2 and 3.3 of [9].

The layout of the paper is as follows. In Section 2 the characteristic of a Fitting class is defined and some recent results from [3] quoted. We then obtain a special case of the Theorem, by showing that certain centre-by-Černikov groups possess a unique conjugacy class of \mathfrak{F} -injectors for each Fitting class \mathfrak{F} . In Section 3 this allows us to invoke the inverse limit arguments of [4] to obtain the main result, the proof being somewhat similar to the corresponding one in [9].

2. The characteristic of a Fitting class

If \mathfrak{F} is a Fitting class of \mathfrak{Y} -groups then as in [3] we define the *finite characteristic* of \mathfrak{F} to be the set of primes p such that $C_p \in \mathfrak{F}$, where C_p is the cyclic group of order p . The finite characteristic will be denoted by $C_f(\mathfrak{F})$. The *infinite characteristic*, $C_i(\mathfrak{F})$, of \mathfrak{F} is defined to be $\{\infty\}$ if \mathfrak{F} contains the infinite cyclic group; otherwise $C_i(\mathfrak{F}) = \emptyset$, the empty set. The *characteristic* of \mathfrak{F} is defined to be $C_f(\mathfrak{F}) \cup C_i(\mathfrak{F})$ and is denoted by $C(\mathfrak{F})$. Hence $C(\mathfrak{F})$ is a subset of $\mathbb{P} \cup \{\infty\}$, where \mathbb{P} denotes the set of all primes.

If $G \in \mathfrak{Y}$ and $x \in G$ then clearly x^G is a soluble Černikov-by-cyclic group. The following is therefore a straightforward consequence of [3, lemma 2.1].

Lemma 2.1. *Let $G \in \mathfrak{Y}$*

- (i) *If G contains an element of prime order p then there are subgroups $K \triangleleft H \text{ sn } G$ such that $H/K \cong C_p$.*
- (ii) *If G contains an element of infinite order then there are subgroups $K \triangleleft H \text{ sn } G$ such that $H/K \cong C_\infty$.*

A standard argument (see [3, Theorem 2.2, 3, Theorem 2.4]) now shows:

Lemma 2.2. *Let \mathfrak{F} be a Fitting class of \mathfrak{Y} -groups*

- (i) *If there is a group $G \in \mathfrak{F}$ which contains a p -element then $C_p \in \mathfrak{F}$.*
- (ii) *If there is a group $G \in \mathfrak{F}$ which contains an element of infinite order then $C_\infty \in \mathfrak{F}$.*
- (iii) *If $C_\infty \in \mathfrak{F}$ then $C(\mathfrak{F}) = \mathbb{P} \cup \{\infty\}$.*
- (iv) *Either $C(\mathfrak{F}) = \pi \subseteq \mathbb{P}$ and every \mathfrak{F} -group is a π -group or $C(\mathfrak{F}) = \mathbb{P} \cup \{\infty\}$.*
- (v) *If $C(\mathfrak{F}) = \pi \subseteq \mathbb{P}$ then \mathfrak{F} contains all locally nilpotent π -groups in \mathfrak{Y} .*
- (vi) *If $C(\mathfrak{F}) = \mathbb{P} \cup \{\infty\}$ then \mathfrak{F} contains all locally nilpotent \mathfrak{Y} -groups.*

(Of course the proof of the latter two facts depend on the fact that all subgroups of a locally nilpotent group are serial.)

These results concerning characteristic will only be required in Černikov-by-(free abelian of finite rank) \mathfrak{Y} -groups. Such groups are of course \mathfrak{S}_1 -groups (in the sense of [3]). The full results are included here for the sake of completeness.

To obtain results in the Černikov-by-(free abelian of finite rank) case we require the following preliminary result on serial subgroups:

Lemma 2.3. *Let G be a centre-by-Černikov group. Then every serial subgroup of G is ascendant.*

Proof. Let Z denote the centre of G and let $S \text{ ser } G$. We show that $SZ/Z \text{ ser } G/Z$ from which it follows that $S \triangleleft SZ \text{ asc } G$. In fact, if $(\Lambda_\sigma, V_\sigma: \sigma \in \Sigma)$ is a series of G containing S then there is a series of G containing SZ consisting of the subgroups $(\bigcap_{\tau > \sigma} V_\tau Z, \Lambda_\sigma Z, V_\sigma Z: \sigma \in \Sigma)$. This follows from elementary set theory. The fact that $\Lambda_\sigma Z \triangleleft \bigcap_{\tau > \sigma} V_\tau Z$ follows because $[\Lambda_\sigma Z, \bigcap_{\tau > \sigma} V_\tau Z] \leq \bigcap_{\tau > \sigma} [\Lambda_\sigma, V_\tau] \leq \bigcap_{\tau > \sigma} V_\tau = \Lambda_\sigma$. \square

If \mathfrak{F} is a Fitting class of \mathfrak{Y} -groups and $G \in \mathfrak{Y}$ we shall let $G_{\mathfrak{F}}$ denote the \mathfrak{F} -radical of G . Then $G_{\mathfrak{F}}$ is the largest normal \mathfrak{F} -subgroup of G . The following result is important for our purposes:

Theorem 2.4. *Let G be an extension of a central finitely generated abelian group Z by a soluble Černikov group and let \mathfrak{F} be a Fitting class of \mathfrak{Y} -groups. Then*

- (a) *G has a unique conjugacy class of \mathfrak{F} -injectors.*
- (b) *If $C(\mathfrak{F}) = \mathbb{P} \cup \{\infty\}$ then G has only finitely many \mathfrak{F} -injectors.*
- (c) *If $C(\mathfrak{F}) = \pi \subseteq \mathbb{P}$ then the cardinality of $\text{Inj}_{\mathfrak{F}} G$ is the same as the cardinality of $|T: N_T(V)|$ where V is a specific \mathfrak{F} -injector of G and T is the subgroup of G consisting of elements of finite order.*

Proof. (a) Since G is centre-by-Černikov it is a CC-group and G' is Černikov, by [7, Theorem 4.21, Corollary 2]. Hence the set T of elements of finite order in G forms a normal subgroup of G . Since $T \cap Z$ is finite and TZ/Z is Černikov it follows that T is Černikov. If $C(\mathfrak{F}) = \mathbb{P} \cup \{\infty\}$ and if H/Z is the divisible part of G/Z then $H \in \mathfrak{F}$ by 2.2(vi). Hence $G/G_{\mathfrak{F}}$ is finite and the result now follows by a special case of [3, Theorem 4.4]. Part (b) clearly follows from this since the \mathfrak{F} -injectors must contain $G_{\mathfrak{F}}$. If, on the other hand, $C(\mathfrak{F}) = \pi \subseteq \mathbb{P}$, then T has an \mathfrak{F} -injector V , by [3, Theorem 4.4]. In fact $V \in \text{Inj}_{\mathfrak{F}} G$. For, if $A \text{ ser } G$ then by 2.3, $A \text{ asc } G$. If $V \cap A \leq X \leq A$ with $X \in \mathfrak{F}$ then X is a π -group by 2.2(i). Hence $X \leq T$ so $X \leq A \cap T$. Since $A \cap T \text{ asc } T$ and $V \in \text{Inj}_{\mathfrak{F}} T$ it follows that $V \cap (A \cap T) \in \text{Max}_{\mathfrak{F}}(A \cap T)$. Hence $V \cap A = X$ and then it follows that G has \mathfrak{F} -injectors. Since the \mathfrak{F} -injectors must all lie in T , conjugacy also follows, by [3, Theorem 4.4]. The remainder of part (c) is now evident, since any two \mathfrak{F} -injectors of G are actually \mathfrak{F} -injectors of T . \square

3. Proof of the theorem

It is convenient to split the proof of the main theorem in two pieces, namely a proof of existence and a proof of local conjugacy of the injectors. The main technical result required in our proofs is [8, Theorem 2.1] which requires that we make certain sets into compact topological spaces. In general this would not be possible, but for Černikov groups such a topology always exists (see [4]) and it is this fact which is important here.

Theorem 3.1. *Let \mathfrak{F} be a Fitting class of \mathfrak{Y} -groups. Then the \mathfrak{Y} -group G possesses \mathfrak{F} -injectors.*

Proof. The group G has a local system, \mathcal{L} , consisting of soluble normal subgroups,

each of which is Černikov-by-(free abelian of finite rank) and finitely generated as a G -operator group. Hence each $N \in \mathcal{L}$ is an extension of a central finitely generated abelian group by a soluble Černikov group. It follows that each $N \in \mathcal{L}$ has a unique conjugacy class of \mathfrak{F} -injectors. If $N \in \mathcal{L}$ let $A(N)$ denote the set of \mathfrak{F} -injectors of N . Notice that if $C(\mathfrak{F}) = \mathbb{P} \cup \{\infty\}$ then each $A(N)$ is a finite set and the proof follows as in [9, Theorem 3.2]. So we may assume $C(\mathfrak{F}) = \pi \subseteq \mathbb{P}$, by 2.2(iv). Let $N \in \mathcal{L}$ and $V \in \text{Inj}_{\mathfrak{F}} N$. By 2.4(iii), $A(N)$ is in 1-1 correspondence with the set of cosets of $N_N(V)$ in N via the correspondence:

$$V^t \leftrightarrow tN_N(V) \quad (t \in N).$$

The sets $A(N)$ may be partially ordered by defining $A(N) \leq A(M)$ if and only if $N \leq M$ whenever $N, M \in \mathcal{L}$. Clearly, if $W \in A(M)$ and $A(N) \leq A(M)$ then $W \cap N \in A(N)$ so we may define a projection $\pi_{MN}: A(M) \rightarrow A(N)$ by

$$(W)\pi_{MN} = W \cap N.$$

If $N \in \mathcal{L}$ the set $A(N)$ can be made into a compact topological T_1 -space as follows: The group $\bar{N} = N/Z(N)$ is Černikov so has the coset topology, defined in [4], and \bar{N} is then a compact, topological T_1 -space whose closed sets are in fact finite unions of right cosets of subgroups, by [4, Lemma 2.1]. Let $N^* = N_N(V)$ denote the set of cosets of $N_N(V)$ in N . Since $Z(N) \leq N_N(V)$ there is a natural map $\Theta_N: \bar{N} \rightarrow N^*$, defined by $nZ(N) \mapsto nN_N(V)$, for $n \in N$. The set N^* is given the quotient topology induced by Θ_N and hence N^* is also a compact topological T_1 -space. Since $A(N)$ is in 1-1 correspondence with N^* there is an induced topology on $A(N)$ and in this way $A(N)$ is a compact topological T_1 -space.

The topology induced on $A(N)$ is independent of the choice of $V \in \text{Inj}_{\mathfrak{F}} N$. This follows using an argument similar to that given in [4, Theorem 3.8]. Also if $N \leq M$ with $M, N \in \mathcal{L}$ and $W \in A(M)$ then there is a natural map

$$\beta_{MN}: \frac{M}{N_M(W)} \rightarrow \frac{N}{N_N(W \cap N)}.$$

For, by a Frattini argument, $M = N \cdot N_M(W \cap N)$ so if $x \in M$ then $x = yt$ for some $y \in N$ and $t \in N_M(W \cap N)$ and one can define β_{MN} by:

$$(xN_M(W))\beta_{MN} = yN_N(W \cap N).$$

It is easy to see that β_{MN} is well defined and that it is a closed, continuous map. It follows that $\pi_{MN}: A(M) \rightarrow A(N)$ is also closed and continuous since the topology induced on both $A(M)$ and $A(N)$ does not depend on the choice of injectors. Hence $\varprojlim A(N)$ is non-empty by [8, Theorem 2.1]. If $(V_N) \in \varprojlim A(N)$ then for $N \leq M$, we have $V_M \cap N = V_N$ so that $V = \bigcup_{N \in \mathcal{L}} V_N$ is a subgroup of G . Moreover $V \in \mathfrak{F}$ since $V \cap N = V_N \triangleleft V$ so V is generated by normal \mathfrak{F} -subgroups. Finally if $S \text{ ser } G$ then $S \cap N \text{ ser } N$ for each $N \in \mathcal{L}$. Suppose $V \cap S \leq X \leq S$ with $X \in \mathfrak{F}$. Then $V \cap S \cap N = X \cap N$ since $V \cap N \in \text{Inj}_{\mathfrak{F}} N$ and it is now clear that $V \cap S \in \text{Max}_{\mathfrak{F}} S$. Hence V is the required \mathfrak{F} -injector. \square

The local conjugacy of the \mathfrak{F} -injectors is easy to establish using the argument given by Alcázar and Otal [1, Theorem 1] and the result established in 2.4; we therefore simply state:

Theorem 3.2. *Let \mathfrak{F} be a Fitting class of \mathfrak{V} -groups. Then the \mathfrak{F} -injectors of $G \in \mathfrak{V}$ are locally conjugate in G .*

In conclusion, we remark that one can now extend many of the standard results from the theory of Fitting classes in finite soluble groups to the class \mathfrak{V} .

Acknowledgement. The author would like to thank the referee for simplifying the proof of 2.3 and for correcting other parts of Section 2.

REFERENCES

1. J. ALCÁZAR and J. OTAL, Sylow subgroups of groups with Černikov conjugacy classes, *J. Algebra*, **110** (1987), 507–513.
2. J. C. BEIDLEMAN and M. J. KARBE, Injectors of locally soluble FC-groups, *Monatsh. Math.* **103** (1987), 7–13.
3. J. C. BEIDLEMAN, M. J. KARBE and M. J. TOMKINSON, Fitting classes of \mathfrak{S}_1 -groups, I, to appear.
4. M. R. DIXON, Some topological properties of residually Černikov groups, *Glasgow Math. J.* **23** (1982), 65–82.
5. B. H. NEUMANN, Isomorphisms of Sylow subgroups of infinite groups, *Math. Scand.* **6** (1958), 299–307.
6. YA. D. POLOVICKII, Groups with extremal classes of conjugate elements, *Sibirsk. Mat. Zh.* **5** (1964), 891–895.
7. D. J. S. ROBINSON, *Finiteness Conditions and Generalized Soluble Groups* (Springer-Verlag, Berlin, 1972).
8. I. STEWART, Conjugacy theorems for a class of locally finite Lie algebras, *Compositio Math.* **30** (1975), 181–210.
9. M. J. TOMKINSON, \mathfrak{F} -injectors of locally soluble FC-groups, *Glasgow Math. J.* **10** (1969), 130–136.

DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF ALABAMA
 P.O. Box 1416
 TUSCALOOSA, AL 35487
 USA