
INSTRUCTION TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, *Journal of Helminthology*,
London School of Hygiene and Tropical Medicine,
Winches Farm Field Station,
395 Hatfield Road,
St Albans, Herts AL4 0XQ,
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the *Journal*. The maximum size an illustration will be printed is 12.0 × 20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the *Journal* (maximum size 12.0 × 20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Composition, secretion, and fate of the glands in the miracidium and sporocyst of <i>Fasciola hepatica</i> L. (Research Note) GERALD R. BUZZELL	79-84
Mucosal lesions in the rat associated with the attachment of <i>Moniliformis</i> (Acanthocephala). JEAN MARTIN, D. W. T. CROMPTON and ARUNA SINGHVI	85-90
Metacercaria of <i>Galactosomum ussuriense</i> Oshmarin, 1963 (Trematoda: Heterophyidae) from brackish water fishes of Bheemunipatnam (India). Z. REKHARANI and R. MADHAVI	91-94
Trapping of large numbers of larvae in the livers of <i>Toxocara canis</i> -reinfected mice. KAZUO SUGANE and TOMOO OSHIMA	95-99
<i>Taenia crassiceps</i> : ultrastructural observations on the oncosphere and associated structures. M. W. K. CHEW	101-113
Immunological studies of amphistomiasis in sheep and goats II. Viability of amphistome metacercariae after irradiation. (Research Note) MD. HAFEEZ and B. V. RAO	115-116
Population distribution of <i>Heterakis pusilla</i> in <i>Gallus gallus</i> L. from India. SANDEEP K. MALHOTRA	117-126
Immunocytochemical and ultrastructural studies on <i>Dipetalonema viteae</i> (Filarioidea). A. PRÜSSE, S. VOLLMER and H. J. DIESFELD	127-142
Redescription of <i>Thometrema magnifica</i> (Szidat, 1954) Gibson & Bray, 1979 (Trematoda: Hemiuroidea), with comments on the validity of <i>T. portoalegrensis</i> Amato, 1968. A. KOHN, O. M. FRÖES, B. M. M. FERNANDES and B. ABRAMSON	143-148
Morphological and histochemical observations on the oocyst and oviducal transport of oocytes in <i>Paramphistomum cervi</i> (Zeder, 1790) (Digenea: Trematoda). B. C. GUPTA, V. R. PARSHAD and S. S. GURAYA	149-153
Immunological regulation of <i>Angiostrongylus cantonensis</i> infections in rats: modulation of population density and enhanced parasite growth following one or two superimposed infections. W. K. YONG and COLIN DOBSON	155-165
Larval <i>Anisakis simplex</i> (Rudolphi, 1809, det. Krabbe, 1878) and larval <i>Hysterothylacium</i> sp. (Nematoda: Ascaridoidea) in euphausiids (Crustacea: Malacostraca) in the North-East Atlantic and northern North Sea. JOHN W. SMITH	167-177
SUPPLEMENTARY REVIEW ARTICLE	
Ascaridoid nematodes of amphibians and reptiles: <i>Typhlophorus</i> , <i>Hartwichia</i> and <i>Trispiculascaris</i> . J. F. A. SPRENT	179-189