XVII Canadian Congress of Neurological Sciences

Four Seasons Hotel Toronto, Ontario June 23-26, 1982

- 1. Pre-Congress Courses
- 2. Special Lectures
- 3. Abstracts of the Scientific Program

For Further Information Contact:

Dr. R.P. Humphreys Room 1504 Hospital for Sick Children 555 University Avenue Toronto, Ontario, Canada M5G 1X8



PAEDIATRIC NEUROLOGIST

There is an opportunity for a full time academic paediatric neurologist with advanced specialty training in neurophysiology. The candidate should have documented expertise in paediatric and neonatal electroencephalography and in sensory evoked potential techniques. Appointment will be at Assistant Professor level in the Division of Neurology, Department of Paediatrics, University of Toronto. Interested candidates should send their Curriculum Vitae and two supporting letters by July 1, 1982 to:

William J. Logan, M.D., Neurologist-In-Chief The Hospital for Sick Children, 555 University Avenue, Toronto, Ontario. M5G 1X8

In Parkinson Therapy:

4 parts L-dopa: 1 part benserazide

For the response you expect... without the frequency of peripheral side effects

The 4:1 ratio provides

- Excellent clinical response in the management of Parkinsonian disability. 1,2
- Significantly fewer side effects nausea and vomiting - than a 10:1 ratio Levodopa/Čarbidopa preparation during the first six months of treatment. 1,2
- Patient preference over the 10:1 ratio Levodopa/Carbidopa preparation, with respect to nausea and vomiting.1

References:

- 1) Rinne UK, Mölsä P. Neurology, 1979; 29:1584-1589. 2) Pakkenberg H et al, Acta Neurol. Scand 1976; 53:376-385.

right from the start



Original Research in Medicine and Chemistry Can. 0094 Reg. Trade Mark

See page vii for brief prescribing information.



Information and companionship for Parkinson patients is available from the Parkinson Foundation of Canada. Please write or call: Suite 232, ManuLife Centre, 55 Bloor St. West, Toronto, Ontario M4W 1A6. Telephone: (416) 964-1155.



To help control refractory generalized tonic-clonic seizures without excessive sedation

