
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology
London School of Hygiene and Tropical Medicine
Keppel Street, London WC1E 7HT
England

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5 × 18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 13.5 × 18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and physiology of parasites* (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
MATOSSIAN, R. M., SALT, I. and STEPHAN, E. Variation in serum immunoglobulin levels in acute trichinosis	1—4
FRANSEN, F. Investigations of the unimicrobial infection of <i>Schistosoma intercalatum</i> in snails and the infection of the final host using cercariae of one sex	5—10
McVICAR, A. H. Intestinal helminth parasites of the ray <i>Raja naevus</i> in British waters	11—21
PANITZ, E. An evaluation of ethyl-6-ethoxybenzothiazole-2-carbamate (Sch 18099) for anthelmintic activity in domestic animals	23—30
BEVERIDGE, I. On two new linstowiid cestodes from Australian dasyurid marsupials	31—40
GIBBONS, L. M. Revision of the genera <i>Longistrongylus</i> Le Roux 1931, <i>Kobusinema</i> Ortlepp, 1963 and <i>Bigalakenema</i> Ortlepp, 1963 (Nematoda: Trichostrongylidae)	41—62
MOBEDI, I. and GHADIRIAN, E. <i>Schizorchis arfaei</i> sp. n. (Cestoda: Anoplocephalidae) from the wood-mouse, <i>Apodemus sylvaticus</i> , in caspian area of Iran	63—67
ROWNTREE, S. and JAMES, C. Research Note. Single sex cercariae of <i>Schistosoma mansoni</i> , a comparison of male and female infectivity	69—70
SCAPIN, M. and TENDLER, M. Research Note. Immunoprecipitins in human schistosomiasis detected with adult worm antigens released by 3M KC1	71—72
PRAH, S. K. and JAMES, C. The influence of physical factors on the survival and infectivity of miracidia of <i>Schistosoma mansoni</i> and <i>S. haematobium</i> I. Effect of temperature and ultra-violet light	73—85
HASHIGUCHI, Y. and HIRAI, H. Influence of immunosuppressants on the establishment of <i>Paragonimus miyazakii</i> in albino rats	87—94