

Rumen development process in goats as affected by supplemental feeding v. grazing: age-related anatomic development, functional achievement and microbial colonisation

Jinzheng Jiao^{1,2}, Xiaopeng Li^{1,2}, Karen A. Beauchemin³, Zhiliang Tan^{1*}, Shaoxun Tang¹ and Chuanshe Zhou¹

¹Key Laboratory of Agro-Ecological Processes in Subtropical Region, South-Central Experimental Station of Animal Nutrition and Feed Science, Ministry of Agriculture, Institute of Subtropical Agriculture, The Chinese Academy of Sciences, Changsha, Hunan 410125, People's Republic of China

²Graduate University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

³Lethbridge Research Centre, Agriculture and Agri-Food Canada, Lethbridge, Alberta, Canada T1J 4B1

(Submitted 2 July 2014 – Final revision received 20 October 2014 – Accepted 7 December 2014 – First published online 26 February 2015)

Abstract

The aim of the present study was to describe age-related changes in anatomic, functional and microbial variables during the rumen development process, as affected by the feeding system (supplemental feeding v. grazing), in goats. Goats were slaughtered at seven time points that were selected to reflect the non-rumination (0, 7 and 14 d), transition (28 and 42 d) and rumination (56 and 70 d) phases of rumen development. Total volatile fatty acid (TVFA) concentration ($P=0.002$), liquid-associated bacterial and archaeal copy numbers ($P<0.01$) were greater for supplemental feeding v. grazing, while rumen pH ($P<0.001$), acetate molar proportion ($P=0.003$) and solid-associated microbial copy numbers ($P<0.05$) were less. Rumen papillae length ($P=0.097$) and extracellular ($P=0.093$) and total ($P=0.073$) protease activity potentials in supplemented goats tended to be greater than those in grazing goats. Furthermore, from 0 to 70 d, irrespective of the feeding system, rumen weight, rumen wall thickness, rumen papillae length and area, TVFA concentration, xylanase, carboxymethylcellulase activity potentials, and microbial copy numbers increased ($P<0.01$) with age, while the greatest amylase and protease activity potentials occurred at 28 d. Most anatomic and functional variables evolved progressively from 14 to 42 d, while microbial colonisation was fastest from birth to 28 d. These outcomes suggest that the supplemental feeding system is more effective in promoting rumen development than the grazing system; in addition, for both the feeding systems, microbial colonisation in the rumen is achieved at 1 month, functional achievement at 2 months, and anatomic development after 2 months.

Key words: Feeding systems: Anatomic development: Functional achievement: Microbial colonisation

Development of the rumen is an important physiological challenge for young ruminants. It entails growth and cellular differentiation of the rumen, and results in a major shift in the pattern of nutrients being delivered to the intestines and liver, and thus to the peripheral tissues of the animal⁽¹⁾. Numerous studies have shown that the rumen development process consists of anatomic development (increase in rumen mass and growth of the rumen papillae)^(2,3), functional achievement (fermentation capacity and enzyme activity)^(4,5), as well as microbial colonisation (bacteria, fungi, methanogenic archaea and protozoa)^(6,7). Insufficient rumen development will negatively affect nutrient digestion and absorption, or even lead to diseases such as respiratory disease and diarrhoea⁽¹⁾. In contrast,

complete rumen development facilitates digestion of feed components, thereby providing nutrients for the physiological requirements of the animal.

Previous studies^(8,9) have demonstrated that the rumen development process occurs in three phases: non-rumination phase (0–3 weeks); transition phase (3–8 weeks); rumination phase (from 8 weeks onwards). In young calves, ruminal enzyme activities have been observed as early as 2 d of age⁽⁵⁾. It has also been found that highest xylanase and amylase activities in the rumen occurred at 7 and 10 d of age, respectively, suggesting degradation capacity, even before the availability of substrate⁽⁵⁾. In contrast, Siddons⁽¹⁰⁾ claimed that ruminal amylase activities increased with age, which was

Abbreviations: CMCCase, carboxymethylcellulase; LAM, liquid-associated microbes; SAM, solid-associated microbes; TVFA, total volatile fatty acid; VFA, volatile fatty acid.

* **Corresponding author:** Z. Tan, fax +86 731 4612685, email zltan@isa.ac.cn

confirmed by observations that the ruminal amylolytic bacterial community increased with age⁽¹¹⁾. Furthermore, it has been noted that colonisation of bacteria in the rumen is sequential, and begins in the first week of life^(12,13). Compared with studies focusing on ruminal bacteria, less attention has relatively been given to colonisation mechanisms of archaea, fungi and protozoa, especially by using non-culture-based molecular methods. Collectively, despite the importance of rumen development, there is still little information available on changes in rumen anatomic development and functional achievement, especially the microbial colonisation process, during these three different phases of rumen development.

In intensive farming, supplemental feeding is a preferred method of providing nutrients with emphasis on offering young ruminants solid starter concentrates at a relative early age^(5,13,14). During the past few decades, research on rumen development has been mainly focused on this type of feeding system. Concomitantly, factors affecting rumen development processes in ruminants receiving supplemental feeding have been extensively illustrated⁽¹⁵⁾, with primary attention on diet composition^(16–19). For instance, total volatile fatty acid (TVFA) concentrations were higher for calves fed ground- *v.* coarse-grain starter⁽¹⁶⁾. Moreover, addition of starter concentrates to forage diets positively influenced the anatomic structure of the rumen wall⁽¹⁹⁾.

Feed supplementation for young ruminants is not prevalent in many areas of the world where ruminant production is based primarily on grazing pastures with limited supplemental feed^(20,21). It has been reported that an artificial feeding system with a milk replacer led to lower rumen weight, lower volatile fatty acid (VFA) concentrations and greater pH, when compared with a natural system wherein the offspring remained with its mother⁽²²⁾. However, it is not clear whether the information available on rumen development in supplement-fed ruminants is relevant for grazing ruminants. There is a paucity of knowledge on rumen development in grazing animals, and no studies have been reported in the peer-reviewed literature on differences between these two feeding systems.

Therefore, the present study aimed at (1) describing the age-related sequential dynamic changes in anatomic, functional and microbial variables during time intervals that span the three phases of rumen development, and (2) determining the effect of the feeding system on the rumen development process. In the present study, solid feed (20 d) and weaning (40 d) trials were conducted during the transition phase. The effect of age was tested from birth to 70 d, while the effect of the feeding system was tested after the solid feed was provided (from 20 to 70 d).

Materials and methods

The experiment was approved by the Animal Care Committee, Institute of Subtropical Agriculture, the Chinese Academy of Sciences, Changsha, China.

Animals, diets and management

A total of forty-four 1-d-old male and female goats were separated from their dams. The experiment start for each

goat was staggered to accommodate the differing birth dates. To ensure that the environmental conditions were similar throughout the experiment, all the goats were housed in a well-ventilated room with controlled temperature and humidity. The goats weighed an average of 1.35 (SEM 0.12) kg at birth. Once each birth was noticed, the kid was immediately placed in an individual pen to avoid direct contact with adult animals. The goats were then maintained within an individual pen for the duration of the study. The animals allocated to the grazing procedure (see below) were returned to their individual pens overnight. From birth to 20 d, all the goats consumed only goat milk *ad libitum*. Each goat was offered twice daily a bucket of 1 litre of goat milk (per meal) at 08.00 and 17.00 hours. A total of four goats were slaughtered at each of the following ages: 0, 7 and 14 d. The remaining thirty-two goats were randomly divided into two treatment groups, based on different feeding systems: supplemental feeding and grazing. For the supplemented group, between 20 and 40 d, the goats were gradually weaned off goat milk and fed a diet of forage supplemented with starter concentrate. Feeds were provided individually twice daily at 08.00 and 17.00 hours; at each meal, the animals received a bucket of goat milk (0.5 litres), and a separate bucket containing a mixture of forage (fresh grass; 0.04 kg DM/meal) and starter concentrate (0.12 kg DM/meal). After 40 d, the goats consumed only forage (0.06 kg DM for each of the two meals provided daily at 08.00 and 17.00 hours) supplemented with starter concentrate (0.17 kg DM at each meal) until 70 d.

The starter concentrate (per kg DM) was composed of 74.1 g whey powder, 211 g maize flour, 320 g bean meal, 65 g fishmeal, 220 g fat powder, 51 g milk powder, 8.6 g CaCO₃, 25.3 g CaHPO₄, 5 g NaCl and 20 g premix. The forage fed to the supplemented group was harvested from the same pasture as grazed by the grazing group. Between 20 and 40 d, goats in the grazing group were weaned off goat milk and fed a diet of only forage, which was achieved by grazing on pasture. The goats were offered goat milk (0.5 litres/meal) for 10 min at 08.00 and 17.00 hours, and grazed 8 h daily on pasture. After 40 d, goats in the grazing group received no milk, and just grazed on pasture for 8 h daily until 70 d. In both the groups, four goats were slaughtered at each of the following ages: 28, 42, 56 and 70 d. All goats had free access to water.

Collection procedures and sampling

After slaughtering the goats, live weight and eviscerated hot carcass weight (hide removed) were measured. The rumen was removed, and the content was divided into three parts. The first one-third of the rumen content was filtered through four layers of cheesecloth, and the rumen fluid pH was immediately determined (pH meter model 2000; Beckman Instruments, Inc.). A 4 ml sample of the rumen fluid was collected and stored at –20°C for the analysis of NH₃-N and VFA.

Another one-third of the rumen content was used to isolate solid-associated microbes (SAM) and liquid-associated microbes (LAM), according to the procedures of Yang *et al.*⁽²³⁾.

To avoid sample contamination between samples collected on the same day, the rumen tissue samples were taken using individual sterilised scalpels, and put into individual DNase- and RNase-free centrifuge tubes. Similarly, each rumen content sample was immediately put into an individual container prefilled with CO₂, and the isolation procedure of SAM and LAM was conducted under anaerobic conditions. The isolated SAM and LAM were stored in individual DNase- and RNase-free centrifuge tubes at –80°C until extraction of DNA. Microbial isolation was performed because LAM and SAM differed in composition and function, and could be affected by dietary factors. Many previous studies have demonstrated that SAM account for 70% of all rumen contents, and thus they are considered to be pivotal to feed digestion in the rumen^(23,24).

The final one-third of the rumen content was used to isolate extracellular and intracellular enzymes, as described by Ha *et al.*⁽²⁵⁾. Extracellular enzymes play an essential role in feed digestion. Intracellular enzymes also play a similar role, and finding ways to release these enzymes may also increase ruminal forage fermentation⁽²⁵⁾. The reason for measuring both extracellular and intracellular enzymes was to establish how age and the feeding system affect the relative proportions, as a means of understanding the development of digestive capacity within the rumen. After the rumen was emptied of its contents, it was rinsed repeatedly with PBS (pH 7.4) until clean, and was then drained of excess PBS and reweighed. Overall, four sections of the rumen wall (approximately 1 cm²) were taken according to the procedures suggested by Lesmeister *et al.*⁽²⁶⁾, and fixed in 10% formalin solution for anatomic analysis.

Assessment of anatomic structure

Rumen tissue samples were removed from buffered formalin, trimmed and dehydrated using graded ethanol (50, 70, 80, 90 and 100%) and xylene. Samples were then embedded in paraffin, sectioned and stained with haematoxylin and buffered eosin⁽²⁾. Rumen papillae length and area were measured on the stained sections at 4× magnification by digital planar morphometry, using a fluorescence microscope (Olympus). A minimum of twenty intact and well-oriented papillae were measured for each sample.

Chemical analysis

The NH₃-N concentrations of the rumen fluid were determined by the phenol-hypochlorite method using a UV–visible spectrophotometer (UV-2450; Shimadzu) at 550 nm. VFA (acetate, propionate, butyrate, valerate, isobutyrate and isovalerate) concentrations were analysed by a gas chromatograph (7890A; Agilent). Procedures have been described in detail in our previous work⁽²⁷⁾.

Determination of enzyme activity potentials

The preparation of samples for enzyme activity potential measurements was done as suggested by Ha *et al.*⁽²⁵⁾. Extracellular and intracellular enzyme activity potentials of

carboxymethylcellulase (CMCase), xylanase and amylase in rumen contents were determined by measuring the release of reducing sugars from substrates (carboxymethylcellulose, xylan and starch, respectively)⁽⁵⁾. The reaction times were 30, 15 and 15 min, respectively. One enzyme activity unit (U) was defined as the amount of enzyme required to release 1 μmol of reducing sugars (xylose or glucose equivalents)/min per g of wet rumen content. Extracellular and intracellular protease activity potentials were assayed using azocasein as a substrate, according to the method of Eun & Beauchemin⁽²⁸⁾. The hydrolysis of azocasein released an azo group, which induced a coloration measured by spectrophotometry at 420 nm. For each enzyme, the potential of total enzyme activity is the sum of the extracellular and intracellular enzyme activities.

DNA extraction and real-time quantitative PCR

Genomic DNA of LAM and SAM was extracted from 1 ml of fluid using the QIAamp DNA Stool Mini Kit (Qiagen GmbH), according to the manufacturer's instructions with a slight modification. The fluid was incubated at 95°C for 10 min instead of the original 70°C for 5 min after the addition of ASL buffer, in order to lyse both Gram-positive and Gram-negative microbial cells. The quality and quantity of DNA were measured on the basis of absorbance at 260 and 280 nm using a NanoDrop ND1000 (NanoDrop Technologies, Inc.). Primers used for quantification of bacteria and fungi⁽²⁹⁾, methanogenic archaea⁽³⁰⁾, and protozoa⁽³¹⁾ were from previous studies. The quantitative PCR was performed using procedures detailed in our earlier work⁽²⁷⁾. Briefly, a standard curve was generated for each microbial group, using plasmid DNA containing the exact 16S or 18S ribosomal RNA gene inserts. The quantitative PCR assays were performed on an ABI 7900HT system (Applied Biosystems) with a total volume of 10 μl, using SYBR[®] Premix Ex Taq[™] (Takara). By relating the C_t value to the standard curves, the final copy numbers of targeted bacteria, fungi, methanogenic archaea and protozoa per g of rumen contents were calculated as described by Chen *et al.*⁽³²⁾. The values were converted to log₁₀ for further statistical analysis.

Statistical analysis

The effect of the feeding system (supplemental feeding and grazing) was examined from 28 to 70 d. Data were analysed as a completely randomised design using the MIXED procedure of SAS (SAS Institute, Inc.) with a model that included the fixed effects of feeding system, age and feeding system × age interaction, with individual animal as the experimental unit because each goat was individually fed and measurements were taken from individual goats. The slice option was used when the feeding system × age interaction was significant, to partition and test the effect of the feeding system on age. To test the effects of age on rumen development of goats from 0 to 70 d, the MIXED procedure of SAS (SAS Institute, Inc.) was used, with animal nested within age as the random effect and individual animal as the experimental unit. Orthogonal contrasts were used to test for

linear, quadratic and cubic effects of age. Quartic, quintic and sextic effects were not examined because they could not be interpreted biologically. If there was no feeding system × age interaction from 28 to 70 d, the linear and quadratic effects of age from 0 to 70 d were averaged over the two feeding systems (supplemental feeding and grazing). If the interaction was significant, the effects of age from 0 to 70 d for the supplemented and grazing groups were presented separately. Statistical significance was accepted at $P < 0.05$, and a trend was considered at $P < 0.10$. All presented data are expressed as least square means.

Results

In the present study, goats were offered fixed amounts of goat milk until 40 d (1 litre/meal from 0 to 20 d and 0.5 litres/meal from 20 to 40 d), and typically all of this milk was consumed. Intakes of the fixed amounts of solid feed provided as forage and starter concentrate to the supplemented group were not recorded in the present study. In addition, it was not possible to monitor the intake for the grazing group under the grazing conditions at the farm. Nonetheless, average weight gain was nearly identical between the supplemented and grazing groups (Table 1); therefore, energy and N intakes were probably similar. Furthermore, for the supplemented goats, a parallel trial (data not shown) indicated that during the period when only goat milk was offered, the goats consumed almost all of the goat milk provided. This separate trial also showed that both during and after weaning, forage and concentrate consumption increased with age.

Live weight and carcass weight of goats at different ages as affected by the different feeding systems

There were no interaction effects between feeding system and age on live weight ($P = 0.984$) and carcass weight ($P = 0.918$), and the feeding system did not affect these parameters ($P > 0.10$) (Table 1). In both groups of goats, an increasing cubic trend of age was observed from 0 to 70 d on live weight ($P = 0.051$), while a quadratic increasing effect of age was observed on carcass weight ($P = 0.009$), with the greatest rate of increase occurring between 14 and 28 d.

Anatomic development of rumen parameters of goats at different ages as affected by the different feeding systems

As presented in Table 2, the feeding system did not affect rumen weight ($P = 0.655$), rumen wall thickness ($P = 0.330$) or rumen papilla number ($P = 0.160$), and there were no feeding system × age interactions ($P > 0.10$) for these variables. Irrespective of the feeding system, rumen weight and rumen wall thickness increased cubically with age ($P < 0.05$), rumen papilla number decreased quadratically with age ($P < 0.001$), and decreased rapidly between 14 and 28 d. Rumen papilla length in both the groups increased linearly with age ($P < 0.001$), and its value tended to be greater ($P = 0.097$) for supplemental feeding *v.* grazing.

Table 1. Live weight and carcass weight of the goats at different ages as affected by the different feeding systems (Least square means and standard errors)

Items	System	Age (d)							P*				P for age†			
		0‡	7‡	14‡	28	42	56	70	SEM§	System	Age	System × age	SEM	L	Q	C
Live weight (kg)	S	1.35	1.54	1.97	4.24	5.96	6.54	7.43	0.397	0.838	<0.001	0.984	0.308	<0.001	0.059	0.051
	G				4.25	6.10	6.44	7.61								
Carcass weight (kg)	S	0.53	0.54	1.05	2.09	2.49	2.68	2.86	0.266	0.829	0.021	0.918	0.213	<0.001	0.009	0.817
	G				1.99	2.46	2.47	3.03								

L, linear effect of age; Q, quadratic effect of age; C, cubic effect of age; S, supplemental feeding; G, grazing.

* P value for both feeding systems from 28 to 70 d.

† P for age from 0 to 70 d.

‡ During 0, 7 and 14 d, all goats consumed only milk. After 20 d, goats were randomised into two groups and received solid feed either in S or as G. Four goats were analysed at 0, 7 and 14 d, while eight goats (four goats each for diets S and G) were analysed at 28, 42, 56 and 70 d. The data for 0, 7 and 14 d were merged for S and G because these two groups were fed similarly during this period.

§ For system × age interaction: from 28 to 70 d.

|| For age: from 0 to 70 d.

Table 2. Rumen anatomic parameters of goats at different ages as affected by the different feeding systems (Least square means and standard errors)

Items	System	Age (d)								P*				P for age†		
		0‡	7‡	14‡	28	42	56	70	SEM§	System	Age	System × age	SEM	L	Q	C
Rumen weight (g)	S	6.12	6.50	8.25	40.75	89.31	101.83	118.00	7.583	0.655	<0.001	0.984	5.344	<0.001	0.885	<0.001
	G				40.00	93.07	103.71	122.82								
Rumen wall thickness (mm)	S	0.06	0.07	0.07	0.11	0.16	0.20	0.20	0.015	0.330	<0.001	0.946	0.012	<0.001	0.822	0.013
	G				0.11	0.18	0.21	0.22								
Rumen papillae length (mm)	S	ND	0.20	0.34	0.63	0.74	0.81	0.87	0.064	0.097	0.002	0.849	0.053	<0.001	0.111	0.261
	G				0.54	0.63	0.72	0.86								
Rumen papillae area (mm ²)	S	ND	0.03	0.04	0.13	0.20	0.22	0.21	0.014	<0.001	<0.001	0.016	0.012	<0.001	<0.001	0.427
	G				0.07	0.14	0.15	0.23					0.011	<0.001	0.396	0.065
Papillae number (number of rumen papillae per mm ² area)	S	ND	6.59	5.93	3.67	2.39	2.22	1.62	0.294	0.160	<0.001	0.327	0.256	<0.001	<0.001	0.341
	G				2.73	2.51	1.98	1.48								

L, linear effect of age; Q, quadratic effect of age; C, cubic effect of age; S, supplemental feeding; G, grazing; ND, not detected due to insufficient sample size.

* P value for both feeding systems from 28 to 70 d.

† P for age from 0 to 70 d.

‡ During 0, 7 and 14 d, all goats consumed only milk. After 20 d, goats were randomised into two groups and received solid feed either in S or as G. Four goats were analysed at 0, 7 and 14 d, while eight goats (four goats each for diets S and G) were analysed at 28, 42, 56 and 70 d. The data for 0, 7 and 14 d were merged for S and G because these two groups were fed similarly during this period.

§ For system × age interaction: from 28 to 70 d.

|| For age: from 0 to 70 d.

In contrast, an interaction between feeding system and age ($P=0.016$) for rumen papillae area was observed, with the papillae area being greater for the supplemented group than for the grazing group at 28, 42 and 56 d, but less than that for the grazing group at 70 d. Furthermore, from 0 to 70 d, in the supplemented goats, rumen papillae area increased quadratically with age ($P<0.001$), while in the grazing goats, it tended to increase cubically with age ($P=0.065$).

Rumen fermentation capacity of goats at different ages as affected by the different feeding systems

There was no feeding system × age interaction ($P>0.10$) on rumen pH or $\text{NH}_3\text{-N}$ concentration, and the grazing group had higher pH ($P<0.001$) and lower $\text{NH}_3\text{-N}$ concentrations ($P<0.001$) than those of the supplemented group (Table 3). From 0 to 70 d, in both groups, age had no effect on pH ($P=0.103$), but had a quadratic increasing effect ($P=0.005$) on $\text{NH}_3\text{-N}$ concentration.

No VFA were detected in rumen samples collected on the day of birth. No feeding system × age interactions were observed for TVFA concentration ($P=0.650$) and acetate molar proportion ($P=0.281$). The molar proportion of acetate was higher ($P=0.003$) while TVFA concentration was lower ($P=0.002$) for grazing *v.* supplemental feeding. From 7 to 70 d, in both groups, age had a quadratic effect on TVFA concentration ($P=0.005$), while a cubic effect was observed on acetate molar proportion ($P=0.011$). The feeding system × age interaction tended to be significant ($P=0.088$) for the acetate:propionate ratio. Age tended to have a cubic effect on the acetate:propionate ratio in the supplemented group ($P=0.086$), and had a cubic effect on the acetate:propionate ratio in the grazing group ($P=0.001$).

Interaction effects between feeding system and age ($P<0.05$) were observed on other molar proportions of VFA except acetate because at different ages, there was no consensus on whether the supplemented group exceeded the grazing group. From 7 to 70 d, age had cubic effects on propionate molar proportion in the grazing goats ($P=0.004$), butyrate molar proportion in both groups ($P<0.05$), and isobutyrate ($P<0.001$), valerate ($P=0.012$) and isovalerate ($P<0.001$) molar proportions in the supplemented goats. Furthermore, a quadratic effect of age ($P<0.001$) was observed on the molar proportion of isobutyrate and isovalerate in the grazing goats, and their values fluctuated with age.

Potentials of ruminal enzyme activities of goats at different ages as affected by the different feeding systems

The enzyme activity potentials of rumen contents were too low to be detected on the day of birth, and at 7 d, amylase and protease could not be detected because the sample size was insufficient to conduct the analytical procedures. The feeding system × age interaction was observed for intracellular xylanase ($P=0.022$), extracellular CMCCase ($P=0.001$) and total CMCCase ($P=0.012$) because most of these activities were greater at 28 and 42 d, while less at 56 and 70 d for grazing *v.* supplemental feeding (Table 4). In addition,

Table 3. pH, NH₃-N and volatile fatty acids (VFA) of the rumen fluid of goats at different ages as affected by the different feeding systems (Least square means and standard errors)

Items	System	Age (d)							SEM§	P*			SEM	P for age†		
		0‡	7‡	14‡	28	42	56	70		System	Age	System × age		L	Q	C
pH	S	6.47	6.66	6.37	6.53	6.33	6.23	5.98	0.182	<0.001	0.020	0.822	0.139	0.103	0.169	0.774
	G				7.25	6.88	6.61	6.62								
NH ₃ -N (mg/ml)	S	1.51	4.75	5.75	23.23	20.61	24.09	23.10	2.172	<0.001	0.699	0.690	1.583	<0.001	0.005	0.355
	G				11.49	11.01	10.09	13.70								
Total VFA (mm)	S	N	16.61	17.77	46.26	83.60	68.86	91.12	4.258	0.002	<0.001	0.650	3.599	<0.001	0.005	0.437
	G				35.36	71.98	64.61	76.17								
Acetate (%)¶	S	N	58.93	79.93	67.20	63.40	59.14	54.05	2.508	0.003	0.078	0.281	2.699	0.087	0.033	0.011
	G				68.02	70.35	64.39	64.82								
Propionate (%)	S	N	20.79	10.36	24.62	20.30	17.89	26.94	1.684	0.001	0.138	0.015	2.336	0.014	0.457	0.318
	G				17.26	17.27	19.28	17.19								
Acetate:propionate	S	N	3.09	8.07	3.10	3.16	3.37	2.03	0.326	0.001	0.156	0.088	1.500	0.300	0.622	0.004
	G				3.98	4.08	3.36	3.79								
Butyrate (%)	S	N	9.40	3.16	4.63	10.87	15.74	13.76	1.128	0.157	<0.001	0.018	1.352	<0.001	0.148	<0.001
	G				7.99	8.67	11.55	12.13								
Isobutyrate (%)	S	N	3.48	2.31	1.11	1.75	2.15	1.29	0.154	0.699	0.080	<0.001	0.207	<0.001	0.001	<0.001
	G				2.13	1.06	1.49	1.78								
Valerate (%)	S	N	2.15	1.20	0.86	1.29	1.79	1.82	0.213	0.736	0.209	0.021	0.253	0.549	0.006	0.012
	G				1.66	1.20	1.25	1.45								
Isovalerate (%)	S	N	5.24	3.96	1.60	2.39	3.29	2.13	0.236	0.663	0.027	<0.001	0.269	<0.001	<0.001	<0.001
	G				2.95	1.44	2.08	2.65								

Rumen development process in goats

L, linear effect of age; Q, quadratic effect of age; C, cubic effect of age; S, supplemental feeding; G, grazing; N, not detectable because the values were too low.

* P value for both feeding systems from 28 to 70 d.

† P for age from 0 to 70 d.

‡ During 0, 7 and 14 d, all goats consumed only milk. After 20 d, goats were randomised into two groups and received solid feed either in S or as G. Four goats were analysed at 0, 7 and 14 d, while eight goats (four goats each for diets S and G) were analysed at 28, 42, 56 and 70 d. The data for 0, 7 and 14 d were merged for S and G because these two groups were fed similarly during this period.

§ For system × age interaction: from 28 to 70 d.

|| For age: from 0 to 70 d.

¶ Molar proportions of VFA.

Table 4. Enzyme activity potentials of the rumen contents of goats at different ages as affected by the different feeding systems (Least square means and standard errors)

Enzymes	Item	System	Age (d)								SEM§	P*			SEM	P for age†		
			0‡	7‡	14‡	28	42	56	70	System		Age	System × age	L		Q	C	
Xylanase (U)¶	E	S	N	N	0.84	1.06	1.73	1.48	2.10	0.133	0.004	<0.001	0.058	0.115	<0.001	0.766	0.273	
		G					1.08	1.01	1.34	1.74				0.138	<0.001	0.190	0.385	
	I	S	N	N	0.23	0.28	1.12	0.79	1.56	0.166	0.610	0.009	0.022	0.189	<0.001	0.716	0.598	
		G					0.75	1.49	0.55	0.62				0.229	0.431	0.008	0.291	
	T	S	N	N	1.06	1.34	2.85	2.27	3.66	0.316	0.096	0.001	0.068	0.259	<0.001	0.895	0.387	
		G					1.83	2.50	1.89	2.36				0.311	0.017	0.127	0.253	
CMCase (U)	E	S	N	N	0.14	0.20	0.22	0.37	0.43	0.037	0.005	0.014	0.001	0.037	<0.001	0.355	0.574	
		G					0.24	0.22	0.28	0.16				0.030	0.461	0.007	0.557	
	I	S	N	N	0.05	0.07	0.12	0.26	0.29	0.048	0.254	0.057	0.075	0.049	0.001	0.556	0.413	
		G					0.19	0.27	0.20	0.24				0.036	0.004	0.019	0.202	
	T	S	N	N	0.19	0.27	0.34	0.63	0.72	0.079	0.465	0.028	0.012	0.081	<0.001	0.430	0.440	
		G					0.43	0.49	0.48	0.39				0.059	0.025	0.006	0.599	
Amylase (U)	E	S	N	ND	1.00	1.77	0.53	1.49	0.94	0.124	<0.001	<0.001	<0.001	0.133	0.351	0.376	0.256	
		G					0.57	0.74	0.52	0.38				0.112	0.003	0.644	0.160	
	I	S	N	ND	0.57	1.76	0.40	0.97	0.66	0.186	<0.001	0.002	0.017	0.237	0.431	0.248	0.042	
		G					0.51	0.41	0.21	0.23				0.067	<0.001	0.845	0.220	
	T	S	N	ND	1.57	3.53	0.93	2.47	1.60	0.281	<0.001	0.001	0.001	0.335	0.352	0.242	0.059	
		G					1.08	1.15	0.73	0.61				0.131	<0.001	0.614	0.544	
Protease (U)	E	S	N	ND	0.59	1.21	0.47	1.23	1.07	0.211	0.093	0.053	0.313	0.190	0.328	0.760	0.296	
		G					1.09	0.59	0.58	0.66								
	I	S	N	ND	0.47	1.22	0.42	0.19	1.08	0.099	0.418	<0.001	<0.001	0.091	0.510	0.027	<0.001	
		G					0.97	0.21	0.80	0.70				0.094	0.330	0.677	0.072	
	T	S	N	ND	1.06	2.42	0.89	1.42	2.14	0.240	0.073	<0.001	0.411	0.223	0.524	0.595	0.014	
		G					2.06	0.80	1.38	1.36								

L, linear effect of age; Q, quadratic effect of age; C, cubic effect of age; E, extracellular; S, supplemental feeding; N, not detectable because the values were too low; G, grazing; I, intracellular; T, total; CMCase, carboxymethylcellulase; ND, not detected due to insufficient sample size.

* P value for both feeding systems from 28 to 70 d.

† P for age from 0 to 70 d.

‡ During 0, 7 and 14 d, all goats consumed only milk. After 20 d, goats were randomised into two groups and received solid feed either in S or as G. Four goats were analysed at 0, 7 and 14 d, while eight goats (four goats each for diets S and G) were analysed at 28, 42, 56 and 70 d. The data for 0, 7 and 14 d were merged for S and G because these two groups were fed similarly during this period.

§ For system × age interaction: from 28 to 70 d.

|| For age: from 0 to 70 d.

¶ For xylanase, CMCase and amylase, 1 U enzyme activity is the amount of enzyme required to release 1 µmol of reducing sugars (xylose or glucose equivalents)/min per g of rumen content; 1 U protease activity is expressed as mg of azocasein hydrolysed/min per g.

Table 5. Microbial copy numbers (log₁₀ copies/DM rumen contents) of rumen contents of goats at different ages as affected by the different feeding systems (Least square means and standard errors)

Microbes	Item	System	Age (d)							SEM§	P*			SEM	P for age†		
			0‡	7‡	14‡	28	42	56	70		System	Age	System × age		L	Q	C
Bacteria	Liquid	S	7.68	9.33	10.07	10.69	10.77	10.80	11.01	0.153	0.001	0.480	0.524	0.156	<0.001	<0.001	<0.001
	Solid	S	7.26	9.15	9.40	9.20	9.91	9.30	9.39	0.180	0.001	0.054	0.547	0.300	<0.001	<0.001	0.017
Archaea	Liquid	S	4.41	5.28	5.97	6.69	7.00	7.43	7.03	0.245	0.007	0.389	0.603	0.268	<0.001	<0.001	0.383
	Solid	S	4.13	4.86	5.47	5.94	6.31	5.97	5.48	0.187	0.008	0.016	0.422	0.262	<0.001	<0.001	0.693
Fungi	Liquid	S	N	4.03	3.68	5.14	4.85	4.40	4.35	0.241	0.384	0.063	0.609	0.215	0.077	0.007	0.314
	Solid	S	N	4.10	3.20	4.84	3.63	3.19	3.66	0.292	<0.001	0.106	0.024	0.347	0.260	0.530	0.146
Protozoa	Liquid	S	N	4.78	5.15	5.65	6.13	6.66	6.27	0.252	0.415	0.036	0.014	0.294	0.001	0.043	0.057
	Solid	S	N	4.35	4.71	5.97	5.73	5.77	5.26	0.265	0.019	0.124	0.977	0.296	0.198	<0.001	0.523
		G				6.32	6.16	6.33	5.80						0.006	0.294	

L, linear effect of age; Q, quadratic effect of age; C, cubic effect of age; S, supplemental feeding; G, grazing; N, not detectable because the values were too low.

* P value for both feeding systems from 28 to 70 d.

† P for age from 0 to 70 d.

‡ During 0, 7 and 14 d, all goats consumed only milk. After 20 d, goats were randomised into two groups and received solid feed either in S or as G. Four goats were analysed at 0, 7 and 14 d, while eight goats (four goats each for diets S and G) were analysed at 28, 42, 56 and 70 d. The data for 0, 7 and 14 d were merged for S and G because these two groups were fed similarly during this period.

§ For system × age interaction: from 28 to 70 d.

|| For age: from 0 to 70 d.

interactions tended to be significant for extracellular xylanase ($P=0.058$), total xylanase ($P=0.068$) and intracellular CMCase ($P=0.075$). From 14 to 70 d, in the supplemented goats, the activity potentials of xylanase and CMCase (extracellular, intracellular and total) increased linearly with age ($P<0.01$), reaching maximum values at 70 d. In grazing goats, the activity potentials of extracellular xylanase ($P<0.001$) and total xylanase ($P=0.017$) increased linearly with age, while age had quadratic effects on the activity potentials of CMCase ($P<0.05$) and intracellular xylanase ($P=0.008$).

For amylase, there were interactions between feeding system and age on the activity potentials of extracellular ($P<0.001$), intracellular ($P=0.017$) and total ($P=0.001$) amylase: the supplemented group exceeded the grazing group for amylase activity potentials at 28, 56 and 70 d, but it was inferior to the grazing group at 42 d. From 14 to 70 d, in the supplemented goats, age had a cubic effect on intracellular amylase ($P=0.042$) and a cubic effect tendency on total amylase ($P=0.059$). In contrast, in the grazing goats, amylase activity potentials decreased linearly with age ($P<0.01$). Similarly, the feeding system \times age interaction ($P<0.001$) was also detected on intracellular protease, as its value in the supplemented group was higher than that in the grazing group at 28, 42 and 70 d, while lower at 56 d. From 14 to 70 d, despite the fact that protease activity potentials were highest at 28 d in both groups, age had cubic effects on intracellular protease activity potentials in the supplemented goats ($P<0.001$), and on total protease activity potentials in both groups ($P=0.014$). Furthermore, extracellular protease ($P=0.093$) and total protease ($P=0.073$) activity potentials in the supplemented goats tended to be greater than those in grazing goats.

Ruminal microbial copy numbers of goats at different ages as affected by the different feeding systems

As illustrated by quantitative PCR results, no feeding system \times age interactions ($P>0.10$) were observed on rumen bacterial and methanogenic archaeal copy numbers (Table 5). The copy numbers of liquid-associated bacteria ($P=0.001$) and archaea ($P=0.007$) were less, while copy numbers of solid-associated bacteria ($P=0.001$) and archaea ($P=0.008$) were greater for the grazing group *v.* the supplemented group. From 0 to 70 d, irrespective of the feeding system, there were increased cubic effects of age ($P<0.05$) on liquid-associated and solid-associated bacterial copy numbers. Furthermore, both liquid-associated and solid-associated archaeal copy numbers increased quadratically with age ($P<0.001$).

Fungi and protozoa were first detected in rumen samples collected at 7 and 14 d, respectively. Feeding system \times age interactions were observed on copy numbers of solid-associated fungi ($P=0.024$) and liquid-associated protozoa ($P=0.014$), for the solid-associated fungi copy number was less while the liquid-associated protozoa copy number was greater for grazing *v.* supplemental feeding at only 28 d. Furthermore, there were no feeding system \times age interactions ($P>0.10$) on liquid-associated fungi and solid-associated

protozoa copy numbers. The solid-associated protozoa copy number ($P=0.019$) was greater for grazing *v.* supplemental feeding. From 7 to 70 d, irrespective of the feeding system, a quadratic effect of age ($P=0.007$) was observed on the liquid-associated fungi copy number, with values increasing from 7 to 28 d, and declining thereafter. The solid-associated fungi copy number in the supplemented group was highest at 28 d, while it increased linearly ($P=0.001$) with age in the grazing group. Moreover, except for the liquid-associated protozoa from 14 to 70 d, copy number in the supplemented goats (cubic tendency, $P=0.057$), age had quadratic effects on protozoa copy number in both groups ($P<0.01$), with their values showing an increase with age during the first 56 d, and a slight decline at 70 d.

Discussion

The rumen development process of young ruminants plays a vital role in host health and nutrition. In ruminants, one of the primary roles of the ruminal microbial ecosystem is to secrete various enzymes that break down plant polymers, and to ferment the released sugars into VFA^(6,33). Furthermore, VFA (mainly butyrate and propionate) are chemical stimuli for the development of the rumen epithelium and papillae, promoting its structural development and absorption activity, thereby providing energy to the host⁽³⁴⁾. Moreover, the rumen epithelium provides a critical barrier between the host and the rumen milieu, and epithelial cells play a key role in recognising the rumen microbiome⁽³⁵⁾. Taken together, these observations suggest that anatomic, functional and microbial development in the rumen is an integrated system, and it is essential to study them together for a better understanding of the process of rumen development.

Effects of the feeding systems on anatomic development, functional achievement and microbial colonisation during the rumen development process

To our knowledge, the present study was the first to demonstrate that supplemental feeding is better than grazing in the promotion of rumen development in young goats. Our outcomes revealed that relative to supplemented goats, more of SAM and less of LAM colonised the rumen of grazing goats. This finding may indicate that the grazing goats consumed more forage than the supplemented goats due to the absence of starter concentrate, and to digest plant polymers, more SAM were needed to colonise the forage⁽²³⁾. This confirms the findings of previous studies on cattle⁽³⁶⁾ and dairy cows⁽³⁷⁾, which indicate that diet has a clear effect on shaping microbial communities. It has also been reported that the solid-associated bacteria account for approximately 70% of the total bacteria⁽²³⁾, thus playing a pivotal role in fibre digestion. The present study revealed that irrespective of the feeding system, liquid-associated bacteria exceeded solid-associated bacteria in numbers in young goats. A primary reason for this might be the fact that the feed intake of the goats in the present study was not as much as that of adult goats, given that live weight at 70 d was only one-third of mature weight. Nevertheless, in

grazing goats, solid-associated fungi exceeded liquid-associated fungi in numbers, suggesting the significance of fungi in the digestion of fibre during early days of ruminant development⁽⁶⁾.

Simultaneously, variation in the feeding system also led to the observation that the rumen fluid in the supplemented group had greater TVFA concentrations and propionate molar proportions than those in the grazing group. This implies that the consumption of some concentrate from 28 d resulted in an earlier initiation of rumen fermentation⁽²⁾. The greater TVFA accumulation and propionate molar proportions in the rumen of supplemented goats probably accounted for the greater rumen papillae area in the present study. Moreover, greater concentrate consumption can lead to higher amylase activity potentials⁽³⁸⁾. Therefore, it is not surprising to find that both extracellular and intracellular amylase activity potentials were higher in supplemental feeding *v.* grazing. Intriguingly, the greater activity potentials of extracellular xylanase and CMCase in supplemented goats suggest that the optimal fibre-degrading capacity occurred when supplemental concentrate was offered⁽⁴⁾. In general, more effective fibre digestion is related to greater solid-associated bacteria number in mature ruminants because solid-associated bacteria exceeded liquid-associated bacteria in number (about 70 *v.* 30%), thus playing a pivotal role in rumen feed digestion^(23,24). Nonetheless, the present study demonstrated in young ruminants the significance of liquid-associated bacteria in digesting fibre, given that the numbers of liquid-associated bacteria exceeded those of solid-associated bacteria, which contrasts with the situation in mature animals. Despite this observation, greater xylanase activity potentials were observed at 28 d, and greater CMCase activity potentials at 28 and 42 d in the grazing group, reflecting enhanced ability to degrade fibre in grazing goats during the transition phase.

During weaning, forage provision was also a major determinant, affecting the rumen development process. In young calves, providing chopped hay (3–4 cm) improved fibre digestibility to a greater extent than finely ground hay (2 mm) during the week after weaning, suggesting that the physical form of forage could affect rumen development⁽³⁹⁾. Meanwhile, provision of chopped hay to calves can promote solid feed DM intake and rumen development (with higher pH)^(17,40). In the present study, although the forage (fresh grass) offered to both supplemented and grazing goats were of the same chemical composition, different forage intake levels (not measured) might also account for variations in rumen development.

Anatomic development, functional achievement and microbial colonisation during the rumen development process

The present study confirmed that the anatomic, functional and microbial development of the rumen is a temporal and successional process, with age being an important factor, irrespective of the feeding system⁽¹³⁾. The ruminal milieu of newborn goats was devoid of VFA and enzyme activities with very low NH₃-N concentration, suggesting the absence of fermentation capacity and enzyme degradation, which concurred

with previous data for calves⁽⁵⁾. Not surprisingly, bacteria colonised rapidly during the first week, similar to a previous observation on calves⁽⁴¹⁾.

In the study of Rey *et al.*⁽⁴¹⁾, the animals were separated from their dams immediately after birth, and placed in individual pens to avoid contact between calves and their dams. Since attendance was not that complete during the birth of the goats in the present study, there was still a possibility of some suckling being unnoticed.

Evidence from previous research on lambs grazing on pasture with dams showed that archaeal colonisation was found at 1–3 d after birth and their population densities reached near 10⁴ copy numbers per g at 1 week of age⁽⁴²⁾, within the range of the present study, even though there was a possibility of contamination with adult animals and between lambs^(7,14). In lambs, whether raised with dams or separately in individual pens, fungi and protozoa appeared relatively later (1–3 weeks), a finding supported by the present results. The maternal vagina microflora directly transmitted to the kid goat^(41,43) and the mother's milk^(44,45) were two main sources of microbes on the first day. As reported in a study on calves⁽⁴⁶⁾, growth of the rumen papillae was also minimal and only slight VFA concentrations were detected in goats during the non-rumination phase. This slow evolution of fermentation capacity could be associated with the intake of only milk without solid feed⁽¹⁸⁾. The present results showed a slight increase in acetate molar proportion at 14 d, thereafter a surge in the acetate:propionate ratio. Such a modification in the VFA profile could be attributed to the fact that microbes could have degraded lactose and other oligosaccharides in the milk into acetate and lactate, as supported by previous studies in calf faeces⁽⁴⁷⁾. Furthermore, amylase activity potentials were relatively high at 14 d, suggesting a vigorous activity of amylolytic microbes that needs further investigation. This might demonstrate the existence of starch degradation capacity even before the availability of solid feed substrate. Collectively, these results suggest that microbes colonise before functional achievements.

During the transition period (3–8 weeks), feed intake was not recorded in the present study. However, it probably increased with age, as the intake of solid feed usually increases significantly after the first month of life^(11,14). Solid feed provides substrate for microbial fermentation in the rumen, which would account for the surge in microbial copy numbers. Fungi and protozoa colonised fastest during this period (28 d), implying that solid feed plays an indispensable role in their colonisation. On the contrary, Fonty *et al.*⁽⁷⁾ claimed that fungi disappeared in lambs when a solid diet was given. Variations in animal species (lambs *v.* goats), diet composition, housing conditions, as well as measurement methods (culture method *v.* molecular method) might account for the discrepancy between the findings of Fonty *et al.*⁽⁷⁾ and the present study.

Along with colonisation of microbes, NH₃-N concentration was found to be comparable with that in adult goats⁽⁴⁸⁾, and TVFA concentration evolved progressively, which agreed with previous studies on calves^(5,49). In accordance with the literature⁽¹¹⁾, we observed a decline in the acetate:propionate

ratio from 8.1 to 3.1–4.0 from 14 to 28 d, which could be attributed to the shift in the goats' diet from milk to forage supplemented with starter concentrate. Lactate resulting from amylolytic microbes that degrade starch in the concentrate and forage can be further converted to propionate by lactate-utilising microbes⁽⁵⁰⁾. Possible enhanced lactate production could account for the increase in the molar proportion of propionate at 28 d.

Presumably increased propionate and butyrate concentrations stimulated the morphological development of the rumen⁽⁴⁶⁾. The rumen papillae length and area increased progressively from 14 to 42 d, which was similar to the findings of a study in calves⁽³⁸⁾. It has also been observed that the number of papillae per unit surface area of rumen declined as the goats matured and grew in size, resulting from the development of the rumen (almost ten times larger in size compared with that at 14 d), with higher age and intake, which stretched the basal epithelial tissue and extended the separation distance between the papillae⁽³⁾. Furthermore, xylanase and CMCase activity potentials increased markedly during the transition phase, indicating a direct reflection of the surge in fibre-degrading capacity.

Presumably such changes in cellulolytic activities reflect changes in cellulolytic bacteria, which reached their highest values at 30 d in lambs⁽⁷⁾. Another explanatory factor could be the progressive increase in fungi, which possess the ability to secrete cellulolytic activity and degrade fibre⁽⁶⁾. In the present study, fungi numbers at 42 d were almost thirty times greater than that at 14 d. In contrast, amylase and protease activity potentials surged from 14 to 28 d, followed by a sharp decline at 42 d. In calves, a similar trend in amylase potential has been observed pre- and post-weaning⁽⁴⁹⁾, which might be due to the stress of weaning.

From 8 weeks onwards, the goats appeared to be mature ruminants with respect to functional achievement of the rumen. At 56 and 70 d, most of the functional parameters did not evolve any further, irrespective of the feeding system, which was in accord with a previous report on calves⁽⁴⁹⁾, demonstrating that functional achievement occurred at 2 months in goats. The quantitative PCR did not detect changes in microbial numbers after 2 months. Similarly, in foals⁽⁴⁾ and lambs⁽⁴²⁾, microbial numbers fluctuated very slightly. In contrast, goat weight gain and rumen enlargement continued beyond 2 months of age⁽⁵⁾. Thus, rumen weight continued to increase, and rumen papillae length and area continued to enlarge, despite complete functional achievement and microbial colonisation of the rumen. Moreover, Tamate *et al.*⁽⁴⁶⁾ also observed in calves an increase in rumen empty weight with a decline in the number of papillae per mm² from 8 to 12 weeks. These results suggest that rumen anatomic development is achieved after 2 months.

It was not feasible to measure the forage intake of grazing goats in the present study, and this is an unfortunate constraint on understanding and interpreting the various responses observed. In addition, variation between individuals is an important determinant during the rumen development process. This is particularly noticeable when presenting microbial data. For example, even though most of the

microbial copy numbers of the same group were similar, we still observed that bacterial copy number from one goat at 14 d was almost sixty times greater than that of another milk-fed goat of the same age. Chen *et al.*⁽³²⁾ also found a significant variation in epimural bacterial diversity among individuals within the same group. This suggests that individual variation needs to be taken into account when exploring the interactions between host and commensal microbes.

Conclusion

From 20 d onwards, when solid feed was started to be offered to the goats, supplemental feeding was superior to grazing, in shaping rumen development, as it promoted greater rumen papillae area, fermentation capacity and enzyme activity potentials. Moreover, irrespective of the feeding system, most of the anatomic, functional and microbial parameters of the rumen development process showed an increase with age, with anatomic and functional parameters evolving progressively from 14 to 42 d of age, while microbial colonisation occurring from birth to 28 d. These results imply that microbial colonisation occurred earlier than functional achievement, with anatomic development occurring last. Offering solid feed and weaning played a vital role in shaping the rumen development process. Further investigations are required to determine the changes in microbial composition and diversity, as well as their interactions with anatomic and functional development.

Acknowledgements

The authors appreciate G. E. Lobley (Rowett Institute of Nutrition and Health, University of Aberdeen, Bucksburn, Aberdeen, UK) for assistance with the revision of the paper.

The authors acknowledge the financial support received from the National Natural Science Foundation of China (grant no. 31320103917), the 'Strategic Priority Research Program – Climate Change: Carbon Budget and Relevant Issues' (grant no. XDA05020700), the 'CAS Visiting Professorship for Senior International Scientists' (grant no. 2010T2S13, 2012T1S0009) and the Hunan Provincial Creation Development Project (2013TF3006).

The authors' contributions are as follows: J. J. and Z. T. designed the research; J. J. and X. L. conducted the research; C. Z. was involved in the animal experiments; J. J., K. A. B. and S. T. performed the statistical analysis and interpreted the results; J. J. wrote the initial draft of the manuscript; K. A. B. critically revised the article for important intellectual content. All authors read and approved the final manuscript.

There are no conflicts of interest to declare.

References

1. Baldwin RLVI, McLeod KR, Klotz JL, *et al.* (2004) Rumen development, intestinal growth and hepatic metabolism in the pre- and postweaning ruminant. *J Dairy Sci* **87**, E55–E65.

2. Wang YH, Xu M, Wang FN, *et al.* (2009) Effect of dietary starch on rumen and small intestine morphology and digesta pH in goats. *Livest Sci* **122**, 48–52.
3. Reynolds C, Dürst B, Lupoli B, *et al.* (2004) Visceral tissue mass and rumen volume in dairy cows during the transition from late gestation to early lactation. *J Dairy Sci* **87**, 961–971.
4. Faubladiere C, Julliard V, Danel J, *et al.* (2013) Bacterial carbohydrate-degrading capacity in foal faeces: changes from birth to pre-weaning and the impact of maternal supplementation with fermented feed products. *Br J Nutr* **110**, 1040–1052.
5. Rey M, Enjalbert F & Monteils V (2012) Establishment of ruminal enzyme activities and fermentation capacity in dairy calves from birth through weaning. *J Dairy Sci* **95**, 1500–1512.
6. Fouts DE, Szpakowski S, Purushe J, *et al.* (2012) Next generation sequencing to define prokaryotic and fungal diversity in the bovine rumen. *PLOS ONE* **7**, e48289.
7. Fonty G, Gouet P, Jouany J-P, *et al.* (1987) Establishment of the microflora and anaerobic fungi in the rumen of lambs. *J Gen Microbiol* **133**, 1835–1843.
8. Lane M, Baldwin R & Jesse B (2002) Developmental changes in ketogenic enzyme gene expression during sheep rumen development. *J Anim Sci* **80**, 1538–1544.
9. Wardrop I & Coombe J (1960) The post-natal growth of the visceral organs of the lamb I. The growth of the visceral organs of the grazing lamb from birth to sixteen weeks of age. *J Agric Sci* **54**, 140–143.
10. Siddons RC (1968) Carbohydrase activities in the bovine digestive tract. *Biochem J* **108**, 839–844.
11. Anderson K, Nagaraja T & Morrill J (1987) Ruminal metabolic development in calves weaned conventionally or early. *J Dairy Sci* **70**, 1000–1005.
12. Li RW, Connor EE, Li C, *et al.* (2012) Characterization of the rumen microbiota of pre-ruminant calves using metagenomic tools. *Environ Microbiol* **14**, 129–139.
13. Jami E, Israel A, Kotser A, *et al.* (2013) Exploring the bovine rumen bacterial community from birth to adulthood. *ISME J* **7**, 1069–1079.
14. Belanche A, Balcells J, De La Fuente G, *et al.* (2010) Description of development of rumen ecosystem by PCR assay in milk-fed, weaned and finished lambs in an intensive fattening system. *J Anim Physiol Anim Nutr* **94**, 648–658.
15. Owens FN, Dubeski P & Hanson C (1993) Factors that alter the growth and development of ruminants. *J Anim Sci* **71**, 3138–3150.
16. Coverdale J, Tyler H, Quigley JD III, *et al.* (2004) Effect of various levels of forage and form of diet on rumen development and growth in calves. *J Dairy Sci* **87**, 2554–2562.
17. Khan MA, Weary DM & von Keyserlingk MA (2011) Hay intake improves performance and rumen development of calves fed higher quantities of milk. *J Dairy Sci* **94**, 3547–3553.
18. Cozzi G, Gottardo F, Mattiello S, *et al.* (2002) The provision of solid feeds to veal calves: I. Growth performance, forestomach development, and carcass and meat quality. *J Anim Sci* **80**, 357–366.
19. Suárez B, Van Reenen C, Stockhofe N, *et al.* (2007) Effect of roughage source and roughage to concentrate ratio on animal performance and rumen development in veal calves. *J Dairy Sci* **90**, 2390–2403.
20. Liu SM, Cai YB, Zhu HY, *et al.* (2012) Potential and constraints in the development of animal industries in China. *J Sci Food Agric* **92**, 1025–1030.
21. Haenlein GF (2001) Past, present, and future perspectives of small ruminant dairy research. *J Dairy Sci* **84**, 2097–2115.
22. Abecia L, Ramos-Morales E, Martinez-Fernandez G, *et al.* (2014) Feeding management in early life influences microbial colonisation and fermentation in the rumen of newborn goat kids. *Anim Prod Sci* **54**, 1449–1454.
23. Yang WZ, Beauchemin KA & Rode LM (2001) Effect of dietary factors on distribution and chemical composition of liquid- or solid-associated bacterial populations in the rumen of dairy cows. *J Anim Sci* **79**, 2736–2746.
24. Chen XL, Wang JK, Wu YM, *et al.* (2008) Effects of chemical treatments of rice straw on rumen fermentation characteristics, fibrolytic enzyme activities and populations of liquid-and solid-associated ruminal microbes *in vitro*. *Anim Feed Sci Technol* **141**, 1–14.
25. Ha JK, Lee SS, Ahn BH, *et al.* (2003) Effects of non-ionic surfactants on enzyme distributions of rumen contents, anaerobic growth of rumen microbes, rumen fermentation characteristics and performances of lactating cows. *Asian-Aust J Anim Sci* **16**, 104–115.
26. Lesmeister K, Tozer P & Heinrichs A (2004) Development and analysis of a rumen tissue sampling procedure. *J Dairy Sci* **87**, 1336–1344.
27. Jiao JZ, Wang PP, He ZX, *et al.* (2014) *In vitro* evaluation on neutral detergent fibre and cellulose digestion by post-ruminal microorganisms in goats. *J Sci Food Agric* **94**, 1745–1752.
28. Eun J-S & Beauchemin K (2005) Effects of a proteolytic feed enzyme on intake, digestion, ruminal fermentation, and milk production. *J Dairy Sci* **88**, 2140–2153.
29. Denman SE & McSweeney CS (2006) Development of a real-time PCR assay for monitoring anaerobic fungal and cellulolytic bacterial populations within the rumen. *FEMS Microbiol Ecol* **58**, 572–582.
30. Hook SE, Northwood KS, Wright A-D, *et al.* (2009) Long-term monensin supplementation does not significantly affect the quantity or diversity of methanogens in the rumen of the lactating dairy cow. *Appl Environ Microbiol* **75**, 374–380.
31. Sylvester JT, Karnati SK, Yu Z, *et al.* (2004) Development of an assay to quantify rumen ciliate protozoal biomass in cows using real-time PCR. *J Nutr* **134**, 3378–3384.
32. Chen YH, Penner GB, Li MJ, *et al.* (2011) Changes in bacterial diversity associated with epithelial tissue in the beef cow rumen during the transition to a high-grain diet. *Appl Environ Microbiol* **77**, 5770–5781.
33. Krause D, Nagaraja T, Wright A, *et al.* (2013) Board-invited review: rumen microbiology: leading the way in microbial ecology. *J Anim Sci* **91**, 331–341.
34. Bergman E (1990) Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol Rev* **70**, 567–590.
35. Malmuthuge N, Li MJ, Fries P, *et al.* (2012) Regional and age dependent changes in gene expression of Toll-like receptors and key antimicrobial defence molecules throughout the gastrointestinal tract of dairy calves. *Vet Immunol Immunopathol* **146**, 18–26.
36. Petri RM, Schwaiger T, Penner GB, *et al.* (2013) Characterization of the core rumen microbiome in cattle during transition from forage to concentrate as well as during and after an acidotic challenge. *PLOS ONE* **8**, e83424.
37. de Menezes AB, Lewis E, O'Donovan M, *et al.* (2011) Microbiome analysis of dairy cows fed pasture or total mixed ration diets. *FEMS Microbiol Ecol* **78**, 256–265.
38. Suárez B, Van Reenen C, Beldman G, *et al.* (2006) Effects of supplementing concentrates differing in carbohydrate

- composition in veal calf diets: I. Animal performance and rumen fermentation characteristics. *J Dairy Sci* **89**, 4365–4375.
39. Montoro C, Miller-Cushon EK, DeVries TJ, *et al.* (2013) Effect of physical form of forage on performance, feeding behavior, and digestibility of Holstein calves. *J Dairy Sci* **96**, 1117–1124.
 40. Castells L, Bach A, Aris A, *et al.* (2013) Effects of forage provision to young calves on rumen fermentation and development of the gastrointestinal tract. *J Dairy Sci* **96**, 5226–5236.
 41. Rey M, Enjalbert F, Combes S, *et al.* (2014) Establishment of ruminal bacterial community in dairy calves from birth to weaning is sequential. *J Appl Microbiol* **116**, 245–257.
 42. Skillman LC, Evans PN, Naylor GE, *et al.* (2004) 16S ribosomal DNA-directed PCR primers for ruminal methanogens and identification of methanogens colonising young lambs. *Anaerobe* **10**, 277–285.
 43. Mändar R & Mikelsaar M (1996) Transmission of mother's microflora to the newborn at birth. *Neonatology* **69**, 30–35.
 44. Hunt KM, Foster JA, Forney LJ, *et al.* (2011) Characterization of the diversity and temporal stability of bacterial communities in human milk. *PLoS ONE* **6**, e21313.
 45. Wise GH & Anderson GW (1939) Factors affecting the passage of liquids into the rumen of the dairy calf. I. Method of administering liquids: drinking from open pail versus sucking through a rubber nipple. *J Dairy Sci* **22**, 697–705.
 46. Tamate H, McGilliard A, Jacobson N, *et al.* (1962) Effect of various dietaries on the anatomical development of the stomach in the calf. *J Dairy Sci* **45**, 408–420.
 47. Shimomura Y & Sato H (2006) Fecal D- and L-lactate, succinate, and volatile fatty acid levels in young dairy calves. *J Vet Med Sci* **68**, 973–977.
 48. Zeng B, Tan ZL, Tang SX, *et al.* (2011) Effects of alkyl polyglycoside, a nonionic surfactant, and forage-to-concentrate ratio on rumen fermentation, amino acid composition of rumen content, bacteria and plasma in goats. *Arch Anim Nutr* **65**, 229–241.
 49. Sahoo A, Kamra D & Pathak N (2005) Pre- and postweaning attributes in faunated and ciliate-free calves fed calf starter with or without fish meal. *J Dairy Sci* **88**, 2027–2036.
 50. Bernalier-Donadille A (2010) Fermentative metabolism by the human gut microbiota. *Gastroenterol Clin Biol* **34**, S16–S22.