

Effects of a novel combination of two mutated acetolactate synthase (ALS) isozymes on resistance to ALS-inhibiting herbicides in flixweed (*Descurainia sophia*)

Research Article

Cite this article: Xu Y, Xu L, Shen J, Li X, Zheng M (2021) Effects of a novel combination of two mutated acetolactate synthase (ALS) isozymes on resistance to ALS-inhibiting herbicides in flixweed (*Descurainia sophia*). *Weed Sci.* **69**: 430–438. doi: [10.1017/wsc.2021.26](https://doi.org/10.1017/wsc.2021.26)

Received: 13 August 2020
Revised: 6 February 2021
Accepted: 21 March 2021
First published online: 5 April 2021

Associate Editor:

Chenxi Wu, Bayer U.S. – Crop Science

Keywords:

ALS isozymes; double mutations; gene expression; tribenuron-methyl resistance

Author for correspondence:

Mingqi Zheng, China Agricultural University, College of Science, No. 2 of Yuan Ming Yuan Xilu, Haidian District, Beijing, 100193, China. (Email: mqzheng@cau.edu.cn)

Yufang Xu¹ , Lina Xu¹, Jing Shen¹, Xuefeng Li² and Mingqi Zheng²

¹Graduate Student, China Agricultural University, College of Science, Beijing, China and ²Professor, China Agricultural University, College of Science, Beijing, China

Abstract

Flixweed [*Descurainia sophia* (L.) Webb ex Prantl] is a notorious broadleaf weed that is widely distributed in winter wheat-growing areas of China and has evolved resistance to tribenuron-methyl mainly due to target-site resistance (TSR) mutations in acetolactate synthase (ALS). In the current research, two ALS genes were identified in tribenuron-methyl-susceptible (TS) or tribenuron-methyl-resistant (TR) *D. sophia*. Resistance mutations of Asp-376-Glu and Pro-197-Ala were identified on ALS1 and ALS2 isozymes in TR *D. sophia*, respectively. The TR *D. sophia* evolved 10,836.3-fold resistance to tribenuron-methyl and displayed cross-resistance to multiple ALS-inhibiting herbicides with different chemical structures. Dose response experiments and ALS activity assay indicated that two mutated ALS isozymes contributed differentially in resistance to tribenuron-methyl, flucetosulfuron, and pyribenzoxim. In addition, the relative expression level of the *ALS1* gene was 2.2- and 1.6-fold higher than *ALS2* genes in TR *D. sophia* at 1 and 7 d after tribenuron-methyl treatment, respectively. In contrast, the relative expression level of *ALS1* and *ALS2* in TS *D. sophia* is similar. This is the first research that explored different roles of ALS isozymes in resistance to ALS-inhibiting herbicides, which might provide a new perspective for the weed resistance management.

Introduction

Acetolactate synthase (ALS) (EC 2.2.1.6), also recognized as acetohydroxyacid synthase (AHAS; EC 4.1.3.18), catalyzes the formation of 2-acetolactate or 2-aceto-2-hydroxybutyrate in the biosynthesis of branched-chain amino acids (Duggleby et al. 2008). It is well known that ALS is the target of ALS-inhibiting herbicides, which have been used worldwide since the early 1980s. Based on their different chemical structures, ALS-inhibiting herbicides are divided into the following classes: sulfonylurea (SU), pyrimidinyl-thiobenzoate (PTB), sulfonylamino-carbonyl-triazolinone (SCT), triazolopyrimidine (TP), and imidazolinone (IMI). To date, more than 160 weed species have evolved resistance to ALS-inhibiting herbicides worldwide due to continuous and intensive application (Heap 2020). Resistance mutations in ALS isozymes that reduce the binding ability of ALS with herbicides are the most important and universal target-site resistance (TSR) mechanisms (Powles and Yu 2010). To date, resistance mutations have been reported at the Ala-122, Pro-197, Ala-205, Asp-376, Arg-377, Trp-574, Ser-653, or Gly-654 loci of the ALS gene (corresponding to the sequence of ALS in *Arabidopsis thaliana*) (Heap 2020; Yu and Powles 2014).

Multiple ALS isozymes have been reported in many plants (Table 1). Existing evidence suggests that different ALS isozymes might play different roles in the growth and development of plants. For example, the *ALS1* and *ALS3* genes are constitutively expressed in all somatic and reproductive tissues of oilseed rape (*Brassica napus* L.), while *ALS2* transcripts have only been detected in flowers and young siliques and are regulated in an organ-specific manner (Ouellet et al. 1992). Two ALS genes (*SurA* and *SurB*) have been detected in tobacco (*Nicotiana tabacum* L.), while the *SurB* gene is consistently expressed at higher levels than the *SurA* gene (Keeler et al. 1993). Similarly, two or more ALS isozymes were also found in various ALS-inhibiting herbicide-resistant weeds, and up to six ALS isozymes were found in quinoa (*Chenopodium quinoa* Willd.) (Mestanza et al. 2015) and monochoria [*Monochoria vaginalis* (Burm. f.) C. Presl ex Kunth] (Imaizumi et al. 2008) (Table 1). In resistant weeds, resistance-endowing mutations harbor only one of the ALS isozymes in a single plant, and the other isozyme is usually a wild-type copy. Although there are some reports for double resistance mutations in an individual plant, whether the mutations occur in the same or different target isozymes is still uncertain. However, in our previous studies, two ALS isozymes carrying resistance mutations were found in an individual tribenuron-methyl-resistant (TR) flixweed

© The Author(s), 2021. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Table 1. The information and characteristics of multiple acetolactate synthase (ALS) isozymes in plants, including weeds.

Plant species	ALS isozymes	Characteristic difference on ALS gene	Reference
Annual bluegrass (<i>Poa annua</i> L.)	Two unnamed ALS	The resistance mutations occurred only in one ALS gene.	McElroy et al. 2013
Bog bulrush [<i>Schoenoplectiella mucronata</i> (L.) J. Jung & H.K. Choi]	ALS1, ALS2, and ALS3	The ALS1 gene was more frequently detected in leaves than the ALS2 and ALS3 genes, and the resistance mutations were identified only in the ALS1 gene.	Scarabel et al. 2010
Cotton (<i>Gossypium hirsutum</i> L.)	A19, A5, and another four unnamed ALS	The A19 and A5 genes constitutively expressed and encoded the main housekeeping ALS gene, and the expression of the other four genes was tissue specific.	Gruła et al. 1995
False flax (<i>Camelina microcarpa</i> Andr. ex DC.)	ALS1 and ALS2	Resistance mutations were identified only in the ALS1 gene.	Hanson et al. 2004
Flixweed [<i>Descurainia sophia</i> (L.) Webb ex Prantl]	ALS1, ALS2, ALS3, and ALS4	The ALS1 and ALS2 genes were constitutively expressed in all populations, while the ALS3 and ALS4 genes were expressed only in some populations.	Xu et al. 2020
Hairy beggarticks (<i>Bidens pilosa</i> L.)	ALS1, ALS2, and ALS3	The resistance mutation of Trp-574-Leu was identified in only one of three ALS genes.	Lamego et al. 2009
Mayweed chamomile (<i>Anthemis cotula</i> L.)	ALS1 and ALS2	Different resistance mutations were identified only in the ALS1 gene.	Intanon et al. 2011
Monochoria [<i>Monochoria vaginalis</i> (Burm f.) C. Presl ex Kunth]	ALS1, ALS2, ALS3, ALS4, ALS5, and ALS6	The resistance mutations occurred only in the ALS1 or ALS3 gene.	Imaizumi et al. 2008
Oilseed rape (<i>Brassica napus</i> L.)	ALS1, ALS2, ALS3, ALS4, and ALS5	The ALS1 and ALS3 genes were constitutively expressed in somatic and reproductive tissues, while the ALS2 gene was expressed only in reproductive tissues. The ALS4 and ALS5 genes were pseudogenes.	Ouellet et al. 1992
Quinoa (<i>Chenopodium quinoa</i> Willd.)	CqAHAS1, CqAHAS2, CqAHAS3, CqAHAS4, CqAHAS5, and CqAHAS6	The CqAHAS1 and CqAHAS2 genes were the functional genes; the other four genes were silent or expressed in a specific tissue.	Mestanza et al. 2015
Rice barnyardgrass [<i>Echinochloa phyllopogon</i> (Stapf) Koso-Pol.]	ALS1 and ALS2	The expression level of the ALS2 gene was significantly higher than that of the ALS1 gene in rapidly growing organs, such as roots, culms, leaf sheaths, and lamina joints.	Iwakami et al. 2012
Russian thistle (<i>Salsola tragus</i> L.)	Two unnamed ALS	The resistance mutation occurred only in one ALS gene.	Warwick et al. 2010
Shortawn foxtail (<i>Alopecurus aequalis</i> Sobol.)	ALS1, ALS2, ALS3, and ALS4	The ALS1 and ALS2 genes were constitutively expressed in all populations, while the ALS3 and ALS4 genes were pseudogenes and were expressed only in some populations.	Iwakami et al. 2017
Sunflower (<i>Helianthus annuus</i> L.)	AHAS1, AHAS2, and AHAS3	Resistance mutations occurred only in the AHAS1 gene, and inducible expression of three AHAS genes was tissue specific and gene dependent.	Breccia et al. 2013; Kolkman et al. 2004
Tobacco (<i>Nicotiana tabacum</i> L.)	SurA and SurB	The SurB gene was consistently expressed at higher levels than the SurA gene in all tobacco organs examined.	Keeler et al. 1993
Wheat (<i>Triticum aestivum</i> L.)	ahas-D1, ahas-B1, and ahas-A1	The resistance mutation of Ser-653-Asn occurred only in the ahas-D1 or ahas-B1 gene, and the mutation on the ahas-D1 gene coded for more-resistant ALS activity.	Pozniak et al. 2004

[*Descurainia sophia* (L.) Webb ex Prantl] (Deng et al. 2017). In the current study, a novel combination of two mutated ALS isozymes is identified in a single TR *D. sophia* plant.

Descurainia sophia is a troublesome weed infesting winter wheat (*Triticum aestivum* L.) and has evolved resistance to tribenuron-methyl since its introduction into China in 1988. Amino acid substitutions in one or multiple ALS isozymes are responsible for *D. sophia*'s resistance to tribenuron-methyl. In addition, enhanced metabolism in TR *D. sophia* mediated by cytochrome P450s heightens *D. sophia* resistance (Deng et al. 2014, 2015, 2017; Xu et al. 2020; Yang et al. 2016, 2018a, 2018b). In the current study, a new resistance mutation combination of mutated ALS1 (Asp-376-Glu) and ALS2 (Pro-197-Ala) is identified in individual *D. sophia*. Do weeds accumulate multiple mutated ALS enzymes in a single plant as a mechanism for combating higher herbicide selection pressure in fields? Are resistance mutations occurring on ALS isozymes randomly, or are some ALS isozymes more preferentially mutated? Do different ALS isozymes play different roles in resistance evolution? Studies of these problems will help to reveal the resistance mechanisms to ALS-inhibiting herbicides. The current research aims to evaluate the contributions of different mutated ALS1 and ALS2 in resistance to different ALS-inhibiting herbicides in *D.*

sophia through (1) determining the herbicide concentration that inhibits 50% of the activity of ALS1 and ALS2 (I_{50}) and (2) comparing the expression levels of ALS1 and ALS2 genes in *D. sophia*.

Materials and Methods

Plant Materials

In 2016, seeds of the TR *D. sophia* population (SD1637) were collected from winter wheat fields in Liaocheng City, China (36.71°N, 115.84°E), where tribenuron-methyl had been used repeatedly for more than 20 yr. The tribenuron-methyl-susceptible (TS) *D. sophia* population (BJ1602) was harvested from the roadside in Beijing (40.21°N, 116.33°E), where tribenuron-methyl is unlikely to be used. To confirm the frequency of resistance mutations, the ALS genes for 50 plants in each population were sequenced, and all plants investigated had the same mutation combination.

Seeds of *D. sophia* were immersed in 20% H₂O₂ solution for 30 min, followed by a 24-h soak in 0.03% gibberellin solution, and germinated in petri dishes at room temperature for 4 d after rinsing with water. Sixteen germinated seeds were planted in one pot containing moist soil, which was kept in artificial climate chambers at

Table 2. Information on primers for *ALS* gene cloning (F_1/R_1), reference gene *18S rRNA* (F_2/R_2) and *ALS* expression determination (F_3/R_3 , F_4/R_4) in tribenuron-methyl-susceptible (TS) and tribenuron-methyl-resistant (TR) *Descurainia sophia*.

Amplification gene	Primers ^a	Sequence(5'→3')	Amplification size
<i>ALS1</i> and <i>ALS2</i>	F_1 R_1	CGCTCCTCTCTGAAGCTACCA CAAACAAACAGCAGTAGCGTCTGAAG	bp 2004 (<i>ALS1</i>)/1998 (<i>ALS2</i>)
<i>18S rRNA</i>	F_2 R_2	TAGTTGGTGAGCGATTGTCTG CTAAGCGGCATAGTCCCTCTAAG	114
<i>ALS1</i>	F_3 R_3	CTCCACCACTTCTCTCTCT CGAAAACACCTCCTTGTT	236
<i>ALS2</i>	F_4 R_4	CCACTTCTTCTCCAGCGA ACCTGAGGATCGAGCGTA	257

^aF, forward primer; R, reverse primer.

25/20 C (light/dark) for a 16-h photoperiod with a photosynthetic photon flux density of approximately 270 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

Determination of Resistance Mutations in *ALS* Genes

A DNA extraction kit (Plant Genomic DNA Kit[®], DP305, Tiangen China, No. 86, Shuang Ying West Road, Changping District, Beijing, China) was used to extract genomic DNA from fresh leaves of TS and TR *D. sophia*. An F_1/R_1 primer pair was designed to amplify the *ALS* genes to full length (Table 2). The polymerase chain reaction (PCR) mixtures and program were the same as those described by Deng et al. (2017). The PCR products were purified from agar gel with a purification kit (TIANGel Midi Purification Kit[®], DP209, Tiangen China). The purified PCR products were ligated to the pLB-Simple vector and then transformed into TOP10 competent cells. Six clones of each DNA sample were selected for sequencing. Resistance mutations were identified by comparing the *ALS* sequence with that of susceptible *D. sophia* (accession no. JQ868736). The *ALS* genes for 50 plants in each population were sequenced to determine the proportion of plants carrying the resistance mutations.

Whole-Plant Dose Response Experiments for *ALS*-inhibiting Herbicides

Whole-plant dose response experiments were used to determine the resistance or cross-resistance levels of TR and TS *D. sophia* populations to different *ALS*-inhibiting herbicides, including tribenuron-methyl (SU), flucetosulfuron (SU), pyribenzoxim (PTB), flucarbazone-sodium (SCT), flumetsulam (TP), and imazethapyr (IMI). *Descurainia sophia* plants at the 4-leaf stage were used for whole-plant dose response experiments. The herbicides were diluted to a series of concentrations with 0.2% Tween-80 solution. The spray doses used for *ALS*-inhibiting herbicides are listed in Table 3. Notably, due to the great difference in the susceptibility to tribenuron-methyl, the rates varied between TS and TR *D. sophia* populations. Herbicides were applied by using an automatic cabinet sprayer (ASP-1098 automatic sprayer, Zhejiang University Xinnong Pesticide Model Technology Development, Hangzhou, China) at a spray volume of 600 L ha⁻¹. Control individuals were treated with water containing 0.2% Tween-80. The aboveground fresh *D. sophia* seedlings were weighed at 21 d after treatment. The experiments were conducted twice with three replicates in one dose.

In Vitro *ALS* Extraction and Activity Assay

Approximately 40 d after planting, 4 g of leaf material was used for *ALS* extraction and activity assays in vitro according to the

methods of Deng et al. (2014). *ALS* activity in vitro was determined by colorimetry (520 nm) with a microplate photometer (Thermo Fisher, Waltham, MA, USA) by measuring acetoin production. The final concentrations in the reaction mixtures for the different herbicides are listed in Table 3. The experiments were repeated twice using independent enzyme extracts with three replicates for each herbicide concentration.

Relative Expression of *ALS1* and *ALS2* in TR and TS *Descurainia sophia*

Each *D. sophia* plant at 40 d after transplanting was treated evenly with 5 mg L⁻¹ technical tribenuron-methyl (15 μl per plant) dissolved in a mixture of acetone and water (4:6 v/v) by using a micro-applicator (Hamilton PB 600 dispenser, Hamilton, Lancaster, PA, USA). The TS population exhibited no observable adverse response to the applied dose. Three plants were selected for RNA extraction by an RNA extraction kit (RNAprep pure Plant Kit[®], DP432, Tiangen China) before tribenuron-methyl treatment (BT) or at 1, 3, 5, and 7 d after tribenuron-methyl treatment (DAT). Plants treated with only an equal volume of 40% acetone were used as controls.

18S rRNA was selected as a reference gene that was confirmed to be stably expressed in *D. sophia* (Yang et al. 2018b). Two primer pairs (F_3/R_3 and F_4/R_4) were designed for quantitative real-time polymerase chain reaction (qPCR) according to the differences between *ALS1* and *ALS2* gene sequences (Table 2). First-strand complementary DNA (cDNA) was synthesized according to the instructions of the FastQuant RT Kit (TIANSscript II RT Kit[®], KR107, Tiangen China). The expression level for *ALS1* or *ALS2* was determined by performing qPCR. The reaction mixtures with a volume of 20 μl consisted of 0.4 μl 50 \times ROX reference dye, 0.6 μl primers, 1 μl diluted cDNA, 7.4 μl ribonuclease-free distillation-distillation H₂O (RNase-free ddH₂O), and 10 μl 2 \times SuperReal PreMix Plus (SuperReal PreMix Plus[®], FP205, Tiangen China). There were four replicates per cDNA. qPCR was carried out with programs of 15-min incubation at 95 C, 40 cycles at 95 C for 10 s, 60 C for 20 s, and 72 C for 32 s. There were four replicates per cDNA and three cDNA samples at each time point in the TS or TR populations.

Statistical Analysis

Whole-Plant Dose Response Experiments and In Vitro *ALS* Activity Assay

The GR₅₀ (herbicide dose causing 50% plant growth reduction) and I₅₀ were calculated using GraphPad Software (v. 5.0, San Diego, CA, USA) with Equation 1. This equation is based on a

Table 3. Herbicide rates of acetolactate synthase (ALS)-inhibiting herbicides in whole-plant dose response experiments and in vitro ALS activity assay.

Herbicide	Rates in whole-plant dose response experiments ^f		Rates for in vitro ALS activity assay
	TS	TR	TS and TR
	g ai ha ⁻¹		μM
Tribenuron-methyl ^a	0.0006, 0.0023, 0.0092, 0.036, 0.15, 0.59, 2.34, 9.38	0.036, 0.15, 0.59, 2.34, 9.38, 37.5, 75, 150	1.0 × 10 ⁻⁵ , 1.0 × 10 ⁻⁴ , 1.0 × 10 ⁻³ , 1.0 × 10 ⁻² , 0.1, 1, 10, 100, 200, 400, 800
Flucetosulfuron ^a	0.0005, 0.0012, 0.0047, 0.019, 0.075, 0.3, 1.2	0.075, 0.3, 1.2, 4.8, 19.2, 76.8, 307.2	1.0 × 10 ⁻⁵ , 1.0 × 10 ⁻⁴ , 1.0 × 10 ⁻³ , 1.0 × 10 ⁻² , 0.1, 1, 10, 100, 500, 1,000
Pyribenzoxim ^b	0.019, 0.075, 0.3, 1.2, 4.8, 19.2, 76.8	0.075, 0.3, 1.2, 4.8, 19.2, 76.8, 307.2	1.0 × 10 ⁻⁵ , 1.0 × 10 ⁻⁴ , 1.0 × 10 ⁻³ , 1.0 × 10 ⁻² , 0.1, 1, 10, 50, 100, 500, 1,000
Flucarbazone-sodium ^c	0.0002, 0.0009, 0.0036, 0.015, 0.059, 0.024, 0.098	0.94, 3.75, 15, 60, 240, 480, 960	1.0 × 10 ⁻⁵ , 1.0 × 10 ⁻⁴ , 1.0 × 10 ⁻³ , 1.0 × 10 ⁻² , 0.1, 1, 10, 100, 500, 1,000
Flumetsulam ^d	0.0017, 0.019, 0.075, 0.3, 1.2, 4.8, 19.2	0.075, 0.3, 1.2, 4.8, 19.2, 76.8, 307.2	1.0 × 10 ⁻⁵ , 1.0 × 10 ⁻⁴ , 1.0 × 10 ⁻³ , 1.0 × 10 ⁻² , 0.1, 1, 10, 100, 500, 1,000
Imazethapyr ^e	0.0012, 0.0047, 0.019, 0.075, 0.3, 1.2, 4.8	0.019, 0.075, 0.3, 1.2, 4.8, 19.2, 76.8	1.0 × 10 ⁻⁵ , 1.0 × 10 ⁻⁴ , 1.0 × 10 ⁻³ , 1.0 × 10 ⁻² , 0.1, 1, 10, 100, 500, 1,000

^{a,b,c,d,e}indicate the herbicide subfamily of sulfonyleurea (SU), pyrimidinyl-thiobenzoate (PTB), sulfonamino-carbonyl-triazolinone (SCT), triazolopyrimidine (TP), and imidazolinone (IMI), respectively.

^fTS, tribenuron-methyl-susceptible; TR, tribenuron-methyl-resistant.

double-sigmoid model, which is constructed as the sum of two different logics. The double-sigmoid model suggests the coexistence of two inhibition targets with different susceptibilities.

$$y = \text{Frac} \times \frac{100}{1 + 10^{(a-x)}} + (1 - \text{Frac}) \times \frac{100}{1 + 10^{(b-x)}} \quad [1]$$

In this equation, y is the percentage of fresh weight or ALS activity (% control); x is the log of the herbicide dose; and a and b are the logarithms of the GR₅₀ (or I₅₀) of two single-sigmoid curves ($b > a$). The *Frac* value is considered to be a putative proportion of the resistance contribution of more-susceptible ALS (Lipovetsky 2010; Tsuneki et al. 2004; Yamato et al. 2013).

Relative Expression of ALS1 and ALS2

The relative expression ratio (as $2^{-\Delta\Delta C_T}$) was calculated by the cycle threshold (C_T) method (Schmittgen and Livak 2008), where $\Delta C_T = C_T$ target gene – C_T internal control gene. Data for the relative expression level of ALS genes were tested for normality using the Kolmogorov-Smirnov test with SPSS software (v. 16.0, IBM, Armonk, NY, USA), and homogeneity of variance was confirmed by Levene's test with SPSS. The data obtained from the relative expression of genes met the assumptions of an independent-samples t -test and ANOVA by testing. The relative expression of two ALS genes between TS or TR *D. sophia* was analyzed by the independent-samples t -test ($P < 0.05$). Dunnett's test at the 5% level of significance in ANOVA was carried out to compare the relative expression of the ALS1 or ALS2 gene at different times in TS or TR *D. sophia*.

Results and Discussion

Determination of Resistance Mutations in ALS Genes

Two ALS genes with full lengths of 2,004 bp (ALS1) and 1,998 bp (ALS2) were cloned from TS (BJ1602) and TR (SD1637) *D. sophia*. A new combination of resistance mutations (ALS1 with Asp-376-Glu, ALS2 with Pro-197-Ala) was identified from a single TR *D. sophia* plant in the current study. The ALS1 and ALS2 genes in TS and TR *D. sophia* show high homology. A single different nucleotide caused the resistance mutation at the site of Pro-197 (in ALS2) or Asp-376 (in ALS1). Fifty TS or TR *D. sophia* plants

were sequenced, and the ALS1 or ALS2 of each plant was the same in each *D. sophia* population, which indicated that the genetic backgrounds of the different plants were very similar or the same.

Two resistance mutations on ALS in a single plant were also reported in kochia [*Bassia scoparia* (L.) A.J. Scott; syn. *Kochia scoparia* (Linn.) Schrad] (Warwick et al. 2008), rigid ryegrass (*Lolium rigidum* Gaudin) (Kaundun et al. 2012), and Palmer amaranth (*Amaranthus palmeri* S. Watson) (Singh et al. 2019), but it is not clear whether these mutations occur in the same ALS isozyme or in different ALS isozymes. To the best of our knowledge, a total of eight different amino acid mutations have been identified at Pro-197 (substituted by Ser, Thr, Leu, His, Ala, or Arg), Asp-376 (by Glu), and Trp-574 (by Leu) in ALS isozymes in TR *D. sophia* (Xu et al. 2020). Initially, resistance mutations were only identified in one ALS isozyme, and the most common resistance mutation was identified at the site of Pro-197 (Xu et al. 2020). However, resistance mutations at Asp-376 or Trp-574 in ALS, which can cause higher levels of resistance to tribenuron-methyl and broader cross-resistance to ALS-inhibiting herbicides than resistance mutations at Pro-197, were identified only in a few TR *D. sophia* (Deng et al. 2017; Xu et al. 2015; Yang et al. 2018a). Deng et al. (2017) first reported that two mutated ALS isozymes (Trp-574-Leu in ALS1, Pro-197-Thr in ALS2) in a single TR *D. sophia* plant exhibit a higher resistance level (789.3-fold) to tribenuron-methyl than TR *D. sophia* populations carrying a single mutated isozyme (211.8- and 366.3-fold) and exhibit cross-resistance to more ALS-inhibiting herbicides. In the current study, a novel combination of two mutated ALS isozymes (Asp-376-Glu in ALS1 and Pro-197-Ala in ALS2) was identified in a single TR *D. sophia* plant. Hence, we speculated that the accumulation of two or multiple mutated ALS isozymes in a single plant results in a higher level of resistance.

Whole-Plant Dose Response Experiments for ALS-inhibiting Herbicides

Whole-plant dose response experiments established the different sensitivities of TS (BJ1602) and TR (SD1637) *D. sophia* populations to different ALS-inhibiting herbicides (Table 4; Figure 1). The TS *D. sophia* were killed completely by tribenuron-methyl at a dose of 0.59 g ai ha⁻¹, while all TR *D. sophia* survived at the highest dose of 150 g ai ha⁻¹ (Figure 1A). The resistance index

Table 4. The GR₅₀ and I₅₀ values of acetolactate synthase (ALS)-inhibiting herbicides for tribenuron-methyl-susceptible (TS) and tribenuron-methyl-resistant (TR) *Descurainia sophia*.^a

Herbicide	TS		TR					
	GR ₅₀	I ₅₀	GR ₅₀	Frac ^b	RI	I ₅₀	Frac	RI
	g ai ha ⁻¹	μM	g ai ha ⁻¹			μM		
Tribenuron-methyl	0.03	0.17	325.09	0.09	10,836.33	508.39	0.08	2,990.53
Flucetosulfuron	0.03	3.07	14.51	0.19	483.67	295.12	0.16	96.13
Pyribenzoxim	1.29	0.13	14.00	0.18	10.85	107.52	0.16	827.08
Flucarbazone-sodium	0.01	0.03	28.39	NA	2,839.00	1.97	NA	65.66
Flumetsulam	0.59	0.45	20.15	NA	34.15	18.12	NA	40.27
Imazethapyr	0.05	5.75	3.49	NA	69.80	29.62	NA	5.15

^aAbbreviations: GR₅₀, herbicide rates causing 50% plant growth reduction; I₅₀, herbicide rates inhibiting 50% ALS activity; RI, resistance index, GR₅₀ value or I₅₀ value of TR populations divided by that of TS population; NA, not applicable.

^bFrac is considered to be a putative proportion of resistance contribution of more susceptible ALS.




Figure 1. Dose response curves of tribenuron-methyl-susceptible (TS, BJ1602-TS) and tribenuron-methyl-resistant (TR, SD1637-TR) *Descurainia sophia* treated with acetolactate synthase (ALS)-inhibiting herbicides: (A) tribenuron-methyl, (B) flucetosulfuron, (C) pyribenzoxim, (D) flucarbazone-sodium, (E) flumetsulam, and (F) imazethapyr. The experiments had three replicates per herbicide dose and were repeated twice. Vertical bars represent the standard error of two experiments.

(RI) of TR *D. sophia* (SD1637) was 10,836.33, which is higher than that of the TR populations carrying any other known resistance mutations (Deng et al. 2017; Yang et al. 2018a). For example, the populations with a single resistance mutation of Pro-197-His, Pro-197-Leu, Pro-197-Thr, Pro-197-Ser, Trp-574-Leu, or Asp-376-Glu in the ALS1 isozyme exhibited 205.8-, 250.0-, 708.4-, 263.7-, 485.3-, or 2,844.7-fold resistance to tribenuron-methyl, respectively (Yang et al. 2018a), and the TR *D. sophia* population with two mutated ALS isozymes (Trp-574-Leu in ALS1, Pro-197-Thr in ALS2) evolved a 789.3-fold resistance to tribenuron-methyl (Deng et al. 2017). Therefore, the presence of two mutated ALS isozymes in individual plants helped weeds survive higher doses of ALS-inhibiting herbicide compared with weeds carrying a single mutated ALS isozyme. In addition, the accumulation of multiple ALS mutations in individual plants was also observed in *A. palmeri* (Singh et al. 2019). In recent years, double mutations have been reported in a variety of resistant weeds, and these double mutations lead to weeds evolving higher resistance levels than weeds with a single resistance mutation. However, it is not clear whether double

resistance mutations occur on the same or different target isozymes. For example, double mutations (Ala-251-Val and Phe-273-Val) in *psbA* resulted in *Amaranthus* spp. evolving a higher resistance to linuron and diuron compared with each single substitution (Davis et al. 2020). Goosegrass [*Eleusine indica* (L.) Gaertn.] with double mutations (Thr-102-Ile and Pro-106-Ser in 5-enolpyruvylshikimate-3-phosphate synthase [EPSPS]) developed a more than 32-fold higher resistance to glyphosate than *E. indica* carrying a single mutation (Pro-106-Ser) (Yu et al. 2015). Similar to *E. indica*, a different EPSPS double mutation (Thr-102-Ile and Pro106-Thr) conferring a high level of glyphosate resistance was reported in tetraploid beggarticks (*Bidens pilosa* L.) (Takano et al. 2020). Notably, a triple mutation (Thr-102-Ile, Ala-103-Val, and Pro-106-Ser) in EPSPS was reported, providing a 314-fold resistance to glyphosate compared with the wild-type strain in smooth pigweed (*Amaranthus hybridus* L.) (Perotti et al. 2018).

The results obtained indicate that TR *D. sophia* exhibits obvious cross-resistance to representative herbicides of ALS-inhibiting




Figure 2. The acetolactate synthase (ALS) activity in vitro of tribenuron-methyl-susceptible (TS, BJ1602-TS) and tribenuron-methyl-resistant (TR, SD1637-TR) populations inhibited by ALS-inhibiting herbicides: (A) tribenuron-methyl, (B) flucetosulfuron, (C) pyribenzoxim, (D) flucarbazone-sodium, (E) flumetsulam, and (F) imazethapyr. The experiments had three replicates per herbicide concentration and were repeated twice. Vertical bars represent the standard error of two experiments.

herbicides with different chemical structures (Table 4). The cross-resistance levels to flucarbazone-sodium (SCT) and flucetosulfuron (SU) are much higher than to imazethapyr (IMI), flumetsulam (TP), and pyribenzoxim (PTB). Although the ALS isozymes are the common targets of these herbicides, their binding ability with ALS isozymes may be different due to different chemical structures. Therefore, TR *D. sophia* displays different cross-resistance levels to different ALS-inhibiting herbicides. The dose response of TR *D. sophia* to different ALS-inhibiting herbicides is different (Figure 1). The whole-plant dose response to tribenuron-methyl, flucetosulfuron, and pyribenzoxim can be fit to double-sigmoid models, demonstrating that two ALS isozymes in TR *D. sophia* have different sensitivities to these herbicides (Figure 1A–C). However, the curves fit for flucarbazone-sodium, flumetsulam, and imazethapyr are more consistent with single-sigmoid models, indicating that the ALS isozymes in TR *D. sophia* have similar sensitivities to these herbicides (Figure 1D–F). *Frac* is usually considered to be the proportion of resistance contributed by a more susceptible ALS isozyme to ALS-inhibiting herbicides (Yamato et al. 2013). The *Frac* value of TR *D. sophia* to tribenuron-methyl, flucetosulfuron, and pyribenzoxim indicates that the resistance contribution of the two ALS isozymes to these herbicides is obviously different. The resistance contributions to flucarbazone-sodium, imazethapyr, and flumetsulam are similar for the two ALS isozymes (Table 4).

In Vitro ALS Activity Assay

The results for I_{50} demonstrate that the herbicide concentrations inhibiting ALS activity in TR *D. sophia* increase greatly compared with those in TS *D. sophia* (Table 4). The curve-fitting models for the whole-plant dose response to and ALS activity inhibition by the same herbicide are nearly consistent for the TR population (Figures 1 and 2). The inhibition of ALS activity by

tribenuron-methyl, flucetosulfuron, and pyribenzoxim can be fit by double-sigmoid models, indicating that two ALS isozymes in TR *D. sophia* contributed differently to the resistance to tribenuron-methyl, flucetosulfuron, and pyribenzoxim (Figure 2A–C). The ALS activity inhibition by flucarbazone-sodium, flumetsulam, and imazethapyr can be fit to single-sigmoid models, demonstrating that the two ALS isozymes exhibit similar contributions to the resistance to flucarbazone-sodium, flumetsulam, and imazethapyr (Figure 2D–F). One double-sigmoid curve can be regarded as the combination of two independent single-sigmoid curves. Compared with the single-sigmoid curves, the double-sigmoid curves for tribenuron-methyl, flucetosulfuron, and pyribenzoxim in TR *D. sophia* have a plateau between 1.0 to 10 μM , 10 to 100 μM , and 1.0 to 100 μM , respectively (Figure 2A–C), indicating that the more susceptible of the two ALS isozymes was saturated first. The activity of another ALS isozyme is gradually saturated with increasing herbicide concentration. In addition, the R^2 obtained by weighted least-squares for the GR_{50} and I_{50} values for six ALS-inhibiting herbicides are determined to be 0.908, which indicates that the reduced ALS activity in response to herbicides is highly related to herbicide resistance in TR *D. sophia*. In particular, the I_{50} -based RI value of flucarbazone-sodium is much lower than the GR_{50} -based RI value. Given that the two ALS isozymes contribute equally to resistance to flucarbazone-sodium, the higher RI value (GR_{50} -based) may also be due to non-target site resistance (NTSR) mechanisms. To date, NTSR to flucarbazone-sodium has been reported only in wild oat (*Avena fatua* L.) (Burns et al. 2018). Further research is needed to determine whether the NTSR mechanisms confer flucarbazone-sodium resistance in TR *D. sophia*.

Reduced ALS affinity with ALS-inhibiting herbicides, which is caused by resistance mutations in ALS isozymes, is the major TSR mechanism for *D. sophia* and other weed species (Deng et al. 2014, 2017; Heap 2020; Xu et al. 2015; Yang et al. 2018a). However, most




Figure 3. The relative expression of *ALS1* and *ALS2* genes in (A) tribenuron-methyl-susceptible (TS) or (B) tribenuron-methyl-resistant (TR) *Descurainia sophia* plants before tribenuron-methyl treatment (BT) or at 1, 3, 5, and 7 d after tribenuron-methyl treatment (DAT). Each data is the mean \pm SE of four replicates. Means with different capital letters indicate the relative expression level of *ALS1* was significantly higher than that of *ALS2* at the same time point on level of $P < 0.05$. Means without letters show the relative expression level of two *ALS* genes at the same time point exhibited no significant difference.




Figure 4. The relative expression levels of *ALS1* (A) or *ALS2*, B) gene before tribenuron-methyl treatment (BT) or at 1, 3, 5, and 7 d after tribenuron-methyl treatment (DAT) in tribenuron-methyl-susceptible (TS) or (B) tribenuron-methyl-resistant (TR) *Descurainia sophia*. Each data is the mean \pm SE of four replicates. Means with different capital (or lowercase) letters mean the relative expression level of *ALS* genes in TS (or TR) plants exhibited significant differences ($P < 0.05$) at BT or at 1, 3, 5, and 7 DAT, respectively.

studies on the TSR mechanism for ALS have mainly focused on the sites and amino acids of resistance mutations in ALS isozymes. With the presence of multiple mutations in a single gene or multiple mutated target isozymes in an individual plant, the resistance patterns and mechanisms will become more complex. For example, the EPSPS crystal structure from *Escherichia coli* revealed that the mutation Thr-97-Ile (Thr-102 in plants) in the presence of Ser-101 (Ser-106 in plants) causes a shift of residue Gly-96 toward the glyphosate binding site, which impairs the efficient binding of glyphosate (Funke et al. 2009). Compared with EPSPS with a double mutation, EPSPS with three mutations may form a new conformational structure with a smaller active site, leading to a higher exclusion of glyphosate (Perotti et al. 2018). The results here also indicate that the resistance contribution of two mutated ALS isozymes to tribenuron-methyl, flucetosulfuron, and pyribenzoxim is different, but there is no obvious difference in resistance to flucarbazone-sodium, flumetsulam, and imazethapyr (Table 4).

Relative Expression of *ALS1* and *ALS2* in TR and TS *Descurainia sophia*

The results in Figure 3 compare the relative expression levels of *ALS1* and *ALS2* in TS or TR *D. sophia* before treatment or at 1, 3, 5, and 7 DAT. At 1 and 7 DAT, the relative expression level of *ALS1* in TR *D. sophia* is significantly higher than that in *ALS2* by 2.2- and 1.6-fold, respectively. No obvious differences are observed before and at 3 and 5 DAT (Figure 3B). In contrast, the relative expression levels of *ALS1* and *ALS2* in TS *D. sophia* are not obviously different before or after tribenuron-methyl treatment (Figure 3A).

The results in Figure 4 compare the relative expression levels of *ALS1* (or *ALS2*) before or at 1, 3, 5, and 7 DAT in the same population. Compared with BT, the relative expression level of *ALS1* in TR *D. sophia* is significantly increased 1.5- and 2.6-fold at 1 and 7 DAT (Figure 4A). *ALS2* is not induced or slightly inhibited before 5 DAT and is increased 1.8-fold at 7 DAT (Figure 4B). In TS

D. sophia, the relative expression level of *ALS1* (or *ALS2*) at 1, 3, 5, and 7 DAT was approximately 0.7- (0.7-), 0.3- (0.4-), 0.4- (0.6-) and 0.6- (0.7-) fold that of BT, respectively (Figure 4).

The expression level of *ALS1* is significantly higher than that of *ALS2* in TR *D. sophia*, and *ALS1* is more easily induced by tribenuron-methyl than *ALS2*. Therefore, *ALS1* might be a preferred mutated target for weeds to gain resistance. The inducible expression of *ALS* has also been reported in other weeds. For example, the transcript levels for the *ALS* genes in *B. scoparia* were 7-fold higher 24 h after chlorsulfuron treatment (Varanasi et al. 2017). The transcript levels for *AHAS3* in the roots of sunflower (*Helianthus annuus* L.) were induced 4- to 5-fold by imazapyr (Breccia et al. 2013). *ALS* expression was upregulated 15.5- to 21.0-fold after mesosulfuron-methyl treatment in shortawn foxtail (*Alopecurus aequalis* Sobol.) (Zhao et al. 2018). Higher gene expression levels may lead to the accumulation of *ALS* isozymes, which can thus enhance the tolerance or resistance of weeds to herbicides by compensating for reduced *ALS* activity. However, enhanced transcription does not necessarily give rise to increased translation level, and posttranslational regulation may affect actual protein levels. Therefore, the contribution of increased *ALS* gene transcription to herbicide resistance needs more supporting data.

In summary, the novel resistance mutation combination (Asp-376-Glu in *ALS1* and Pro-197-Ala in *ALS2*) leads to TR *D. sophia* evolving higher and broader resistance or cross-resistance to *ALS*-inhibiting herbicides than *D. sophia* with a single mutated *ALS* isozyme. The resistance contribution of two mutated *ALS* isozymes is different, but the contribution ratio of each *ALS* isozyme cannot be determined. The presence of two or multiple mutated *ALS* isozymes is probably the result of continuous herbicide selection on the basis of a single resistance mutation, which helps weeds survive higher doses of herbicides. However, it has to be pointed out that this study is based on a TS/TR *D. sophia* population. These results and speculations need to be verified by studying more *D. sophia* populations and even other weed species.

Acknowledgments. This work was sponsored by the National Natural Science Foundation of China (31672047). No conflicts of interest have been declared.

References

- Breccia G, Vega T, Felitti SA, Picardi L, Nestares G (2013) Differential expression of acetoxyacid synthase genes in sunflower plantlets and its response to imazapyr herbicide. *Plant Sci* 208:28–33
- Burns EE, Keith BK, Talbert LE, Dyer WE (2018) Non-target site resistance to flucarbazone, imazamethabenz and pinoxaden is controlled by three linked genes in *Avena fatua*. *Weed Res* 58:8–16
- Davis G, Letarte J, Grainger CM, Rajcan I, Tardif FJ (2020) Widespread herbicide resistance in pigweed species in Ontario carrot production is due to multiple photosystem II mutations. *Can J Plant Sci* 100:56–67
- Deng W, Cao Y, Yang Q, Liu MJ, Mei Y, Zheng MQ (2014) Different cross-resistance patterns to AHAS herbicides of two tribenuron-methyl resistant flixweed (*Descurainia sophia* L.) biotypes in China. *Pestic Biochem Physiol* 112:26–32
- Deng W, Liu MJ, Yang Q, Mei Y, Li XF, Zheng MQ (2015) Tribenuron-methyl resistance and mutation diversity of Pro197 in flixweed (*Descurainia sophia* L.) accessions from China. *Pestic Biochem Physiol* 117:68–74
- Deng W, Yang Q, Zhang YZ, Jiao HT, Mei Y, Li XF, Zheng MQ (2017) Cross-resistance patterns to acetolactate synthase (*ALS*)-inhibiting herbicides of flixweed (*Descurainia sophia* L.) conferred by different combinations of *ALS* isozymes with a Pro-197-Thr mutation or a novel Trp-574-Leu mutation. *Pestic Biochem Physiol* 136:41–45
- Duggleby RG, McCourt JA, Guddat LW (2008) Structure and mechanism of inhibition of plant acetoxyacid synthase. *Pestic Biochem Physiol* 46:309–324
- Funke T, Yang Y, Han HJ, Healy-Fried M, Olesen S, Becker A, Schonbrunn E (2009) Structural basis of glyphosate resistance resulting from the double mutation Thr⁹⁷→Ile and Pro¹⁰¹→Ser in 5-enolpyruvylshikimate-3-phosphate synthase from *Escherichia coli*. *J Biol Chem* 284:9854–9860
- Grula JW, Hudspeth RL, Hobbs SL, Anderson DM (1995) Organization, inheritance and expression of acetoxyacid synthase genes in the cotton allotetraploid *Gossypium hirsutum*. *Plant Mol Biol* 28:837–846
- Hanson BD, Park KW, Mallory-Smith CA, Thill DC (2004) Resistance of *Camelina microcarpa* to acetolactate synthase inhibiting herbicides. *Weed Res* 44:187–194
- Heap I (2020) International Herbicide-Resistant Weed Database. <http://www.weedscience.com>. Accessed: November 19, 2020
- Imaizumi T, Wang GX, Ohsako T, Tominaga T (2008) Genetic diversity of sulfonylurea-resistant and -susceptible *Monochoria vaginalis* populations in Japan. *Weed Res* 48:187–196
- Intanon S, Perez-Jones A, Hulting AG, Mallory-Smith CA (2011) Multiple Pro¹⁹⁷ *ALS* substitutions endow resistance to *ALS* inhibitors within and among mayweed chamomile populations. *Weed Sci* 59:431–437
- Iwakami S, Shimono Y, Manabe Y, Endo M, Shibaike H, Uchino A, Tominaga T (2017) Copy number variation in acetolactate synthase genes of thifensulfuron-methyl resistant *Alopecurus aequalis* (shortawn foxtail) accessions in Japan. *Front Plant Sci* 8:254
- Iwakami S, Uchino A, Watanabe H, Yamasue Y, Inamura T (2012) Isolation and expression of genes for acetolactate synthase and acetyl-CoA carboxylase in *Echinochloa phyllopogon*, a polyploid weed species. *Pest Manag Sci* 68:1098–1106
- Kaundun SS, Dale RP, Bailly GC (2012) Molecular basis of resistance to herbicides inhibiting acetolactate synthase in two rigid ryegrass (*Lolium rigidum*) populations from Australia. *Weed Sci* 60:172–178
- Keeler SJ, Sanders P, Smith JK, Mazur BJ (1993) Regulation of tobacco acetolactate synthase gene expression. *Plant Physiol* 102:1009–1018
- Kolkman JM, Slabaugh MB, Bruniard JM, Berry S, Bushman BS, Olungu C, Maes N, Abratti G, Zambelli A, Miller JF, Leon A, Knapp SJ (2004) Acetoxyacid synthase mutations conferring resistance to imidazolinone or sulfonylurea herbicides in sunflower. *Theor Appl Genet* 109:1147–1159
- Lamego FP, Charlson D, Delatorre CA, Burgos NR, Vidal RA (2009) Molecular basis of resistance to *ALS*-inhibitor herbicides in greater beggarticks. *Weed Sci* 57:474–481
- Lipovetsky S (2010) Double logistic curve in regression modeling. *J Appl Stat* 37:1785–1793
- McElroy JS, Flessner ML, Wang Z, Dane F, Walker RH, Wehtje GR (2013) A Trp⁵⁷⁴ to Leu amino acid substitution in the *ALS* gene of annual bluegrass (*Poa annua*) is associated with resistance to *ALS*-inhibiting herbicides. *Weed Sci* 61:21–25
- Mestanza C, Riegel R, Silva H, Vásquez SC (2015) Characterization of the acetoxyacid synthase multigene family in the tetraploid plant *Chenopodium quinoa*. *Electron J Biotechnol* 18:393–398
- Ouellet T, Rutledge RG, Miki BL (1992) Members of the acetoxyacid synthase multigene family of *Brassica napus* have divergent patterns of expression. *Plant J* 2:321–330
- Perotti VE, Larran AS, Palmieri VE, Martinatto AK, Alvarez CE, Tuesca D, Permingeat HR (2018) A novel triple amino acid substitution in the EPSPS found in a high-level glyphosate resistant *Amaranthus hybridus* population from Argentina. *Pest Manag Sci* 75:1242–1251
- Powles SB, Yu Q (2010) Evolution in action: plants resistant to herbicides. *Annu Rev Plant Biol* 61:317–347
- Pozniak CJ, Birk IT, O'Donoghue LS, Ménard C, Hucl PJ, Singh BK (2004) Physiological, molecular characterization of mutation-derived imidazolinone resistance in spring wheat. *Crop Sci* 44:1434–1443
- Scarabel L, Locascio A, Furini A, Sattina M, Varotto S (2010) Characterisation of *ALS* genes in the polyploid species *Schoenoplectus mucronatus* and implications for resistance management. *Pest Manag Sci* 66:337–344
- Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C_T method. *Nat Protoc* 3:1101–1108

- Singh S, Singh V, Salas-Perez RA, Bagavathiannan MV, Lawton-Rauh A, Romaburgos N (2019) Target-site mutation accumulation among ALS inhibitor-resistant Palmer amaranth. *Pest Manag Sci* 75:1131–1139
- Takano HK, Fernandes VN, Adegas FS, Oliveira RS Jr, Westra P, Gaines TA, Dayan FE (2020) A novel TIPT double mutation in EPSPS conferring glyphosate resistance in tetraploid *Bidens subalternans*. *Pest Manag Sci* 76:95–102
- Tsuneki H, You Y, Toyooka N, Kagawa S, Kobayashi S, Sasaoka T, Nemoto H, Kimura I, Dani JA (2004) Alkaloids indolizidine 235B', quinolizidine 1-epi-207I, and the tricyclic 205B are potent and selective noncompetitive inhibitors of nicotinic acetylcholine receptors. *Mol Pharmacol* 66:1061–1069
- Varanasi VK, Bayramov S, Prasad PVV, Jugulam, M (2017) Expression profiles of *psbA*, *ALS*, *EPSPS*, and other chloroplastic genes in response to PSII-, ALS-, and EPSPS-inhibitor treatments in *Kochia scoparia*. *Am J Plant Sci* 8:451–470
- Warwick SI, Sauder CA, Beckie HJ (2010) Acetolactate synthase (ALS) target-site mutations in ALS inhibitor-resistant Russian thistle (*Salsola tragus*). *Weed Sci* 58:244–251
- Warwick SI, Xu R, Sauder C, Beckie HJ (2008) Acetolactate synthase target-site mutations and single nucleotide polymorphism genotyping in ALS-resistant kochia (*Kochia scoparia*). *Weed Sci* 56:797–806
- Xu X, Liu GQ, Chen SL, Li BH, Liu XM, Wang XY, Fan CQ, Wang GQ, Ni HW (2015) Mutation at residue 376 of AHAS confers tribenuron-methyl resistance in flaxweed (*Descurainia sophia*) accession from Hebei Province, China. *Pestic Biochem Physiol* 125:62–68
- Xu YF, Xu LN, Li XF, Zheng MQ (2020) Investigation of resistant level to tribenuron-methyl, diversity and regional difference of the resistance mutations on acetolactate synthase (ALS) isozymes in *Descurainia sophia* L. from China. *Pestic Biochem Physiol* 169:104653
- Yamato S, Sada Y, Ikeda H (2013) Characterization of acetolactate synthase from sulfonylurea herbicide-resistant *Schoenoplectus juncooides*. *Weed Biol Manag* 13:104–113
- Yang Q, Deng W, Li XF, Yu Q, Bai LY, Zheng M (2016) Target-site and non-target-site based resistance to the herbicide tribenuron-methyl in flaxweed (*Descurainia sophia* L.). *BMC Genomics* 17:551–564
- Yang Q, Deng W, Wang SP, Liu HJ, Li XF, Yu Q, Zheng MQ (2018a) Effects of resistance mutations of Pro197, Asp376 and Trp574 on the characteristics of acetohydroxyacid synthase (AHAS) isozymes. *Pest Manag Sci* 74:1870–1879
- Yang Q, Li JY, Shen J, Xu YF, Liu HJ, Deng W, Li XF, Zheng MQ (2018b) Metabolic resistance to acetolactate synthase inhibiting herbicide tribenuron-methyl in *Descurainia sophia* L. mediated by cytochrome P450 enzymes. *J Agric Food Chem* 66:4319–4327
- Yu Q, Jalaludin A, Han HP, Chen M, Sammons RD, Powles SB (2015) Evolution of a double amino acid substitution in the 5-enolpyruvylshikimate-3-phosphate synthase in *Eleusine indica* conferring high-level glyphosate resistance. *Plant Physiol* 167:1440–1447
- Yu Q, Powles SB (2014) Resistance to AHAS inhibitor herbicides: current understanding. *Pest Manag Sci* 70:1340–1350
- Zhao N, Yan YY, Wang HZ, Bai S, Wang Q, Liu W, Wang JX (2018) Acetolactate synthase overexpression in mesosulfuron-methyl-resistant shortawn foxtail (*Alopecurus aequalis* Sobol.): reference gene selection and herbicide target gene expression analysis. *J Agric Food Chem* 66:9624–9634