

Precise Near-Infrared Radial Velocities

Peter Plavchan¹, Peter Gao², Jonathan Gagne³, Elise Furlan⁴,
Carolyn Brinkworth⁵, Michael Bottom², Angelle Tanner⁶, Guillem
Anglada-Escude⁷, Russel White⁸, Cassy Davison⁸, Sean Mills⁹, Chas
Beichman⁴, John Asher Johnson¹⁰, David Ciardi⁴, Kent Wallace¹¹,
Bertrand Mennesson¹¹, Gautam Vasisht¹¹, Lisa Prato¹², Stephen
Kane¹³, Sam Crawford¹¹, Tim Crawford¹¹, Keeyoon Sung¹¹, Brian
Drouin¹¹, Sean Lin¹¹, Stephanie Leifer¹¹, Joe Catanzarite¹⁴, Todd
Henry¹⁵, Kaspar von Braun¹², Bernie Walp¹⁶, Claire Geneser¹, Nick
Ogden¹, Andrew Stufflebeam¹, Garrett Pohl¹ and Joe Regan¹

¹Missouri State University

email: plavchan@missouristate.edu

²Caltech, ³University of Montreal, ⁴NASA Exoplanet Science Institute, ⁵National Center for
Atmospheric Research/ University Corporation for Atmospheric Research, ⁶Mississippi State
University, ⁷University College London, ⁸Georgia State University, ⁹University of
Chicago, ¹⁰Harvard, ¹¹NASA Jet Propulsion Laboratory, ¹²Lowell Observatory, ¹³San Francisco
State University, ¹⁴SETI Institute, ¹⁵Georgia State University, ¹⁶NASA Ames

Abstract. We present the results of two 2.3 μm near-infrared (NIR) radial velocity (RV) surveys to detect exoplanets around 36 nearby and young M dwarfs. We use the CSHELL spectrograph ($R \sim 46,000$) at the NASA InfraRed Telescope Facility (IRTF), combined with an isotopic methane absorption gas cell for common optical path relative wavelength calibration. We have developed a sophisticated RV forward modeling code that accounts for fringing and other instrumental artifacts present in the spectra. With a spectral grasp of only 5 nm, we are able to reach long-term radial velocity dispersions of $\sim 20\text{--}30 \text{ m s}^{-1}$ on our survey targets.

Keywords. instrumentation: spectrographs, techniques: radial velocities

1. Introduction

We are carrying out an exoplanet search to exploit the “M dwarf opportunity.” Approximately 70% of main sequence stars are M dwarfs, spanning a factor of 5 in mass/radius and $\sim 10^3$ in luminosity. Due to their smaller masses and radii and cooler temperatures compared to FGK stars, transiting exoplanets produced deeper eclipses of M dwarfs, exoplanet habitable zones are closer, and exoplanet radial velocity amplitude reflex motions are larger. More recently, a new “M dwarf opportunity” for finding exoplanets has been identified with the Kepler mission: not only are planets smaller than $4 R_{\oplus}$ more common than Jovian planets, they are predominantly found around lower mass stars (Howard *et al.* 2012, Dressing & Charbonneau 2013).

There are several factors spoiling the “M dwarf opportunity.” First, M dwarfs are red and faint, with $V - K > 3.5$ mag and only 4 M4 or later dwarfs with $V < 12$ mag. Additionally, small planetary RV signals are comparable in amplitude and time-scale to numerous periodic perturbing stellar jitter signals such as rotational spot modulation & magnetic activity cycles (Vanderberg *et al.* 2015, Robertson *et al.* 2015). One solution to both of these challenges is to observe in the NIR, where M dwarfs put out much of their bolometric luminosity, and the amplitude of stellar jitter is diminished (Reiners *et al.* 2010, Plavchan *et al.* 2013a,b,2015, Anglada-Escude *et al.* 2012).




Figure 1. Left: CSHELL NIR RV time-series for GJ 338A ($N_{obs}=6$, $rms=22 \text{ m s}^{-1}$, $\chi_r^2 = 2.4$). Right: CSHELL NIR RV survey results, plotted as a function of RV rms and reduced chi-squared for both young and nearby M dwarfs in red and blue respectively.




Figure 2. A comparison of the potential exoplanet sensitivity of CSHELL and iSHELL. The solid black lines correspond to the RV semi-amplitude of Habitable Zone (HZ) exoplanets with the indicated masses as a function of stellar mass.

2. Results and Conclusions

We have obtained our goal precision of $\sim 30\text{--}60 \text{ m s}^{-1}$ in our young M dwarf survey from 2010–2012, and $\sim 20\text{--}30 \text{ m s}^{-1}$ in 2014 at higher SNR for nearby M dwarfs not previously surveyed by visible RV teams. We present a sample RV curve and our survey results in Figure 1. We are in the process of publishing our detailed survey results.

Looking towards the future, we have made the gas cells for iSHELL ($R \sim 75,000$, spectral grasp of 250 nm), which will replace CSHELL in 2016. We expect to get to $< 5 \text{ m s}^{-1}$ precision with iSHELL due to the increased spectral grasp alone (Figure 2). The increased efficiency, resolution, no fringing, and smaller number of bad pixels will provide further improvements.

References

Anglada-Escude, G., *et al.*, 2012, *PASP*, 124, 586
 Dressing, C. & Charbonneau, D., 2013, *ApJ*, 767, 95
 Howard, A., *et al.* 2012, *ApJS*, 201, 15
 Reiners, A., *et al.*, 2010, *ApJ*, 710, 432
 Plavchan, P., *et al.*, 2015, *NASA ExoPAG report*, arXiv:1503.01770
 Plavchan, P., *et al.*, 2013a, *SPIE*, 8864, 0G
 Plavchan, P., *et al.*, 2013b, *SPIE* 8864, 1J
 Robertson, P., *et al.* 2015, *ApJ* 801, 79
 Vanderberg, A., *et al.*, 2015, in prep.