



original papers

Psychiatric Bulletin (2008), 32, 288–291. doi: 10.1192/pb.bp.107.018424

NIRUJ AGRAWAL, SIMON FLEMINGER, HOWARD RING AND SHOUMITRO DEB

Neuropsychiatry in the UK: national survey of existing service provision[†]

AIMS AND METHOD

To elucidate and describe current neuropsychiatry service provision in the UK. A questionnaire was developed and posted to members of the Royal College of Psychiatrists who had expressed an interest in neuropsychiatry. The responses were tabulated and analysed using

descriptive statistics and SPSS version 11.0 for Windows. The neuropsychiatry services provided, sources of referrals, setting of the services and funding streams are described.

RESULTS

Out of 251 respondents, 70 reported providing a neuropsychiatry service,

21 having been principally appointed as neuropsychiatrists.

CLINICAL IMPLICATIONS

Neuropsychiatry services in the UK are currently based in a few regional centres, representing a patchy and inadequate service provision.

Increasing recognition of the effect of brain disorder on mental health and insights into the indelible inseparability of mind and body (Yudofsky & Hales, 2002) have contributed to a growing interest in neuropsychiatry. This interest has been fuelled by the arbitrariness of the historical separation of neurology and psychiatry, and recognition of the growing overlap between the two, with advancement of scientific knowledge. In the UK, early identification and management of cognitive, behavioural and mood symptoms in people with neurological disorders, in the true Meyerian tradition of psychobiology (Lidz, 1966), is now an important quality requirement of the National Service Framework for long-term conditions (Agrawal & Mitchell, 2005; Department of Health, 2005).

Yet, the current map of neuropsychiatry service provision in the UK is still not clear. Traditionally, such services have been provided at a few regional or national centres and users were expected to travel long distances for a neuropsychiatrist's opinion. This is very much against the ideals of integrated, interdisciplinary and easily accessible neuropsychiatry services (Department of Health, 2005). Geographic distance to accessible neuropsychiatry services has been found to be associated with unmet need (Fleminger *et al*, 2006). This highlights the need for reasonably local neuropsychiatry services with clear referral pathways.

Nationally, there is a dearth of information about neuropsychiatry services, their referral pathways and funding streams. There have been a few papers on individual neuropsychiatry services or a group of services in the UK (Leonard *et al*, 2002; Barrett & Sudharsan, 2005; Fleminger *et al*, 2006), but they do not provide a coherent national picture.

Hence, the Royal College of Psychiatrists' Special Interest Group in Neuropsychiatry decided to survey College members who had expressed an interest in neuropsychiatry, to obtain information on current neuropsychiatry service provision, including staffing levels, services provided, their setting, referral pathways and funding streams. Our aim was to use this information to inform neuropsychiatry service development in the UK.

Method

A questionnaire was developed by a working group of the Special Interest Group in Neuropsychiatry. This was posted to 508 College members by the audit department of South West London and St George's Mental Health Trust on behalf of the Group. An envelope was included for the reply and an additional sheet of paper was enclosed to allow the trainees, who may have felt they were not responsible for providing any services, to respond. The members were mailed after 4 weeks to encourage response from those who did not answer the initial letter. All the responses were collated and entered on to an SPSS database and analysed with the help of the Trust's audit department.

Results

A total of 251 members (49.4%) replied to the survey: 161 (64.1%) after the first mailing and a further 90 (35.9%) following second mailing. Some trainees who were not responsible for providing a neuropsychiatry

[†]See special article pp. 303–306, this issue.



service answered by telephone, as the questionnaire did not apply to them.

Of the 251 respondents, 70 reported (27.9%) that they were responsible for providing some degree of neuropsychiatry service. The rest ($n=181$), who felt that the survey did not apply to them, included trainees (35.1%), members who had an interest in but were not responsible for providing any neuropsychiatry service (30.4%), and others (32.1%). This latter group included 'addressee not known or moved' or questionnaire returns without stating a reason. Only a small proportion of the respondents (2.6%) reported no interest in neuropsychiatry.

In our analysis we focused on 70 respondents who reported being responsible for or providing some neuropsychiatry services. The vast majority of these (77%) were National Health Service (NHS) consultants, a small number were academics (16%) and a few (7%) worked in the private sector (Fig. 1).

Only 21 of these respondents (30%) were principally employed as neuropsychiatrists. The rest had a range of principal employments in various psychiatric subspecialties (Fig. 2). However, the great majority of these nevertheless provided at least 1–2 sessions worth of neuropsychiatry input.

Of those who were principally employed as a neuropsychiatrist ($n=21$), similar proportions were working in the NHS sector and had academic jobs as in Fig. 1. Over half (52.4%) of the neuropsychiatrists were working full-time in clinical neuropsychiatry – one in five did 6–9 sessions and about a quarter did 3–5 sessions. Part-time neuropsychiatrists were either doing split clinical jobs with other psychiatric subspecialties (30%) or academic jobs (40%), with the rest not specifying.

A wide range of neuropsychiatry services (defined as those employing a neuropsychiatrist) were offered (Table 1). The most common of these were out-patient services and specialist clinics. Over half (61.9%) had in-patient beds. Community nursing was only rarely available as a part of neuropsychiatry service. Most of the services covered a wide range of neuropsychiatry areas. The most common of these were brain injury, memory and epilepsy clinics (Table 2).

Of the 21 doctors who were principally appointed as neuropsychiatrists, over 70% of the neuropsychiatry services had existed for more than 10 years, 5% for more than 5 but less than 10 years and the rest had been developed within the last 5 years. Although 52% of the services had expanded in the last 5 years (including the ones which were developed in this period), 38% remained unchanged and 10% were forced to reduce in size.

Over half of the services were principally based in teaching hospitals (56%); the rest were based in brain injury rehabilitation units (14%), regional neurosciences centres (10%), district general hospitals (10%), and other settings (10%). The sources of funding of the services varied widely (Fig. 3). Three-fourths of the funds was, however, channelled through mental health trusts.

The majority of neuropsychiatry services accepted referrals from a wide range of sources (Table 3), including




Fig. 1. Type of employment ($n=70$).




Fig. 2. Clinical field of principal employment ($n=70$).

Table 1. Neuropsychiatry services provided by the 21 neuropsychiatrists¹

Type of services	<i>n</i> (%)
Special interest clinic	12 (57.1)
General hospital liaison psychiatry	10 (47.6)
Specialist neuropsychiatry service	18 (85.7)
Out-patient service	19 (90.5)
Neuroscience liaison psychiatry	13 (61.9)
Day patients	7 (33.3)
Community nursing	5 (23.8)
In-patient beds	13 (61.9)

1. Doctors were providing more than one service.

psychiatrists, neuroscience clinicians (neurologists, neurosurgeons, neuropsychologists, etc.) and specialist units such as brain injury rehabilitation centres. Despite the perceived tertiary nature of neuropsychiatry, three-quarters of the services accepted referrals from primary care.

Discussion

This is the first national survey of its kind. It focused on the members of the Royal College of Psychiatrists with interest in neuropsychiatry. Although it is possible that some psychiatrists who provide neuropsychiatry services may not have registered an interest, for the purpose of



original papers

Table 2. Areas covered by neuropsychiatric services provided by the 21 neuropsychiatrists¹

Type of conditions	n (%)
Brain injury	20 (95.2)
Epilepsy	19 (90.5)
Movement disorder	13 (61.9)
Cognitive and memory disorder	17 (81.0)
Sleep disorder	12 (57.1)
Liaison psychiatry	15 (71.4)
Neuropsychiatry of developmental disorder	15 (71.4)
Neurological unexplained symptoms	13 (61.9)
Other	3 (14.3)

1. Doctors were providing more than one service.




Fig. 3. Sources of funding (n=21 neuropsychiatrists).

the study we assumed it is unlikely that clinicians principally employed as neuropsychiatrists will not have done so. The rate of response to our survey was consistent with other postal surveys. A high number of trainees and other clinicians expressing interest in neuropsychiatry reflects the growing interest in this specialty. This needs to be fostered and a clear neuropsychiatry training programme has to be provided to meet their needs (Mitchell & Agrawal, 2005).

We found that there were 21 consultants who were principally employed as a neuropsychiatrist in the UK. This number is consistent with the number of neuropsychiatrists known to our interest group. The majority were based in a few major regional, national or teaching hospitals. This confirms the patchy nature of current neuropsychiatry service provision.

A range of psychiatric consultants currently provide some degree of neuropsychiatric input, mostly in the form of 1–2 special interest clinics a week. Out of the 21 consultants who were employed as neuropsychiatrists, about half were devoting all their sessions to clinical neuropsychiatry and most of the rest devoted 3–9

Table 3. Referral pathways (N=21 neuropsychiatrists)

Sources of referrals	n (%)
Neuroscience clinicians	18 (85.7)
General practitioners	16 (76.2)
Psychiatrists	19 (90.5)
Other general hospital consultants	17 (81.0)
Specialist units	12 (57.1)

sessions to clinical work. They provided a wide range of neuropsychiatry services, including a number of specialist neuropsychiatry clinics. Referrals were accepted from primary, secondary or tertiary services. Hence, where the neuropsychiatry services existed, they provided a reasonably comprehensive service. However, in-patient neuropsychiatry beds were not commonly available.

One of the most worrying aspects identified was that of the two-thirds of neuropsychiatry services that had existed for more than a decade, a significant proportion had not expanded in recent years and a significant number was forced to reduce in size. This occurred at a time when neurological and psychiatric services went through an unprecedented expansion and the overall numbers of consultants increased in the UK by over 70% (Department of Health, 2004). This could possibly be attributed to the predominant focus of national service framework for mental health (Department of Health, 1999) on providing comprehensive community services. The wide range of funding sources of existing neuropsychiatry services indicate a lack of coherent funding and commissioning arrangements. This can again be a factor contributing to the lack of appropriate neuropsychiatry service development in the UK.

In conclusion, neuropsychiatry services in the UK are currently based in a few regional centres. This represents a grossly inadequate service provision. Although some other psychiatric specialists try and fill in the gap with the help of special interest clinics, this can not be a reliable way to meet the population need. Neuropsychiatry service development seems to have lagged behind other psychiatric and neuroscience services significantly over the past decade. Lack of clarity of funding streams, commissioning arrangements, and appropriate guidelines about what would constitute an adequate neuropsychiatry service could be contributing to this limited and currently inequitable service provision.

Acknowledgements

We thank the Audit Department of the South West London and St George’s Mental Health Trust at Springfield Hospital which provided substantial help and support. In particular, we thank Ms Liz Collins, Ms Jane Dunnell and Ms Grace Chew.

Declaration of interest

S.F. is the lead consultant neuropsychiatrist for two brain injury units which both require primary care trusts to authorise in-patient admissions and out-patient appointments.

References

AGRAWAL, N. & MITCHELL, A. J. (2005) The national service framework for long term conditions: aims to meet psychological, social and emotional needs in neurological disorders. *BMJ*, **330**, 1280–1281.

BARRETT, K. & SUDHARSAN, S. (2005) Service innovations: is there a market for neuropsychiatry? A year in the life of a district based neuropsychiatry service. *Psychiatric Bulletin*, **29**, 465–467.



DEPARTMENT OF HEALTH (1999) *Modern Standards and Service Models. National Service Framework for Mental Health* (http://www.hcsu.org.uk/index.php?option=com_content&task=view&id=243&Itemid=109).

DEPARTMENT OF HEALTH (2005) *National Service Framework for Long-term Conditions* (http://www.dh.gov.uk/en/publicationsandstatistics/publications/publicationspolicyandguidance/dh_4105361).

FLEMINGER, S., LEIGH, E. & MCCARTHY, C. (2006) The size of demand for specialised neuropsychiatry services: rates of referrals to neuropsychiatric services in South Thames region of the United Kingdom. *Journal of Neuropsychiatry and Clinical Neurosciences*, **18**, 121–128.

LEONARD, F., MAJID, S., SIVAKUMAR, K., et al (2002) Service innovations: a neuropsychiatry outreach clinic. *Psychiatric Bulletin*, **26**, 99–101.

LIDZ, T. (1966) Adolf Meyer and development of American psychiatry. *American Journal of Psychiatry*, **159**, 695–704.

MITCHELL, A. J. & AGRAWAL, N. (2005) Training in neuropsychiatry: is it time to reintegrate into mainstream

psychiatry? *Psychiatric Bulletin*, **29**, 361–364.

YUDOFSKY, S. C. & HALES, R. E. (2002) Neuropsychiatry and future of psychiatry and Neurology. *American Journal of Psychiatry*, **159**, 1261–1264.

***Niruj Agrawal** Consultant in Neuropsychiatry, Department of Neuropsychiatry, Clare House, St George's Hospital, London SW17 0QT, email: Niruj.Agrawal@swlstg-tr.nhs.uk, **Simon Fleminger** Consultant in Neuropsychiatry, **Howard Ring** Consultant in Developmental Neuropsychiatry, **Shoumitro Deb** Clinical Professor of Neuropsychiatry and Intellectual Disability

Psychiatric Bulletin (2008), **32**, 291–294. doi: 10.1192/pb.bp.107.017715.

LUKE SOLOMONS, ADRIAN TRELOAR AND RYAN NORONHA

Competence of psychiatric clinicians in interpreting electrocardiograms and QT intervals: can they do this? Does it matter?

AIMS AND METHOD

We assessed the abilities of trainee and consultant psychiatrists in reading and interpreting electrocardiograms (ECGs) and QT intervals using a questionnaire and standardised ECG.

RESULTS

Only 5% of our sample of trainee

and consultant psychiatrists could correctly indicate a QTc interval. Performances on other measures, such as rate were also poor, with senior house officers performing better than consultants.

CLINICAL IMPLICATIONS

The increased awareness of problems caused by antipsychotics has not

been reflected in improved knowledge of ECGs among psychiatrists. Machines do not reliably calculate QT intervals. We therefore urge better training and understanding of ECGs for psychiatrists.

People with mental illnesses, especially schizophrenia, have a higher mortality rate than the general population, the causes of which are yet poorly understood but could be related to side-effects of antipsychotic medication (Waddington et al, 1998). Psychotropic medication has been shown to increase the risk of serious ventricular arrhythmias and thus, sudden cardiac death. Ray et al (2001) and Liperoti et al (2005) showed that moderate doses of antipsychotics were associated with large absolute and relative increases in the risk of sudden cardiac death. Prolonged QT intervals reflect electrical dysfunction of the myocardium and may indicate predisposition to malignant tachycardias and sudden death. The QT interval is prolonged by a slow pulse and being female, and shortened by anxiety. Importantly, drugs and ischaemia can also prolong it. The corrected QT (QTc) gives a better indication of the risks by allowing for the pulse rate. There are various methods of calculating this, but the most commonly used is the Bazett's formula (Box 1) because of its relative simplicity, despite inaccuracies associated with heart rate. Failure to identify a prolonged QT leaves patients at risk of sudden death.

Recent work by Minerotti et al (2006) has shown increased QTc intervals in healthy men with neurotic personality traits, in alcohol dependence syndrome (Borini et al, 2003), and in incident diabetics (Carnethon

et al, 2006). The new consensus statement on high-dose antipsychotic medication from the Royal College of Psychiatrists (2006) recommends assessing cardiovascular risk with electrocardiogram (ECG) monitoring prior to use of antipsychotic medication to reduce risk of arrhythmias – we presume this has increased awareness and ordering of ECGs. Proper monitoring of QTc interval may detect prolongation and reduce occurrence of potentially lethal cardiac arrhythmias.

In this context, doctors have come to be more dependent upon computer ECG interpretation. The reliability of such programs has not been proven (Willems et al, 1991). Standards regarding the accuracy of these readings were developed in the 1980s, but they have been largely ignored. Recent work in this area (Krishnan et al, 2006) has highlighted their deficiencies. The British Heart Foundation (2005) advises that computer-assisted ECG interpretation can identify important anomalies, but errors are common and interpretation should not be accepted without visual inspection. Electrocardiogram departments and cardiologists lack the capacity to provide a QTc validation service. We thus feel that psychiatrists should be competent in this skill – the Royal College of Psychiatrists recommended this in a consensus statement on high-dose antipsychotic medication (Royal College of Psychiatrists, 2006).