

FUNCTION-THEORETIC METRICS AND BOUNDARY BEHAVIOUR OF FUNCTIONS MEROMORPHIC OR HOLOMORPHIC IN THE UNIT DISK

SHINJI YAMASHITA

§ 1. Introduction. The metrics to which the title of the present paper refers are expressed in the form of elements of arc length as follows:

- (i) $|dw|$ in the finite w -plane $W_1 : |w| < \infty$.
- (ii) $\frac{|dw|}{1 + |w|^2}$ in the Riemann w -sphere $W_2 : |w| \leq \infty$.
- (iii) $\frac{|dw|}{1 - |w|^2}$ in the open unit disk $W_3 : |w| < 1$.

Let $D : |z| < 1$ be the open unit disk and let $\Gamma : |z| = 1$ be the unit circle in the z -plane. We fix a constant ρ , $1/2 < \rho < 1$, once and for all and we denote by $\mathcal{D}(\zeta)$ the open disk $\{z; |z - \rho\zeta| < 1 - \rho\}$ for $\zeta \in \Gamma$. By a segment X at $\zeta \in \Gamma$ we mean an open rectilinear segment connecting ζ and a point of D . Let $w = f(z)$ be a function from D into W_j ($j = 1, 2, 3$), being meromorphic or holomorphic in D , and set for $z = re^{i\theta} \in D$,

$$\begin{aligned}\delta_1(r, \theta) &= |f'(re^{i\theta})|; \\ \delta_2(r, \theta) &= \frac{|f'(re^{i\theta})|}{1 + |f(re^{i\theta})|^2}; \\ \delta_3(r, \theta) &= \frac{|f'(re^{i\theta})|}{1 - |f(re^{i\theta})|^2};\end{aligned}$$

corresponding respectively to $j = 1, 2$ and 3 . The word ‘‘capacity’’ always means ‘‘logarithmic capacity’’. Then our result is stated in the following

THEOREM. *Let M be a subset of Γ which is a Borel set in the plane and set*

$$\sigma = \bigcup_{\zeta \in M} \mathcal{D}(\zeta).$$

Received November 27, 1970.

Let $w = f(z)$ be a meromorphic or holomorphic function from D into W_j such that

$$(1) \quad \iint_D \{\delta_j(r, \theta)\}^2 r dr d\theta < \infty \quad (j = 1, 2, 3).$$

Then there exists a subset E_j of M , being of capacity zero^{*}), such that for any $\zeta \in M - E_j$ and for any segment X at ζ we have

$$(2) \quad \int_X \delta_j(r, \theta) |dz| < \infty \quad (z = re^{i\theta} \in X)$$

according as $j = 1, 2, 3$.

The condition (2) for $j = 1, 2, 3$ implies the existence of a limiting value $f(\zeta) \in W_j$ of $f(z)$ as $X \ni z \rightarrow \zeta$ according as $j = 1, 2, 3$. Then by the theorem of Lindelöf-Iversen-Gross [1, p. 5] combined with our condition (1), the function f has the angular limit $f(\zeta)$ at ζ , in other words, ζ is a Fatou point [1, p. 59] of f . It should therefore be noted that our theorem in the case $j = 1, 2$ gives "localization" of Beurling-Tsuji's theorem ([3, Theorems 3 and 4], [4, p. 344]).

An application of the theorem for $j = 3$ is the following. Let $G \subset W_3$ be a Jordan domain whose non-Euclidean area is finite and let $w = \Phi(z)$ be a one-to-one conformal map from D onto G in the w -plane. Furthermore, let $\Phi(\zeta)$ be the Carathéodory extension of Φ to Γ . Then we have $|\Phi(\zeta)| < 1$ except perhaps for a set of $\zeta \in \Gamma$ of capacity zero. Therefore, the boundary of G touches the circle $|w| = 1$ at a "thin" set in this sense.

§ 2. Three lemmas. Let $0 < \alpha < \pi/2$ and let $\Delta = \{re^{i\theta}; 0 < r \leq 1, |\theta| \leq \alpha\}$. We let $\Delta^* \supset \Delta$ be an open disc whose boundary contains the origin and we use the same notation $\delta_j(r, \theta)$ as in § 1 for a function f defined in Δ^* ($j = 1, 2, 3$). We begin with two lemmas [4, p. 342, Theorem VIII. 47 and p. 343, Theorem VIII. 48] expressed in one.

LEMMA $j(j = 1, 2)$. Let $w = f(z)$ be a function from Δ^* into W_j , being meromorphic or holomorphic in Δ^* . Assume that f does not take three distinct points of W_2 in Δ^* and set

$$A_j(\theta) = \int_0^1 \delta_j(r, \theta) dr$$

for $|\theta| \leq \alpha$. Assume furthermore that both $A_j(-\alpha)$ and $A_j(\alpha)$ are finite. Then $A_j(\theta)$ is bounded for $|\theta| \leq \alpha$.

^{*}) In other words, the outer logarithmic capacity of E_j is zero.

The following lemma needs a proof.

LEMMA 3. Let $w = f(z)$ be a holomorphic function from Δ^* into W_3 . Set

$$A_3(\theta) = \int_0^1 \delta_3(r, \theta) dr$$

for $|\theta| \leq \alpha$ and assume that both $A_3(-\alpha)$ and $A_3(\alpha)$ are finite. Then $A_3(\theta)$ is bounded for $|\theta| \leq \alpha$.

Proof. As f is bounded in Δ^* , by the same argument as in the next paragraph to the theorem in §1 the origin is a Fatou point of f at which f has the angular limit $f(0)$ with $|f(0)| < 1$. This implies that we have a positive constant B such that $(1 - |f(re^{i\theta})|^2)^{-1} < B$ on Δ . On the other hand, both $A_1(-\alpha)$ and $A_1(\alpha)$ are finite because of $\delta_3(r, \theta) \geq \delta_1(r, \theta)$ for $|\theta| \leq \alpha$. Lemma 3 follows from Lemma 1 combined with $A_3(\theta) \leq BA_1(\theta)$ for $|\theta| \leq \alpha$.

§ 3. Proof of Theorem. In the following $z = re^{i\theta}$ and $e^{i\omega}$ are always points of D and M respectively. To avoid unnecessary complexity we drop the suffix j of $\delta_j(r, \theta)$ if the argument is true for $j = 1, 2, 3$. We remark that $\delta_2(r, \theta)$ is not defined at the poles of f ; but this is not essential in the following proof.

We set

$$h(r, \theta) = \begin{cases} \delta(r, \theta) & \text{for } z \in \sigma, \\ 0 & \text{for } z \in D - \sigma. \end{cases}$$

Let $\phi \equiv \phi(r, \theta) = \pi - \arg(re^{i\theta} - 1)$, where $0 < r < 1$, $|\theta| \leq \pi$ and $\pi/2 < \arg(re^{i\theta} - 1) < 3\pi/2$. Then by $\tan \phi = r \sin \theta / (1 - r \cos \theta)$ we have

$$\begin{aligned} (3) \quad \frac{\partial \phi}{\partial \theta} &= - \frac{\partial}{\partial \theta} \arg(re^{i\theta} - 1) \\ &= \frac{\partial}{\partial \theta} \operatorname{Im} \log \{1/(re^{i\theta} - 1)\} \\ &= r(\cos \theta - r)/(1 - 2r \cos \theta + r^2). \end{aligned}$$

We next consider the function

$$(4) \quad H(\omega; r, \theta) = h(r, \theta + \omega) \frac{\partial \phi}{\partial \theta}.$$

Then $H(\omega; r, \theta)$, for a fixed ω , is Lebesgue measurable for $0 < r < 1$ and $|\theta| \leq \pi$; and $H(\omega; r, \theta) \geq 0$ in the disk

$$S = \{re^{i\theta}; \cos \theta > r\}$$

and further $H(\omega; r, \theta) \leq 0$ in $D - S$ by (3). Therefore we may consider two integrals:

$$J_1(\omega) = \iint_S H(\omega; r, \theta) dr d\theta \geq 0$$

and

$$J_2(\omega) = - \iint_{D-S} H(\omega; r, \theta) dr d\theta \geq 0$$

for $e^{i\omega} \in M$. We first assert that

(I) $J_2(\omega) < +\infty$ for any $e^{i\omega} \in M$, so that $H(\omega; r, \theta)$ possesses a definite integral on D [2, p. 20] and that

$$(5) \quad J(\omega) \equiv \iint_D H(\omega; r, \theta) dr d\theta = J_1(\omega) - J_2(\omega).$$

We let, for the proof, C_r be the circle $|z| = r$, $0 < r < 1$. Then

$$- \frac{\partial \phi}{\partial \theta} = r(r - \cos \theta)/(1 - 2r \cos \theta + r^2) \leq r/(r + 1) < r$$

for $re^{i\theta} \in C_r - S$. This can be proved by considering $-\frac{\partial \phi}{\partial \theta}$ as a function of $\cos \theta$ (cf. [4, p. 346]). Therefore by (3) and (4) we have

$$(6) \quad -H(\omega; r, \theta) \leq rh(r, \theta + \omega), \quad re^{i\theta} \in C_r - S.$$

We estimate $J_2(\omega)$ upwards by (6) and by Schwarz's inequality as follows:

$$\begin{aligned} J_2(\omega) &= - \int_0^1 dr \int_{C_r-S} H(\omega; r, \theta) d\theta \leq \int_0^1 dr \int_{C_r-S} rh(r, \theta + \omega) d\theta \\ &= \iint_{D-S} h(r, \theta + \omega) r dr d\theta \leq \iint_D h(r, \theta + \omega) r dr d\theta \\ &= \iint_D h(r, \theta) r dr d\theta \leq \pi^{1/2} \left[\iint_D \{h(r, \theta)\}^2 r dr d\theta \right]^{1/2} \\ &= (\pi U)^{1/2} < +\infty, \end{aligned}$$

where

$$(7) \quad U = \iint_D \{h(r, \theta)\}^2 r dr d\theta = \iint_D \{\delta(r, \theta)\}^2 r dr d\theta < +\infty$$

by our assumption (1) in the theorem. This completes the proof of (I).

Let $\mathcal{L}(\omega, \varphi)$ be the chord of the circle $|z - \rho e^{i\omega}| = 1 - \rho$, with one end-point $e^{i\omega}$, making the directed angle φ , $|\varphi| < \pi/2$, with the radius of D at $e^{i\omega}$. We shall use the notation $\mathcal{L}(0, \varphi)$ though $\zeta = 1$ may not be in M . The chord $\mathcal{L}(\omega, \varphi)$ has the length

$$(8) \quad \lambda(\varphi) = (2 - 2\rho) \cos \varphi,$$

being independent of ω . We then set for $-\pi/2 < \varphi < \pi/2$,

$$(9) \quad L(\omega, \varphi) = \int_{\mathcal{L}(\omega, \varphi)} \delta_2(r, \theta) |dz| \quad (z = r e^{i\theta} \in \mathcal{L}(\omega, \varphi))$$

and we consider the function $\chi(\omega)$ on M defined by

$$(10) \quad \chi(\omega) = \int_{-\pi/2}^{\pi/2} L(\omega, \varphi) \cos \varphi d\varphi.$$

(II) *The function $\chi(\omega)$ is Borel measurable on M .*

We shall prove this for $\delta_2(r, \theta)^*$. In other cases the proofs are simpler and hence are omitted.

Let γ_k ($k = 1, 2, \dots$) be the circle $|z| = r_k$, $2\rho - 1 \leq r_k < 1$, such that $r_k \nearrow 1$ and the set $\bigcup_{k=1}^{\infty} \gamma_k$ contains all the poles of f in the half-open ring $\{z; 2\rho - 1 \leq |z| < 1\}$. Let R_ν ($\nu = 1, 2, \dots$) be the open set, being o the form of a summation of ring domains whose boundaries are concentric circles with the centre $z = 0$, such that

$$R_1 \supset R_2 \supset \dots \supset \bigcap_{\nu=1}^{\infty} R_\nu = \bigcup_{k=1}^{\infty} \gamma_k.$$

Let $2\rho - 1 < \beta_1 < \dots < \beta_m < \dots < 1$, $\beta_m \nearrow 1$ and let D_m be the closed ring $\{z; 2\rho - 1 \leq |z| \leq \beta_m\}$. We then set $D_{m\nu} = D_m - R_\nu$ for $m, \nu = 1, 2, \dots$. We note first that

$$(11) \quad L(\omega, \varphi) = \int_{\mathcal{L}(\omega, \varphi)} \delta_2(r, \theta) |dz| = \int_{\mathcal{L}(0, \varphi)} \delta_2(r, \theta + \omega) |dz|$$

($z = r e^{i\theta} \in \mathcal{L}(0, \varphi)$ in the last expression)

and we then consider

$$L_{m\nu}(\omega, \varphi) \equiv \int_{\mathcal{L}(0, \varphi) \cap D_{m\nu}} \delta_2(r, \theta + \omega) |dz|$$

($z = r e^{i\theta} \in \mathcal{L}(0, \varphi) \cap D_{m\nu}$).

* δ_2 may be extended continuously to the poles of f and our proof will be rather simplified (*Added in proof*).

We shall show that for any $e^{i\omega_0} \in M$ we have $L_{m\nu}(\omega, \varphi) \rightarrow L_{m\nu}(\omega_0, \varphi)$ as $\omega \rightarrow \omega_0$ uniformly for $-\pi/2 < \varphi < \pi/2$, so that

$$\chi_{m\nu}(\omega) \equiv \int_{-\pi/2}^{\pi/2} L_{m\nu}(\omega, \varphi) \cos \varphi d\varphi$$

is continuous on M . Indeed,

$$\begin{aligned} & |L_{m\nu}(\omega, \varphi) - L_{m\nu}(\omega_0, \varphi)| \\ & \leq \int_{\mathcal{L}(0, \varphi) \cap D_{m\nu}} |\delta_2(r, \theta + \omega) - \delta_2(r, \theta + \omega_0)| |dz| \\ & \leq \{ \max_{re^{i\theta} \in D_{m\nu}} |\delta_2(r, \theta + \omega) - \delta_2(r, \theta + \omega_0)| \} \times \\ & \times \{ \sup_{|\varphi| < \pi/2} \int_{\mathcal{L}(0, \varphi) \cap D_{m\nu}} |dz| \}, \end{aligned}$$

so that our assertion follows from the uniform continuity of the function $\delta_2(r, \theta)$ on the compact set $D_{m\nu}$. Set

$$L_m(\omega, \varphi) = \int_{\mathcal{L}(0, \varphi) \cap D_m} \delta_2(r, \theta + \omega) |dz|$$

and further set

$$\chi_m(\omega) = \int_{-\pi/2}^{\pi/2} L_m(\omega, \varphi) \cos \varphi d\varphi.$$

Then $\chi_{m\nu}(\omega) \nearrow \chi_m(\omega)$ as $\nu \nearrow \infty$ and $\chi_m(\omega) \nearrow \chi(\omega)$ as $m \nearrow \infty$. This proves our proposition (II).

(III) *The inequality $J_1(\omega) \geq (2\rho - 1)\chi(\omega)$ holds for any $e^{i\omega} \in M$.*

We remember that $\mathcal{D}(1)$ is the disk $|z - \rho| < 1 - \rho$ and we let

$$J_1^*(\omega) = \iint_{\mathcal{D}(1)} H(\omega; r, \theta) dr d\theta.$$

Then $J_1(\omega) \geq J_1^*(\omega)$ since $S \supset \mathcal{D}(1)$ and $H(\omega; r, \theta) \geq 0$ in S . To estimate $J_1^*(\omega)$ downwards, we set for $re^{i\theta} \in \mathcal{D}(1)$,

$$\begin{aligned} t &= |re^{i\theta} - 1| \text{ and } \psi = \pi - \arg(re^{i\theta} - 1) \text{ for} \\ \pi/2 &< \arg(re^{i\theta} - 1) < 3\pi/2. \end{aligned}$$

Then $1 > r = (1 - 2t \cos \psi + t^2)^{1/2}$, and on the chord $\mathcal{L}(0, \psi)$, for a fixed ψ , $|\psi| < \pi/2$, we have

$$\begin{aligned} dr &= (t - \cos \psi)(1 - 2t \cos \psi + t^2)^{-1/2} dt \\ &\geq (\cos \psi - t)(-dt) \text{ (for } dt \leq 0). \end{aligned}$$

We note that r decreases as t increases on $\mathcal{L}(0, \psi)$ and $\cos \psi \geq t$ since $re^{i\theta} \in \mathcal{D}(1) \subset S$. Furthermore, on the circle $C_r : |z| = r, 0 < r < 1$, we have

$$H(\omega; r, \theta)d\theta = h(r, \theta + \omega)d\psi$$

by (4). We therefore obtain

$$\begin{aligned} J_1^*(\omega) &= \int_{2\rho-1}^1 dr \int_{C_r \cap \mathcal{D}(1)} H(\omega; r, \theta)d\theta \\ &= \int_{2\rho-1}^1 dr \int_{C_r \cap \mathcal{D}(1)} h(r, \theta + \omega)d\psi \\ &= \iint_{\mathcal{D}(1)} h(r, \theta + \omega)drd\psi \\ &= \int_{-\pi/2}^{\pi/2} d\psi \int_{\mathcal{L}(0, \psi)} h(r, \theta + \omega)dr \\ &\geq \int_{-\pi/2}^{\pi/2} d\psi \int_0^{\lambda(\psi)} \delta(r, \theta + \omega)(\cos \psi - t)dt \end{aligned}$$

(where $\lambda(\psi)$ is defined in (8); we note that $h(r, \theta + \omega) = \delta(r, \theta + \omega)$ for $re^{i\theta} \in \mathcal{D}(1)$ since $\sigma \supset \mathcal{D}(e^{i\omega})$)

$$\geq (2\rho - 1) \int_{-\pi/2}^{\pi/2} d\psi \int_0^{\lambda(\psi)} \delta(r, \theta + \omega) \cos \psi dt$$

(because of $\cos \psi - t \geq (2\rho - 1) \cos \psi$ for $0 \leq t \leq \lambda(\psi)$)

$$= (2\rho - 1) \int_{-\pi/2}^{\pi/2} L(\omega, \psi) \cos \psi d\psi$$

(cf. (11); the formula (11) is true for δ)

$$= (2\rho - 1)\chi(\omega).$$

(IV) *The set $E = \{e^{i\omega} \in M; \chi(\omega) = +\infty\}$ is of capacity zero.*

By (II) the set E is a Borel set in the plane, so that E is capacitable by the celebrated Choquet theorem. Therefore we have only to prove that E is of inner capacity zero. Assume on the contrary that E contains a closed set F of positive capacity and let

$$u(z) = \int_F \log(1/|z - e^{i\omega}|)d\mu(\omega) \leq V < +\infty$$

be the conductor potential [4, p. 55] of F , where V is a constant and μ is a Borel measure on F of total mass $\mu(F) = 1$. Then we have [4, p. 345]

$$(12) \quad \iint_D \left(\frac{\partial u}{\partial r} \right)^2 r dr d\theta \leq \pi V/2$$

and

$$(13) \quad r \frac{\partial u}{\partial r} = - \int_F \frac{\partial}{\partial \theta} \arg (re^{i\theta} - e^{i\omega}) d\mu(\omega).$$

We next consider the function

$$(14) \quad \begin{aligned} Q(\omega; r, \theta) &\equiv H(\omega; r, \theta - \omega) \\ &= -h(r, \theta) \frac{\partial}{\partial \theta} \arg (re^{i\theta} - e^{i\omega}) \\ &= h(r, \theta) r \{ \cos(\theta - \omega) - r \} / \{ 1 - 2r \cos(\theta - \omega) + r^2 \} \end{aligned}$$

for $re^{i\theta} \in D$ and $e^{i\omega} \in F$ (cf. (3), (4)). Then Q is a Borel measurable function on the product space $D \times F$ and by (13) and (14) we have

$$h(r, \theta) r \frac{\partial u}{\partial r} = \int_F Q(\omega; r, \theta) d\mu(\omega).$$

On the other hand, both $h(r, \theta)$ and $\frac{\partial u}{\partial r}$ are square summable on D with respect to the measure $r dr d\theta$ by (7) and (12). Therefore, we have by Schwarz's inequality,

$$\begin{aligned} J &\equiv \iint_D dr d\theta \int_F Q(\omega; r, \theta) d\mu(\omega) \\ &= \iint_D h(r, \theta) r \frac{\partial u}{\partial r} dr d\theta \neq \pm \infty. \end{aligned}$$

By Fubini's theorem [2, p. 87] applied to the positive and the negative parts of Q respectively we have

$$(15) \quad J = \int_F d\mu(\omega) \iint_D Q(\omega; r, \theta) dr d\theta \neq \pm \infty.$$

Now, by (3), (4), (5) and (14) we have

$$\begin{aligned} J(\omega) &= \iint_D h(r, \theta + \omega) \frac{\partial}{\partial \theta} \{- \arg (re^{i\theta} - 1)\} dr d\theta \\ &= \iint_D h(r, \theta) \frac{\partial}{\partial \theta} \{- \arg (re^{i\theta} - e^{i\omega})\} dr d\theta \\ &= \iint_D Q(\omega; r, \theta) dr d\theta, \end{aligned}$$

so that by (15),

$$J = \int_F J(\omega) d\mu(\omega) \neq \pm \infty.$$

However, by (5), (III) and the very definition of E we have $J(\omega) = +\infty$ for $e^{i\omega} \in F \subset E$. This is a contradiction.

(V) *The set E is the exceptional set in the statement of the theorem.*

Let $e^{i\omega} \in M - E$. Then $\chi(\omega) < +\infty$, so that by the definition of $\chi(\omega)$ (cf. (10)), the quantity $L(\omega, \varphi)$ (cf. (9)) is finite for a.e., φ , $|\varphi| < \pi/2$. Consequently, there are two chords $\sphericalangle(\omega, \varphi_1)$ and $\sphericalangle(\omega, \varphi_2)$, $-\pi/2 < \varphi_1 < \varphi_2 < \pi/2$, at $e^{i\omega}$ such that $L(\omega, \varphi_k) < +\infty$, $k = 1, 2$. By Lemma j for $j = 1, 2, 3$ and by our assumption (1) we know that $L(\omega, \varphi) < +\infty$ for any φ , $\varphi_1 < \varphi < \varphi_2$. Repeating this process, we have the required property (2) at the point $e^{i\omega} \in M - E$.

REFERENCES

- [1] K. Noshiro, Cluster sets, Berlin-Göttingen-Heidelberg (1960).
- [2] S. Saks, Theory of the integral, second edition, Warsaw (1937).
- [3] M. Tsuji, Beurling's theorem on exceptional sets, Tôhoku Math. J. 2 (1950), 113-125.
- [4] M. Tsuji, Potential theory in modern function theory, Tokyo (1959).

*Mathematical Institute
Tôhoku University
Sendai 980, Japan*