

Dietary glycaemic index and glycaemic load among Australian children and adolescents

Jimmy Chun Yu Louie¹, Anette E. Buyken², Kristina Heyer³ and Victoria M. Flood^{1,4*}

¹Cluster for Public Health Nutrition, Boden Institute of Obesity, Nutrition, Exercise and Eating Disorders, University of Sydney, NSW 2006, Australia

²Department of Nutrition and Health, Research Institute of Child Nutrition, Dortmund, Germany

³Faculty of Health Sciences, University of Bielefeld, 33501 Bielefeld, Germany

⁴School of Health Sciences, Faculty of Health and Behavioural Sciences, The University of Wollongong, NSW 2522, Australia

(Received 29 October 2010 – Revised 21 February 2011 – Accepted 21 February 2011 – First published online 18 May 2011)

Abstract

There are no published data regarding the overall dietary glycaemic index (GI) and glycaemic load (GL) of Australian children and adolescents. We therefore aim to describe the dietary GI and GL of participants of the 2007 Australian National Children's Nutrition and Physical Activity Survey (2007ANCNPAS), and to identify the main foods contributing to their GL. Children, aged 2–16 years, who provided two 24 h recalls in the 2007ANCNPAS were included. A final dataset of 4184 participants was analysed. GI of each food item was assigned using a previously published method. GL was calculated, and food groups contributing to the GL were described by age group and sex. The weighted mean dietary GI and GL of the participants were 54 (SD 5) and 136 (SD 44), respectively. Among the nutrients examined, Ca had the highest inverse relationship with GI ($P < 0.001$), while percentage energy from starch was most positively associated with GI. The association between fibre density and GI was modest, and percentage energy from sugar had an inverse relationship with GI. Daily dietary GL contributed by energy-dense and/or nutrient-poor (EDNP) items in subjects aged 14–16 years was more than doubled that of subjects aged 2–3 years. To conclude, Australian children and adolescents were having a high-GI dietary pattern characterised by high-starchy food intake and low Ca intake. A significant proportion of their dietary GL was from EDNP foods. Efforts to reduce dietary GI and GL in children and adolescents should focus on energy-dense starchy foods.

Key words: Dietary glycaemic index: Glycaemic index: Glycaemic load: Australian: Children

Chronic high dietary glycaemic index (GI) and glycaemic load (GL) have been shown to increase the risk of chronic diseases such as diabetes and CVD among adults^(1–3). However, the current evidence regarding dietary GI and GL and their associations with disease risks, among children and adolescents, is limited and mixed. Data from longitudinal studies such as the Dortmund Nutritional and Anthropometric Longitudinally Designed (DONALD) study^(4,5) did not show any prospective association between dietary GI or GL and percentage of body fat or BMI. There has also been no study to investigate the effect of dietary GI or GL in childhood on long-term health risks. On the other hand, low dietary GL has been shown to improve cognitive performance^(6,7), and some evidence from randomised controlled trials^(8,9) also suggested that a low-GL diet was beneficial in the treatment of obesity in children and adolescents.

Despite the potential health benefits, data about the dietary GI and GL of a nationally representative sample of children and adolescents have not been reported in the literature, and there have only been a few investigations of GI and GL among children and adolescents in larger samples or national surveys in other countries^(10–12). Therefore, the aims of the present study are to (1) describe the dietary GI and GL of Australian children and adolescents, (2) analyse their relation to other nutrients and (3) identify the major foods contributing to their GI and GL, using data from the most recent Australian national dataset available to date, the 2007 Australian National Children's Nutrition and Physical Activity Survey (2007ANCNPAS)⁽¹³⁾. Furthermore, we examined age group and sex differences with regard to these aims.

Abbreviations: 2007ANCNPAS, 2007 Australian National Children's Nutrition and Physical Activity Survey; CHO, carbohydrate; DONALD, Dortmund Nutritional and Anthropometric Longitudinally Designed; GI, glycaemic index; GL, glycaemic load; PAL, physical activity level.

* **Corresponding author:** Assistant Professor V. M. Flood, fax +61 2 4221 5945, email vflood@uow.edu.au

Materials and methods

The 2007 Australian National Children's Nutrition and Physical Activity Survey

The 2007ANCNPAS was commissioned in 2007 by the Australian Commonwealth Department of Agriculture, Fisheries and Forestry, and the Australian Food and Grocery Council⁽¹³⁾. The methodology of the 2007ANCNPAS was previously described in detail⁽¹⁴⁾. In brief, the survey measured the dietary intakes of food and beverages as well as the use of supplements using the 24 h recall method, administered twice during the survey period. These data were collected on children aged 2–16 years (*n* 4834) between 22 February and 30 August 2007. Dietary intake data were entered into a purpose-built database, with nutrition compositions based on the AUSNUT2007 database⁽¹⁵⁾.

Data cleaning

Children who completed only one 24 h recall (*n* 179) were excluded from the analysis. The plausibility of the remaining food intake data was assessed using the Goldberg cut-off⁽¹⁶⁾ for specific physical activity level (PAL), which was determined from information collected about physical activity with a pedometer. Where PAL data were not available (*n* 2438), we have utilised the lower 95% CI of a PAL of 1.5 (i.e. 0.93) and the upper 95% CI of a PAL of 1.7 (i.e. 2.73) as the cut-off values for under- and over-reporting. These PAL cut-off points were approximately the 25th and 75th percentile of those children who had a PAL determined from the survey. Participants with the energy intake:BMR ratio outside the 95% CI were excluded from the analysis. We excluded 360 under-reporters and 100 over-reporters based on this method. We also excluded eleven participants who reported unusually high intakes of foods (e.g. six cups of rice in a meal). The final dataset included 4184 participants, where 51% were male. The demographic characteristics of the participants are summarised in Table 1.

Linking published glycaemic index values to 2007 Australian National Children's Nutrition and Physical Activity Survey food items

GI values were assigned to individual food items recorded in the 2007ANCNPAS dataset based on a method previously described by our group⁽¹⁷⁾ with a small modification at step 1, as AUSNUT2007 contains no information about the GI. In short, the four steps involved in the assignment process are as follows:

Step 1. Determine if the item has <5 g of carbohydrates (CHO) per 100 g. If yes, a GI value of 0 was assigned to that item. If no;

Step 2. Determine if there is a 'closely related food item' or an exact match in the four databases used. If yes, assign that GI value. If no;

Step 3. Determine if the median GI value of the food subgroup is available. If yes, assign the median GI value of the subgroup. If no;

Table 1. Demographic characteristics of the subjects included in the analyses

(Number of subjects and percentages, *n* 4184)

	<i>n</i>	%	Australian national statistic (%)*
Sex (<i>n</i>)			
Male	2134	51.0	51.3
Female	2050	49.0	48.7
Age groups (years)			
2–3	1068	25.5	13.4
4–8	1181	28.0	28.0
9–13	1019	24.4	36.4
14–16	916	21.9	22.3
Aboriginal			
Yes	120	2.9	4.3
No	4062	97.1	90.1
Missing	2	0.0	5.6
Region			
Urban	2805	67.0	N/A
Rural	1379	33.0	N/A
State			
ACT	166	4.0	1.6
NSW	1037	24.8	33.0
NT	88	2.1	1.0
QLD	698	16.7	19.7
SA	818	19.6	7.6
TAS	172	4.1	2.4
VIC	826	19.7	24.8
WA	379	9.1	9.9

N/A, not available.

* Data from the 2006 Australian Census for children and adolescents aged 2–16 years.

Step 4. Determine if the item is a 'top carbohydrate contributor'⁽¹⁷⁾.

If yes, assign a GI value of 50 or a GI value of an appropriate closest matched item as decided by the research nutritionists. If no, a GI value of 0 is assigned.

Recipes in the original 2007ANCNPAS database were recorded as individual ingredients, and GI values were assigned to the individual ingredients rather than the recipe as an item. It was not envisaged to significantly affect the final mean dietary GI or GL of the participants. This is because the majority of these recipes (*n* 3332) were sandwiches, where the GI of the recipe was not anticipated to be significantly different from the main CHO component(s) in the recipe.

There were a total of 3418 different food items recorded in the 24 h recalls. Items with similar expected glycaemic effects and nutritional properties were grouped together for this analysis. The details of food groupings used in this analysis can be found in Table S1 of the supplementary material (available online at <http://www.journals.cambridge.org/bjn>).

Following the aforementioned method, 1075 (31.5%) food items that contained <5 g of CHO per 100 g were assigned a GI of 0 (step 1), whereas 1829 (53.5%) items were assigned the GI value of a 'closely related' food item or an exact match in the GI databases used (step 2). A total of 239 (7.0%) items were assigned the median GI of their corresponding food subgroups (step 3), while 275 (8.0%) items were assigned a GI of 0 because they were not the 'top carbohydrate contributors', and ten items in the 'top carbohydrate contributors' list (0.3%) were assigned a GI of 50 (step 4).

Calculation of the glycaemic load and dietary glycaemic index

The GL of each food item was calculated as the corresponding GI (%) × amount (g) of available CHO in a serve of that food. The daily dietary GL of each subject was calculated as Σ GL, and the dietary GI was obtained by (dietary GL/total available CHO intake in the day) × 100%.

Statistical analysis

In order to increase the representativeness of the analyses, specific sample weighting was applied for all statistical analyses to account for over sampling in the age group of 2–3 years and in South Australia. Nutrient intakes and dietary GI and GL were analysed as the mean of the two 24 h recalls. One-way ANOVA was used to test for differences in dietary GI and GL between the groups. Age- and sex-specific energy-adjusted nutrients/dietary GI residuals were calculated by linear regression, with nutrient of interest/dietary GI as the dependent variable and daily energy intake as the independent variable. Pearson's χ^2 test was used to test for differences in numbers of male participants, indigenous participants and participants from urban area across the age groups. Trends for mean dietary GI and GL across age groups were tested by linear regression, with age in years (continuous) as the independent variable and GI and GL residuals as the dependent variable. Pearson's correlation coefficients (by age group) between dietary GI and percentage energy from macronutrients and Ca were calculated: model 1 included adjustment for sex and BMI Z-score; model 2 calculated the correlation between the age- and sex-specific energy-adjusted residuals of dietary GI and percentage energy from macronutrients as well as from Ca, with additional adjustment for BMI Z-score. Per consumer analysis included only subjects who had reported the consumption of food item(s) in the food

groups tested. The Kruskal–Wallis test was used to test for differences in the GL contribution between boys and girls, as these were not normally distributed. A *P* value < 0.05 was considered to indicate statistical significance. All statistical analyses were carried out using Statistical Packages for Social Science version 17.0 (SPSS Australasia Private Limited, North Sydney, NSW, Australia).

Results

Subjects excluded from the analysis (*n* 460) were older (11.1 *v.* 8.3 years; *P* < 0.001) and had higher BMI (22.5 *v.* 18.3 kg/m²; *P* < 0.001) than the included subjects. There was also a higher proportion of girls among the excluded subjects (57.8 *v.* 48.9%; *P* < 0.001).

The weighted mean dietary GI and GL of the study population were 54.1 (SD 4.7) and 135.6 (SD 43.9), respectively. No significant difference by sex was detected for dietary GI, but boys had a significantly higher dietary GL than girls (145.5 (SD 47.4) *v.* 124.9 (SD 37.0); *P* < 0.001). There was no significant difference by state of residence (data not shown). Table 2 outlines the mean daily intake of select macro- and micronutrients by age groups of the participants. The age-dependent increase in total energy intake was accompanied by an increase in percentage energy from starch, as well as a decrease in percentage energy from sugars and CHO. There was no significant trend in GI or GL as age increased, but children aged 2–3 years had significantly lower GI and GL than their older counterparts. Fibre density showed significant decreasing trends.

Table 3 shows Pearson's correlation coefficients between dietary GI and selected macro- and micronutrients, stratified by age groups. Energy from non-glycaemic macronutrients was inversely associated with dietary GI. Among CHO, percentage energy from sugar was inversely associated with dietary GI, while percentage energy from starch was strongly

Table 2. Daily intake of selected macronutrients, fibre and demographics of the subjects by age group (Weighted mean values and standard deviations)

	2–3 years		4–8 years		9–13 years		14–16 years		<i>P</i> *
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
<i>n</i>	1068		1181		1019		916		–
BMI (kg/m ²)	24.1	84.9	18.3	41.2	21.1	44.2	21.5	16.6	0.070
Male (%)	51.1		51.2		51.6		54.6		0.447
Urban (%)	73.0		69.2		67.5		70.0		0.122
Indigenous (%)	2.7		2.4		3.4		3.1		0.492
Dietary GI	52.1	4.8	54.1	4.4	54.7	4.6	54.6	5.0	0.108
Dietary GL	97.8	25.7	122.2	31.8	149.1	40.5	164.3	50.9	0.066
Energy (kJ)	5966.9	1263.1	7240.7	1615.8	8776.2	2003.9	9956.2	2521.5	† < 0.001†
Energy from fat (%)	31.0	5.5	31.2	5.6	31.4	5.4	32.2	5.6	† < 0.001†
Energy from saturated fat (%)	14.1	3.5	14.1	3.4	14.1	3.2	14.1	3.4	0.131
Energy from protein (%)	17.0	3.3	16.8	3.4	17.0	3.7	17.5	4.0	† 0.001†
Energy from carbohydrates (%)	53.1	6.4	53.2	6.7	52.8	6.6	51.3	6.7	↓ < 0.001†
Energy from sugars (%)	27.9	6.3	26.4	6.5	25.3	6.4	24.0	6.6	↓ < 0.001†
Energy from starch (%)	24.8	6.1	26.3	5.9	27.0	6.2	26.8	5.9	† < 0.001†
Fibre density (g/MJ)	2.7	0.9	2.7	0.8	2.5	0.8	2.5	0.8	↓ < 0.001†

GI, glycaemic index; GL, glycaemic load.

* *P* values represent *P* for trend except for male (%), urban (%) and indigenous (%), which were tested by χ^2 test, and for BMI that was tested by one-way ANOVA.

† † and † preceding *P* for trend indicate the direction of trend.

Table 3. Pearson's correlation coefficient between dietary glycaemic index (GI) and selected nutrient intakes of the subjects by age group

	2–3 years		4–8 years		9–13 years		14–16 years	
	Model 1*	Model 2†	Model 1*	Model 2†	Model 1*	Model 2†	Model 1*	Model 2†
Dietary GL	0.44	0.70	0.40	0.64	0.35	0.63	0.44	0.70
Energy from fat (%)	–0.19	–0.20	–0.12	–0.12	–0.15	–0.14	–0.17	–0.18
Energy from saturated fat (%)	–0.28	–0.29	–0.21	–0.21	–0.28	–0.28	–0.25	–0.27
Energy from protein (%)	–0.22	–0.22	–0.14	–0.14	–0.10	–0.11	–0.15	–0.16
Energy from carbohydrates (%)	0.27	0.27	0.17	0.17	0.18	0.17	0.23	0.24
Energy from sugars (%)	–0.16	–0.17	–0.20	–0.20	–0.16	–0.16	–0.10	–0.10
Energy from starch (%)	0.44	0.45	0.41	0.41	0.34	0.34	0.34	0.35
Fibre density (g/MJ)	0.02‡	0.02‡	–0.09	–0.09	–0.08	–0.09	–0.14	–0.13
Ca (mg)	–0.37	–0.44	–0.31	–0.37	–0.34	–0.38	–0.31	–0.37

GL, glycaemic load.

* Correlation between dietary GI and the nutrient variables, adjusted for sex, BMI Z-score.

† Correlation between dietary GI and the nutrient variables (as age-, sex- and energy-adjusted residuals), adjusted for BMI Z-score.

‡ Correlation coefficient values were not statistically significant.

positively associated with dietary GI. A weak inverse correlation between fibre density and dietary GI was also found for all age groups, but not for the age group of 2–3 years, and Ca was found to be strongly inversely associated with dietary GI. Percentage energy from saturated fat was also negatively associated with dietary GI.

Table 4 shows the number of consumers for the top twenty GL-contributing food groups by age groups, while Table 5 shows the mean GL contribution by the top twenty GL-contributing food groups stratified by age groups. At age 2–3 years, the main contributors to the GL were fruit, breakfast cereals and white bread, while at age 14–16 years, white bread, breakfast cereals, cakes, pastries and doughnuts as well as soft drinks ranked among the top contributors. There were significant increases in percentage GL contributed by energy-dense and/or nutrient-poor foods such as soft drinks, white breads, high-fat potatoes and fast foods across age groups, while that for nutrient-dense foods such as fruit, milk and yogurt was found to be decreasing across age groups.

Per consumer analyses found that for all food groups, except fruit and milk, the daily absolute dietary GL contribution was increasing across age groups. Notably, the daily dietary GL contributed by white breads and that by energy-dense and/or nutrient-poor items such as cakes, pastry and doughnut, salty snacks and soft drinks, etc. in subjects aged 14–16 years was nearly or more than doubled that of their 2–3-year counterparts. Increases in nutrient-dense food groups across age group were more modest. When examined by sex (Table 6), per consumer analysis revealed that girls across all age groups had a significantly less percentage GL and daily GL from breakfast cereals, and older girls had less daily GL from confectionery and soft drinks than younger girls.

Discussion

The present study is the first to investigate the dietary GI and GL of a nationally representative sample of children and adolescents. We have shown that Australian children and adolescents were having a significant proportion of their

dietary GL from energy-dense and/or nutrient-poor food items, and these proportions were found to increase with age. The present results also suggest that Australian children and adolescents were having a high-GI dietary pattern characterised by high starch and low Ca intake. Apart from low Ca intake, overall, the nutritional intake of 2007ANCNPAS participants appeared to be satisfactory, with most of the participants meeting the estimated average requirement for key micronutrients⁽¹⁸⁾.

As expected, GI correlates positively with GL of which it is a component (model 1). Correlations are enhanced once potential variations due to differences in energy intake are accounted for (model 2), i.e. when GL can no longer vary due to increases or decreases in CHO. With this adjustment, GL variations are theoretically reduced to those attributable to exchanging CHO for protein or fat intake and to variations in GI itself, which explains the high correlation between this energy-adjusted GL and GI.

The weak negative correlation between percentage energy from sugar and GI suggested that the popular belief that high intake of sugary foods increases the dietary GI may be incorrect, at least among Australian children and adolescents. Fibre density, unlike Buyken *et al.*⁽¹⁹⁾ who showed a strong significant negative trend across GI tertiles, was only modestly correlated with GI in an inverse fashion. While soluble or viscous fibre is able to reduce the digestion rate of CHO by limiting the access of enzyme to the CHO (hence lowering the GI), insoluble fibre, when finely milled, e.g. in wholemeal flour, has no or minimal effect on glucose absorption rate because it does not block the access of enzymes to the CHO. Therefore, foods high in finely milled insoluble fibre may still have a high GI, e.g. wholemeal breads, and the consumption of these foods seemed to be higher in the 2007ANCNPAS population than those in the DONALD study. The AUSNUT2007 database used for this analysis unfortunately did not allow for the separation of the two types of fibre, thus disallowing further analysis.

There have been few investigations into the dietary GI and GL among children and adolescents in large samples or national surveys. A Canadian study of 4936 adolescents aged 9–17 years has reported a mean GI and GL of 55 and 144⁽¹⁰⁾.

Table 4. Number of consumers for the top twenty glycaemic load-contributing food groups by age group

	All subjects			2–3 years			4–8 years			9–13 years			14–16 years		
	Total	Boys	Girls	Total	Boys	Girls	Total	Boys	Girls	Total	Boys	Girls	Total	Boys	Girls
Biscuits*	2859	1464	1395	447	234	213	1079	544	535	909	466	443	424	220	204
Cakes, pastry and doughnuts*	2626	1333	1293	309	165	144	924	448	476	911	461	450	482	259	223
Confectionery*	2752	1391	1361	324	168	156	1021	527	494	926	462	464	481	234	247
Cordial*	1206	642	564	154	83	71	413	214	199	426	216	210	213	129	84
Fast foods*	1541	841	700	133	71	62	463	248	215	571	302	269	374	220	154
High-fat potatoes*	1584	809	775	182	88	94	532	281	251	530	258	272	340	182	158
Salty snacks*	1898	1029	869	174	91	83	725	389	336	660	360	300	339	189	150
Soft drinks*	1829	1036	793	103	53	50	497	274	223	766	431	335	463	278	185
Sugars*	2811	1477	1334	373	192	181	1019	533	486	928	488	440	491	264	227
Breakfast cereals	3032	1666	1366	455	232	223	1149	603	546	954	528	426	474	303	171
Cows' milk (excluding flavoured milk)	3640	1938	1702	517	264	253	1320	683	637	1183	629	554	620	362	258
Fruit and/or vegetable juice and fruit drinks	2522	1301	1221	335	174	161	907	485	422	812	403	409	468	239	229
Fruits	3418	1750	1668	513	262	251	1330	675	655	1062	541	521	513	272	241
Mixed-grain breads	928	468	460	166	92	74	363	176	187	242	121	121	157	79	78
Pasta and noodles	1781	937	844	268	142	126	631	335	296	566	291	275	316	169	147
Rice	984	519	465	135	65	70	338	175	163	337	185	152	174	94	80
Starchy vegetables (including low-fat potatoes)	2090	1058	1032	298	152	146	779	378	401	673	347	326	340	181	159
White breads	2979	1571	1408	340	173	167	1095	563	532	985	513	472	559	322	237
Wholemeal breads	1027	584	533	187	91	96	432	241	191	332	162	160	176	90	86
Yogurt	1330	682	648	283	147	136	560	281	279	331	174	157	156	80	76

* Energy-dense and/or nutrient-poor food groups.

Table 5. Glycaemic load (GL) contribution by the top twenty GL-contributing food groups by age group (Mean values and standard deviations)

Food groups	GL contribution (%)									Per consumer daily GL contribution*									
	2–3 years		4–8 years		9–13 years		14–16 years		$P_{\text{trend}}\dagger$	2–3 years		4–8 years		9–13 years		14–16 years		$P_{\text{trend}}\dagger$	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD	Mean	SD		
All participants (n 4184)																			
Fruit	10.9	8.2	8.1	6.3	5.4	5.2	4.3	5.2	↓ < 0.001	11.4	7.8	10.8	7.4	10.1	7.5	10.6	8.7		↓ 0.011
Breakfast cereals	10.7	8.7	9.9	8.5	10.0	9.5	9.4	10.8	↓ 0.010	13.0	9.2	15.7	11.4	21.9	15.3	26.3	20.7	↑	< 0.001
White breads	9.2	10.6	12.0	11.7	11.7	11.2	12.2	11.9	↑ 0.001	14.8	10.0	19.7	12.9	23.3	14.3	27.7	20.6	↑	< 0.001
Biscuits‡	6.9	6.3	6.4	6.7	4.8	5.7	4.0	5.8	↓ < 0.001	8.3	6.1	10.5	8.1	10.6	8.8	12.3	11.0	↑	< 0.001
Cows' milk (excluding flavoured milk)‡	5.9	4.8	3.7	3.1	3.1	2.6	2.7	2.8	↓ < 0.001	5.9	3.8	4.8	3.2	5.2	3.7	5.4	4.3		0.659
Cakes, pastry and doughnuts‡	5.4	7.4	6.6	8.2	7.8	8.7	7.6	9.0	↑ < 0.001	9.8	8.6	13.2	11.2	17.3	13.6	20.0	14.4	↑	< 0.001
Pasta and noodles	4.4	6.3	4.0	6.3	4.1	6.7	4.3	6.9	0.498	9.0	6.6	11.3	8.9	14.7	11.1	17.0	11.8	↑	< 0.001
Confectionery‡	4.3	6.2	5.5	6.8	5.7	7.0	5.7	7.9	↑ < 0.001	7.6	8.2	10.2	10.0	13.1	12.3	16.1	18.3	↑	< 0.001
Fruit and/or vegetable juice and fruit drinks	4.2	5.5	4.3	5.4	3.7	4.9	3.8	5.0	↓ 0.012	6.8	5.6	8.5	6.9	9.2	7.7	10.3	8.5	↑	< 0.001
Wholemeal breads	3.9	7.0	3.3	6.4	2.3	5.6	1.9	4.7	↓ < 0.001	10.8	6.9	13.2	8.0	13.4	9.2	13.1	9.0		↑ 0.041
Rice	3.4	8.6	3.5	8.1	4.1	9.2	3.8	8.8	↑ 0.040	14.7	14.7	20.7	16.4	27.4	20.9	31.3	25.0	↑	< 0.001
Mixed-grain breads	2.9	5.9	2.3	5.3	1.6	4.3	1.8	4.5	↓ < 0.001	9.2	6.3	10.5	6.8	12.4	8.4	14.1	9.9	↑	< 0.001
Sugars‡	2.9	3.8	3.2	3.8	3.0	4.1	3.0	3.8	0.327	4.3	4.3	5.7	5.2	6.9	7.2	7.8	7.1	↑	< 0.001
High-fat potatoes‡	2.6	4.9	3.1	5.5	3.7	5.8	4.9	7.2	↑ < 0.001	8.0	6.3	10.6	7.4	14.3	8.5	17.8	11.4	↑	< 0.001
Starchy vegetables (including low-fat potatoes)	2.5	3.8	2.6	3.9	2.8	4.1	2.5	4.1	0.895	4.4	3.6	5.7	5.3	8.1	6.0	9.0	7.0	↑	< 0.001
Yogurt	2.3	3.4	1.3	2.3	0.7	1.7	0.6	1.7	↓ < 0.001	4.3	3.2	3.9	3.3	4.3	3.3	4.7	4.0		↑ 0.042
Cordial‡	2.1	5.4	1.8	4.3	2.2	4.8	2.1	4.9	0.286	8.1	10.3	8.7	8.5	10.8	10.2	12.8	12.5	↑	< 0.001
Fast foods‡	1.8	4.5	2.7	5.6	4.2	6.8	5.7	8.5	↑ < 0.001	7.7	7.0	10.6	9.4	15.2	12.0	19.5	15.9	↑	< 0.001
Salty snacks‡	1.3	2.9	2.3	3.5	2.6	4.4	2.2	3.9	↑ < 0.001	4.0	4.1	5.8	5.1	8.0	7.7	8.0	7.7	↑	< 0.001
Soft drinks‡	1.0	2.8	2.4	4.7	5.1	6.7	6.6	8.7	↑ < 0.001	5.8	5.0	8.9	8.0	14.3	11.5	18.9	16.8	↑	< 0.001

* See Table 4 for the number of each cell.

† ↑ and ↓ preceding P for trend indicate the direction of trend.

‡ Energy-dense and/or nutrient-poor food groups.

Table 6. Glycaemic load (GL) contribution by the top twenty GL-contributing food groups by age group and sex (Mean values and standard deviations)

Food groups	Percentage of GL contribution (%)									Per consumer daily GL contribution*								
	2–3 years		4–8 years		9–13 years		14–16 years		<i>P</i> _{trend} †	2–3 years		4–8 years		9–13 years		14–16 years		<i>P</i> _{trend} †
	Mean	SD	Mean	SD	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Boys only (n 2134)																		
Breakfast cereals	11.5	9.4	10.3	8.7	11.2	9.8	11.6	11.3	0.084	14.6	10.6	17.1	13.2	24.8	17.1	29.5	20.8	↑ < 0.001
Fruit	10.9	8.3	7.9	6.2	4.9	4.9	3.9	4.9	↓ < 0.001	11.7	8.2	11.3	7.6	10.1	7.9	10.8	9.0	↓ 0.015
White breads	9.4	10.8	11.8	11.2	12.0	11.4	12.9	11.8	↑ < 0.001	15.6	10.5	20.2	13.4	25.0	15.4	30.5	22.6	↑ < 0.001
Biscuits‡	6.7	6.0	6.4	6.7	4.7	5.7	3.8	5.4	↓ < 0.001	8.3	6.3	11.4	8.4	11.3	9.5	13.3	10.9	↑ < 0.001
Cows' milk (excluding flavoured milk)	5.9	4.6	3.6	2.8	3.3	2.6	3.0	2.8	↓ < 0.001	6.0	3.7	5.0	3.3	5.8	4.0	6.2	4.5	↑ 0.003
Cakes, pastry and doughnuts‡	5.5	7.5	6.0	7.9	7.3	8.5	6.7	8.0	↑ 0.023	9.8	8.4	13.4	11.5	17.6	14.3	20.0	14.6	↑ < 0.001
Fruit and/or vegetable juice and fruit drinks	4.3	5.4	4.4	5.5	3.5	5.8	4.6	6.9	↑ < 0.001	7.2	5.7	9.0	7.4	9.5	6.4	11.3	10.0	↑ < 0.001
Pasta and noodles	4.2	5.9	3.9	6.1	4.0	6.6	4.0	6.4	0.943	8.6	6.4	11.5	9.6	15.5	11.2	18.1	12.5	↑ < 0.001
Confectionery‡	4.1	5.7	5.7	7.2	5.4	6.6	5.1	7.7	0.552	7.2	7.1	10.9	11.0	14.0	12.9	17.9	21.0	↑ < 0.001
Wholemeal breads	3.9	7.1	3.6	6.4	2.2	5.4	1.8	4.5	↓ < 0.001	11.7	7.0	13.8	7.9	13.7	9.9	14.0	8.9	0.359
Mixed-grain breads	3.1	5.8	2.2	5.0	1.5	4.1	1.7	4.5	↓ < 0.001	9.5	6.3	11.0	7.2	12.8	7.9	15.7	11.7	↑ < 0.001
Rice	3.0	7.7	3.6	8.5	4.3	9.4	3.5	8.6	0.189	13.7	11.5	22.3	16.9	28.9	21.8	33.7	30.2	↑ < 0.001
Sugars‡	3.0	3.9	3.2	3.9	3.1	4.2	2.9	3.7	0.919	4.5	4.3	6.0	5.5	7.2	7.6	8.2	7.3	↑ < 0.001
Starchy vegetables (including low-fat potatoes)	2.5	3.6	2.2	3.5	2.8	4.1	2.5	4.0	0.397	4.5	3.4	5.6	5.2	8.6	6.2	10.1	7.7	↑ < 0.001
High-fat potatoes‡	2.5	4.9	3.1	5.3	3.5	5.8	4.6	6.9	↑ < 0.001	8.1	6.1	10.8	7.5	15.6	9.4	18.6	10.8	↑ < 0.001
Cordial‡	2.4	6.2	1.8	4.3	2.1	4.7	2.3	5.2	0.847	9.0	12.2	9.1	9.1	11.2	10.7	13.9	14.1	↑ < 0.001
Yogurt	2.4	3.5	1.3	2.4	0.7	1.7	0.6	1.8	↓ < 0.001	4.5	3.4	4.2	4.0	4.7	3.8	5.1	4.7	0.083
Fast foods‡	2.0	4.8	2.7	5.4	4.1	6.5	6.3	8.6	↑ < 0.001	8.3	8.1	10.7	9.9	15.4	13.0	21.5	15.8	↑ < 0.001
Salty snacks‡	1.3	2.6	2.5	3.9	2.8	4.7	2.2	3.8	↑ 0.030	3.9	3.3	6.2	6.0	8.6	8.4	8.5	7.9	↑ < 0.001
Soft drinks‡	0.9	2.5	2.7	5.1	5.6	6.8	7.6	9.5	↑ < 0.001	5.3	4.2	9.7	8.8	15.5	12.5	21.4	18.6	↑ < 0.001
Girls only (n 2050)																		
Fruit	10.9	8.2	8.2	6.5	6.0§	5.4	4.9§	5.4	↓ < 0.001	11.0	7.4	10.2§	7.3	10.2	7.0	10.5	8.3	0.260
Breakfast cereals	9.9†	8.0	9.4§	8.4	8.6§	8.8	6.6§	9.4	↓ < 0.001	11.4§	7.2	14.0§	8.6	18.4§	11.7	20.6§	19.3	↑ < 0.001
White breads	9.1	10.3	12.2	11.3	11.4	10.9	11.3	12.0	0.236	14.1	9.5	19.2	12.3	21.3§	12.8	23.8§	16.8	↑ < 0.001
Biscuits‡	7.1	6.6	6.3	6.8	4.9	5.7	4.3	6.2	↓ < 0.001	8.4	5.9	9.5	7.6	9.9	7.9	11.2	11.1	↑ < 0.001
Cows' milk (excluding flavoured milk)	6.0	4.9	3.7	3.3	2.9§	2.7	2.2§	2.8	↓ < 0.001	5.7	4.0	4.6§	3.1	4.5§	3.2	4.2§	3.8	↓ < 0.001
Cakes, pastry and doughnuts‡	5.3	7.4	7.2§	8.4	8.3	9.0	8.7§	10.0	↑ < 0.001	9.7	8.8	13.1	10.8	16.9	12.9	20.0	14.3	↑ < 0.001
Pasta and noodles	4.6	6.6	4.0	6.5	4.2	6.8	4.7	7.5	0.357	9.4	6.8	11.1	8.0	13.7	11.0	15.8	10.8	↑ < 0.001
Confectionery‡	4.5	6.8	5.3	6.4	6.0	7.4	6.6§	8.1	↑ < 0.001	8.0	9.4	9.4§	8.7	12.1§	11.7	14.4§	15.1	↑ < 0.001
Fruit and/or vegetable juice and fruit drinks	4.1	5.6	4.0	5.4	3.9	5.3	4.3§	5.1	0.872	6.5	5.5	8.0§	6.2	9.0	8.8	9.2§	6.4	↑ < 0.001
Wholemeal breads	3.9	6.9	3.1	6.4	2.5	5.7	2.1	5.0	↓ < 0.001	9.9	6.7	12.5	8.1	13.1	8.5	12.2	9.0	↑ 0.037
Rice	3.8	9.4	3.3	7.7	3.9	8.9	4.2	9.0	0.108	15.7	17.2	19.1	15.6	25.5	19.6	28.5	16.9	↑ < 0.001
High-fat potatoes‡	2.7	4.9	3.2	5.7	3.9	5.7	5.2	7.6	↑ < 0.001	7.8	6.4	10.3	7.2	13.1§	7.4	16.9	11.9	↑ < 0.001
Mixed-grain breads	2.7	5.9	2.5	5.6	1.7	4.5	1.9	4.4	↓ < 0.001	8.8	6.4	9.9	6.3	12.1	8.9	12.4§	7.5	↑ < 0.001
Sugars‡	2.7	3.8	3.1	3.7	3.0	3.9	3.2	4.0	0.122	4.0	4.2	5.3†	4.8	6.5	6.7	7.4	7.0	↑ < 0.001
Starchy vegetables (including low-fat potatoes)	2.5	3.9	2.9§	4.3	2.8	4.0	2.6	4.2	0.564	4.2	3.9	5.8	5.3	7.6§	5.7	7.8§	5.9	↑ < 0.001
Yogurt	2.2	3.3	1.3	2.2	0.7	1.6	0.7	1.7	↓ < 0.001	4.1	2.8	3.6§	2.4	3.9§	2.7	4.3	2.9	0.303
Cordial‡	1.8	4.4	1.8	4.3	2.3	5.0	1.8	4.4	0.172	7.0	7.4	8.2	7.8	10.4	9.7	11.1	9.2	↑ < 0.001
Fast foods‡	1.7	4.1	2.8	5.8	4.4	7.1	5.0	8.3	↑ < 0.001	7.1	5.3	10.6	8.8	15.0	10.9	16.7§	15.6	↑ < 0.001
Salty snacks‡	1.3	3.1	2.1	3.1	2.4§	4.1	2.2	4.0	↑ < 0.001	4.2	4.9	5.4§	3.9	7.2§	6.6	7.4	7.4	↑ < 0.001
Soft drinks‡	1.1	3.2	2.1§	4.2	4.5§	6.5	5.3§	7.5	↑ < 0.001	6.2	5.7	7.9§	6.8	12.9§	10.0	15.3§	12.9	↑ < 0.001

* See Table 4 for the number of each cell.

† ↑ and ↓ preceding *p* for trend indicate the direction of trend.

‡ Energy-dense and/or nutrient-poor food groups.

§ Mean values were significantly different from those of boys in the same age group (*P* < 0.05).

A small Italian study of 105 children aged 8 years, which assessed the subjects' diets with a validated FFQ, has produced similar findings to the present study. The mean dietary GI and GL were found to be 58 and 145, respectively, and boys were found to have a higher GL but not a higher GI than girls. A Danish study⁽¹²⁾ of 849 children aged 10 and 16 years has also found that boys were having a higher dietary GL than girls. However, the dietary GL reported (mean GL for boys 231 (SD 67)) was much higher than that found in the present study for the 9–16-year-olds, and only 1 d of recall was obtained. The DONALD study⁽⁴⁾ found the dietary GI and GL of children at 2 years to be 52 and 63, and these were shown to have increased to 56 and 113 in 5 years, respectively, though not statistically significant. The present results showed that although there was no trend for GI or GL as age increases, older (≥ 4 years) children had a significantly higher GI and GL than their 2–3-year counterparts. Higher percentage GL from dairy foods and fruits, both having low GI, among participants aged 2–3 years may explain such a difference. The concurrent increase in dietary GI with age also suggests that older respondents were selecting more high/higher-GI foods in their diet.

A decrease in percentage energy from sugars with age was found, which seemed contradictory to the fact that daily GL from energy-dense and/or nutrient-poor foods, which were mostly sugary in nature, was increasing with age. Indeed, the DONALD study⁽⁴⁾ has shown that the percentage energy from added sugars increased from 9.5 to 14.2% in 5 years. Unfortunately, the AUSNUT2007 database used in this analysis did not allow the separation of added sugars from naturally occurring sugars.

An increase in percentage energy from starch with age was also observed, and this may partly explain the increase in dietary GI with age, as most starchy foods, especially those that are highly processed, are of high GI. Most of the subjects' dietary GL was from white breads and other energy-dense and/or nutrient-poor foods, such as soft drinks, high-fat potatoes, high-sugar, low-fibre breakfast cereals, cordial, etc. These foods have a moderate/high GI, and provide a small amount of fibre/nutrients or are high in energy, or both, making them poor dietary choices. Age-stratified analyses revealed that older respondents were having more of their dietary GL from energy-dense and/or nutrient-poor foods, and less from fruits.

Very few data are available for the food types contributing to the dietary GL among children and adolescents. The present findings differed from that of the German DONALD study⁽¹⁹⁾, which showed that German children in the study had high relative contributions to dietary GL by bread and rolls, as well as by milk and dairy products, and that contributions by breakfast cereals and cereal grains were lower than that of the Australian children in the present study. This suggests that the contribution to GL by different foods is specific to the food habits of the respective countries. However, 'tolerated food groups' (similar to the energy-dense and/or nutrient-poor foods described in the present study) contributed to a significant proportion of the subjects' GL in both studies, which raised concerns that the diets of these children were suboptimal.

Both high dietary GI and GL have recently been found to be associated with higher risk for chronic diseases^(1–3,20). The development of chronic diseases usually occurs slowly over a long period, and it has been suggested that long-term dietary behaviour could be shaped during childhood and adolescence^(21–23). Lowering the dietary GL may therefore be an effective, yet simple and easy strategy for Australian children and adolescents to reduce the risk of developing these chronic diseases in later life. In theory, by reducing the GI of the food items and/or the amount of CHO eaten could reduce the dietary GL. However, reducing the amount of CHO eaten could be difficult for people who follow a traditionally high-CHO diet, e.g. Asians. Utilising low- or lower-GI alternatives of their traditional staple foods, e.g. basmati rice for jasmine rice, is therefore a more suitable strategy to be employed⁽²⁴⁾.

A particular strength of the present study is the use of a published method for assigning GI values to the food items in the 2007ANCNPAS food database. Based on this method, we assigned GI values to around 85% of the food items in the first two steps, which utilised the current best available sources of GI values, therefore increasing the reliability of the GI values assigned. The use of a nationally representative sample (through sample weighting) also increased the generalisability of the findings.

However, there are several limitations to the present study. Despite being a suitable dietary assessment method to be used for a large number of subjects, the evidence to support the use of a 24 h dietary recall in children is limited, and it has been argued that an accurate dietary assessment among children is especially difficult⁽²⁵⁾. Parental recall of food intake, as in the case for children aged 8 years or below in the present study, had been suggested to be unreliable, especially when the reporting parent was away from home for more than 4 h/d (e.g. working)⁽²⁵⁾. This is likely to result in under-reporting and may hence contribute to inaccuracy at an individual level^(26–29). Children who reported their own food intake were also likely to inaccurately recall their food intake, due to incorrect identification of foods^(30,31) and unfamiliarity of the food⁽³²⁾, which may lead to misreporting, as well as information overload (e.g. large number of foods to report) that is likely to result in under-reporting⁽³³⁾.

By using the Goldberg cut-off for a specific PAL method⁽¹⁶⁾, we have excluded under- and over-reporters based on a scientifically accepted methodology, which is likely to increase the plausibility of the present findings. However, we were unable to detect incorrectly recalled foods in the database based on this method, which could possibly affect the calculated GI and GL, e.g. a low-GI mixed-grain bread may have been reported as 'brown bread', which could be subsequently coded as wholemeal bread, a high-GI food. We were also unable to exclude the possibility of residual error.

In addition, dietary intake may vary from day to day, and therefore data obtained from two 24 h recalls may not be representative of the habitual intake of an individual^(25,34). A 7 d food record would be ideal, but the large-scale nature of the 2007ANCNPAS made this impractical. It is also highly demanding on the participants and may result in a low response rate, i.e. data are not representative any more.

Also, while the two 24 h recalls for the same participant were collected only 7–21 d apart, the data collection for the whole study population spanned across autumn and winter in Australia, which may have arguably affected the consumption patterns of seasonal foods. Importation may have reduced the likelihood of unavailability of seasonal fruit and vegetables, though the higher price associated with it may still affect the consumption patterns. However, there are no notable differences in the availability of fruit and vegetables between autumn and winter in Australia⁽³⁵⁾.

In conclusion, the present results suggest that Australian children and adolescents were having a high-GI dietary pattern characterised by high starchy food intake and low Ca intake. A significant proportion of their dietary GL was from energy-dense and/or nutrient-poor foods. Effort should be made to encourage the replacement of high-GI, energy-dense and/or nutrient-poor food items for nutrient-dense low-GI foods to simultaneously lower the dietary GI and GL and improve the dietary quality of Australian children and adolescents.

Acknowledgements

The original data of the 2007ANCNPAS were collected by the Australian Commonwealth Scientific and Industrial Research Organization and the University of South Australia. The authors would like to thank the Australian Commonwealth Department of Health and Ageing for providing the survey data via the Australian Social Science Data Archive. The authors declare that those who carried out the original analysis and collection of the data bear no responsibility for further analysis or interpretation included in the manuscript. The present study received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. J. C. Y. L., A. E. B. and V. M. F. contributed to the conception of the study. J. C. Y. L. and K. H. assigned the GI values with input from V. M. F.; J. C. Y. L. and K. H. performed the statistical analyses under the guidance of A. E. B. and V. M. F.; A. E. B., V. M. F. and J. C. Y. L. interpreted the data. J. C. Y. L. drafted the manuscript. All authors were involved in the subsequent edits of the manuscript, and read and approved the final manuscript. The authors declare they have no competing interest.

References

1. Barclay AW, Petocz P, McMillan-Price J, *et al.* (2008) Glycemic index, glycemic load, and chronic disease risk – a meta-analysis of observational studies. *Am J Clin Nutr* **87**, 627–637.
2. Mente A, de Koning L, Shannon HS, *et al.* (2009) A systematic review of the evidence supporting a causal link between dietary factors and coronary heart disease. *Arch Intern Med* **169**, 659–669.
3. Liu S & Chou EL (2010) Dietary glycemic load and type 2 diabetes: modeling the glucose-raising potential of carbohydrates for prevention. *Am J Clin Nutr* **92**, 675–677.
4. Buyken AE, Cheng G, Gunther AL, *et al.* (2008) Relation of dietary glycemic index, glycemic load, added sugar intake, or fiber intake to the development of body composition between ages 2 and 7 y. *Am J Clin Nutr* **88**, 755–762.
5. Cheng G, Karaolis-Danckert N, Libuda L, *et al.* (2009) Relation of dietary glycemic index, glycemic load, and fiber and whole-grain intakes during puberty to the concurrent development of percent body fat and body mass index. *Am J Epidemiol* **169**, 667–677.
6. Benton D, Maconie A & Williams C (2007) The influence of the glycaemic load of breakfast on the behaviour of children in school. *Physiol Behav* **92**, 717–724.
7. Gilseman MB, de Bruin EA & Dye L (2009) The influence of carbohydrate on cognitive performance: a critical evaluation from the perspective of glycaemic load. *Br J Nutr* **101**, 941–949.
8. Ebbeling CB, Leidig MM, Sinclair KB, *et al.* (2003) A reduced-glycemic load diet in the treatment of adolescent obesity. *Arch Pediatr Adolesc Med* **157**, 773–779.
9. Fajcsak Z, Gabor A, Kovacs V, *et al.* (2008) The effects of 6-week low glycemic load diet based on low glycemic index foods in overweight/obese children – pilot study. *J Am Coll Nutr* **27**, 12–21.
10. Forbes LE, Storey KE, Fraser SN, *et al.* (2009) Dietary patterns associated with glycemic index and glycemic load among Alberta adolescents. *Appl Physiol Nutr Metab* **34**, 648–658.
11. Hui LL & Nelson EAS (2005) Meal glycaemic load of normal-weight and overweight Hong Kong children. *Eur J Clin Nutr* **60**, 220–227.
12. Nielsen BM, Bjornsbo KS, Tetens I, *et al.* (2005) Dietary glycaemic index and glycaemic load in Danish children in relation to body fatness. *Br J Nutr* **94**, 992–997.
13. Commonwealth Department of Health and Ageing, Department of Agriculture, Fisheries and Forestry, Australian Food and Grocery Council *et al.* (2008) *The 2007 National Children's Nutrition and Physical Activity Survey [Computer File]*. Canberra: Australian Social Science Data Archive, The Australian National University.
14. University of South Australia, Australian Commonwealth Scientific and Research Organization & i-View Pty Ltd (2009) User Guide – 2007 Australian National Children's Nutrition and Physical Activity Survey. [http://www.health.gov.au/internet/main/publishing.nsf/Content/AC3F256C715674D5CA2574D6000237D/\\$File/user-guide-v2.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/AC3F256C715674D5CA2574D6000237D/$File/user-guide-v2.pdf)
15. Food Standards Australia New Zealand (2008) AUSNUT 2007 <http://www.foodstandards.gov.au/monitoringandsurveillance/foodcompositionprogram/ausnut2007/index.cfm>
16. Goldberg GR, Black AE, Jebb SA, *et al.* (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* **45**, 569–581.
17. Louie JC-Y, Flood V, Turner N, *et al.* (2011) Methodology for adding glycemic index values to 24-hour recalls. *Nutrition* **27**, 59–64.
18. Commonwealth Scientific Industrial Research Organisation (Australia), Preventative Health National Research Flagship & Australia TUoS (2008) *2007 Australian National Children's Nutrition and Physical Activity Surveys – Main Findings*. Canberra: Department of Health and Ageing.
19. Buyken AE, Dettmann W, Kersting M, *et al.* (2005) Glycaemic index and glycaemic load in the diet of healthy school-children: trends from 1990 to 2002, contribution of different carbohydrate sources and relationships to dietary quality. *Br J Nutr* **94**, 796–803.
20. Liu S, Willett WC, Stampfer MJ, *et al.* (2000) A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women. *Am J Clin Nutr* **71**, 1455–1461.
21. Mikkilä V, Räsänen L, Raitakari OT, *et al.* (2005) Consistent dietary patterns identified from childhood to adulthood: the cardiovascular risk in Young Finns Study. *Br J Nutr* **93**, 923–931.

22. Westenhoefer J (2001) Establishing good dietary habits – capturing the minds of children. *Public Health Nutr* **4**, 125–129.
23. Pérez-Rodrigo C & Aranceta J (2001) School-based nutrition education: lessons learned and new perspectives. *Public Health Nutr* **4**, 131–139.
24. Liu S (2006) Lowering dietary glycemic load for weight control and cardiovascular health: a matter of quality. *Arch Intern Med* **166**, 1438–1439.
25. Livingstone MBE & Robson PJ (2000) Measurement of dietary intake in children. *Proc Nutr Soc* **59**, 279–293.
26. Klesges RC, Klesges LM, Brown G, *et al.* (1987) Validation of the 24-hour dietary recall in preschool children. *J Am Diet Assoc* **87**, 1383–1385.
27. Eck LH, Klesges RC & Hanson CL (1989) Recall of a child's intake from one meal: are parents accurate? *J Am Diet Assoc* **89**, 784–789.
28. Basch CE, Shea S, Arliss R, *et al.* (1990) Validation of mothers' reports of dietary intake by four to seven year-old children. *Am J Public Health* **80**, 1314–1317.
29. Baranowski T, Sprague D, Baranowski JH, *et al.* (1991) Accuracy of maternal dietary recall for preschool children. *J Am Diet Assoc* **91**, 669–674.
30. Samuelson G (1970) An epidemiological study of child health and nutrition in a northern Swedish County. II. Methodological study of the recall technique. *Nutr Metab* **12**, 321–340.
31. Emmons L & Hayes M (1973) Accuracy of 24-hr. recalls of young children. *J Am Diet Assoc* **62**, 409–415.
32. Warren JM, Henry CJK, Livingstone MBE, *et al.* (2003) How well do children aged 5–7 years recall food eaten at school lunch? *Public Health Nutr* **6**, 41–47.
33. Baranowski T, Dworkin R, Henske JC, *et al.* (1986) The accuracy of children's self-reports of diet: Family Health Project. *J Am Diet Assoc* **86**, 1381–1385.
34. Biro G, Hulshof KF, Ovesen L, *et al.* (2002) Selection of methodology to assess food intake. *Eur J Clin Nutr* **56**, Suppl. 2, S25–S32.
35. Vegetarian Network Victoria (2009) Seasonal Foods in Australia. <http://www.vnv.org.au/site/files/seasonalfoodcalendar.pdf>