

The role of nutrition and diet in rheumatoid arthritis

Rosemary H. Martin

Human Nutrition Research Group, University of Ulster, Coleraine BT52 1SA, UK

The nutritional implications of rheumatoid arthritis (RA) are due to a combination of the underlying disease pathology, disability and drug treatment. A wide range of conflicting dietary advice is available, yet the lack of scientific information regarding its efficacy often leaves sufferers vulnerable to exploitation. The recent increase in scientific investigations by the medical and related professions in this area represents an extremely welcome approach to remedy this situation.

Characteristics of rheumatoid arthritis

RA is a chronic, systemic inflammatory disease of unknown aetiology which principally affects the joints (Kumar *et al.* 1992). Worldwide, RA affects approximately 1% of the adult population and also occurs among one in 1000 children as juvenile RA (Di Toro & Polito, 1997). RA can strike at any age, yet typically occurs between 40 and 60 years of age (Bodman & Roitt, 1994). Approximately three females are affected for every male, and during pregnancy 70% of women suffering from RA experience remission, with flare-ups after birth (Nelson & Ostenson, 1997). As yet no definitive marker exists for RA. However, the auto-antibody rheumatoid factor is found in 70–80% of patients (Harris, 1990). There is an increased risk of RA among individuals positive for the human leucocyte antigen-DR4 allele of the major histocompatibility complex class II gene involved in antigen expression (Oiller & Thompson, 1992).

RA is characterized by painful, swollen joints, particularly of the fingers, wrists and knees. Symptoms may be 'symmetrical', with the same joints affected on opposite sides of the body. Typically, there is 'morning stiffness' of the joints (Kumar *et al.* 1992). Low-grade fever with flu-like symptoms, such as shivering, aching and fatigue, contributes to reduced appetite, weight loss and general malaise (Semble, 1995). The clinical course features periods of remission and relapse that are highly variable. Some patients remain stable, while sadly in others the disease is extremely aggressive, inflicting early disability (Hilliquin & Menkes, 1994).

Pathophysiology of inflammatory processes in rheumatoid arthritis

Normal human inflammatory responses are an essential component of the body's defence mechanisms against invading pathogens (Coleman *et al.* 1992). However, in RA these processes, once initiated, persist, resulting in the formation of a destructive, tumour-like tissue called 'pannus' (Bodman & Roitt, 1994). This invades soft tissues causing cartilage breakdown and bone erosion (Palmer, 1995). Inflammation is mediated by soluble proteins or cytokines (Panayi, 1993), the actions of which may be crudely simplified into those that are pro-inflammatory, e.g. interleukin-1 β and tumour necrosis factor- α , or anti-inflammatory, e.g. interleukin-10 and transforming growth factor- β (Koch *et al.* 1995).

Factors affecting nutrition in rheumatoid arthritis

Metabolic effects of inflammation

Tumour necrosis factor- α and interleukin-1 β bring about fever, increased catabolism, muscle wasting, connective tissue remodelling and anorexia associated with RA (Roubenoff *et al.* 1994). High levels of reactive oxygen species increase the oxidative stress on antioxidant defences (Grimble, 1990). Elevated plasma Cu, decreased plasma Zn and Fe are features of inflammation (Milanino *et al.* 1993) which are poorly understood, especially in RA. Recently, Roubenoff *et al.* (1997) have published findings indicating abnormal homocysteine metabolism in RA.

Physical disability

Ultimately, RA sufferers experience some degree of physical disability. Hands may become immobilized in a claw-like position (Kumar *et al.* 1992), making daily activities such as food preparation, eating and drinking extremely difficult. Poor mobility increases the dependence on others for shopping and cooking. Reduced physical activity contributes to loss of bone mass in long-standing RA (Hansen *et al.* 1996b), although this also depends on disease activity and drug therapy. Housebound individuals

Abbreviation: RA, rheumatoid arthritis.

Corresponding author: R. H. Martin, fax +44 (0)1265 324965, email RH.martin@ulst.ac.uk

may also be at risk of vitamin D deficiency in cases where there is little exposure to sunlight.

Drug treatment

RA sufferers undergo lifelong drug treatment to alleviate symptoms and delay disease progression. Most of the drugs used have significant side effects, involving the liver, kidneys, gastrointestinal tract and eyes (Semble, 1995). There are also nutritional implications in terms of altered appetite, nausea, vomiting, taste changes and altered nutrient absorption, metabolism and excretion (Brooks, 1994; Semble, 1995). Methotrexate is a known folate antagonist (Dijkmans, 1995), while non-steroidal anti-inflammatory drugs may increase gut mucosal permeability (Parke *et al.* 1996). Increased Ca and vitamin D intakes may protect against bone loss during steroid treatment for RA (Gulko & Mulloy, 1996), particularly in children where poor bone mineralization and growth failure are potential problems (Di Toro & Polito, 1997). Drugs place stress on detoxification systems operating via cytochrome P450, which utilizes riboflavin and thiamin (Parke *et al.* 1996).

Malnutrition in rheumatoid arthritis

The increased catabolism raises resting energy expenditure, which leads to weight loss and reduced lean body mass, especially if energy and protein requirements are not met (Roubenoff *et al.* 1994). This phenomenon is recognized as 'rheumatoid cachexia'. Helliwell *et al.* (1986) identified 26% of their RA group who were malnourished according to established anthropometric and biochemical indices. Malnutrition has been identified also among Finnish (Nenonen, 1995) and Spanish patients (Hernandez-Berriain *et al.* 1996) and, unsurprisingly, was more common among those who had severe disease. Low dietary intakes of Cu, Mg, Zn, vitamins D, E and A, pyridoxine, folate and *n*-3 fatty acids have been consistently reported by different investigators (Kremer & Bigaouette, 1996; Hansen *et al.* 1996a; Morgan *et al.* 1997).

Dietary approaches to rheumatoid arthritis

Dietary approaches to RA range from those based on sound nutritional principles to fads, myths and quackery. Rheumatologists have traditionally been sceptical, although viewpoints are changing. In contrast, RA patients are well known for their enthusiasm for dietary approaches. Of those surveyed, 75% of UK RA sufferers (Garret *et al.* 1993) and 33% of Danish RA sufferers (Haughen *et al.* 1991) believe that food plays a part in the severity of their symptoms. At least 20% will have tried a diet in their attempt to alleviate symptoms (Haughen *et al.* 1991). RA sufferers collectively spend billions of dollars on alternative treatments, including dietary therapies (Panush *et al.* 1983). Unfortunately, much of the advice is contradictory (Darlington *et al.* 1990), leaving RA sufferers confused as to what strategy to follow.

Some 'cultural beliefs' hold that RA is a 'cold' disease to be treated with 'hot' foods such as chilli, garlic and peas,

and 'hot' medicines such as aspirin, cod-liver oil, Fe and vitamins (Champion, 1994). Low-acid diets emerged from the belief that RA is due to acid-alkaline imbalance, and advocate the avoidance of citrus fruits and other acidic foods (Champion, 1994). Some authors have attempted to produce benefits via alteration of bowel flora using 'living' food (holistic) diets rich in lactobacilli (Nenonen, 1995).

'Cleansing', 'detoxifying' diets aim to purge the disease-causing toxins from the body. Fasting is a component of 'detoxifying', which can produce immediate relief from RA symptoms (Palmbad *et al.* 1991). However, the benefits are not lasting and symptoms generally deteriorate on return to normal eating or a vegetarian diet (Kjeldsen-Kragh *et al.* 1991). The mechanism behind such observations is unknown, but may relate to changes in intermediary metabolism or the removal of aggravating food antigens (Parke *et al.* 1996). Fasting is not recommended except under clinical supervision where genuine food allergy is suspected.

Elimination approach

Foods associated with symptoms of rheumatoid arthritis

Oils, fish and fresh vegetables are commonly associated with improvements in symptoms, while citrus fruits, chocolate, alcohol, red meats, flour products, spices and fizzy drinks are often implicated in the aggravation of symptoms (Darlington *et al.* 1990; Haughen *et al.* 1991; Garret *et al.* 1993). Such lists reflect those of the popular 'Dong' diet (Dong & Banks, 1975) for RA, based on the Chinese poor man's diet: rich in seafood, vegetables and rice, but eliminating dairy produce, alcohol, additives, spices and meat, allowing occasional poultry. In a clinical trial of this type of diet, Panush *et al.* (1983) found no consistent effect on disease, with the exception of two patients who experienced considerable improvement. Given that immunological mechanisms are important in RA pathology, and that despite research no 'trigger' antigens have yet been isolated, many patients and some clinicians have asked: 'Are there dietary causes of RA?'

Food intolerance in rheumatoid arthritis

Potential food antigens may be intrinsic to the actual foodstuff or arise from chemicals added to the food as preservatives, fertilizers or pesticides. The gut is a major organ in immune defence, and intestinal antigens may cross the gut barrier, eventually circulating as immune complexes (Buchanan *et al.* 1991). Absorption of dietary antigens from the gut may be enhanced by changes in gut permeability owing to immaturity, immunodeficiency, allergy or concomitant disease such as Crohn's disease (Parke *et al.* 1996).

Evidence to date for the role of food intolerance in RA is inconclusive, derived mainly from anecdotal reports. Milk and dairy produce, cereals, maize, wheat gluten, tartrazine and azo dyes have all been implicated in RA in recent reviews (Buchanan *et al.* 1991; Darlington & Ramsey, 1993). Overall, there seem to exist 'subgroups' of RA patients for whom food intolerance plays a significant role

in their disease. Carefully-designed double-blind clinical trials are needed to distinguish true intolerance from disease which spontaneously relapses and remits.

Dietary supplementation

Nutrient megadosing

The old adage: 'if a little is good, more is better' reflects the practice of nutrient megadosing, whether self administered or prescribed by 'alternative practitioners'. Commonly advocated are vitamin E, Zn and Se (Champion, 1994). Se supplementation trials in RA have improved Se status without any consistent effect on disease activity (Tarp, 1995). Apart from the expense and risk of toxicity, moderate nutritional supplementation may interfere with normal regulation of nutrient bioavailability; for example, high doses of Zn inhibit Cu absorption (Bremner & Beattie, 1995).

Novel food supplements

Novel food supplements are common to the shelves of any health food shop and some pharmacies. There is only slight evidence that the New Zealand green-lipped mussel (*Perna canaliculus*) extract, or 'Seatone' is of value in RA (Larkin, 1994). Devil's claw (*Harpagophytum*)-root extract, sea kelp, ginseng, and lactobacillus are others recommended in books (Darlington *et al.* 1990). Such products undoubtedly have powerful placebo effects, necessitating properly-controlled clinical trials to investigate their reported efficacy in RA.

Fish and plant oils

Oil-based supplements have emerged as the most promising area of dietary manipulation in RA. Pharmacological doses (3–10 g/d) of the long-chain *n*-3 fatty acids found in fish oils reduce stiffness and pain in RA (Kremer, 1996). Their mechanisms of action include the inhibition of pro-inflammatory eicosanoids (De Luca *et al.* 1995) and cytokines (Endres & von Shacky, 1996), although clearly other mechanisms exist. The value of low doses of long-chain *n*-3 fatty acids is unclear; however, frequent intakes of oily fish may reduce the risk of RA (Shapiro *et al.* 1996), or limit its severity (Recht *et al.* 1990).

Summary

Given the lack of understanding of the nutritional requirements in RA, plus the variability in its clinical course, it is difficult to produce specific dietary recommendations for RA. In general, sufferers should consume as varied a diet as possible, based on current Department of Health (1991) guidelines. Dietary counselling is important to help patients achieve this. Self-imposed elimination diets should be avoided and suspected food intolerance tested under strict clinical supervision. Nutrient megadosing is inadvisable, although dietary supplementation with Ca,

vitamin D, folic acid or multivitamins and minerals should be recommended where necessary.

References

- Bodman KB & Roitt IM (1994) The pathophysiology of rheumatoid arthritis. *Fundamental and Clinical Rheumatology* **2**, 73–81.
- Bremner I & Beattie JH (1995) Copper and zinc metabolism in health and disease: speciation and interactions. *Proceedings of the Nutrition Society* **54**, 489–499.
- Brooks PM (1994) Rheumatoid arthritis: pharmacological approaches. In *Rheumatology*, pp. 10.1–10.6 [PM Dieppe and JH Klippel, editors]. London: Mosby Year Book Ltd.
- Buchanan HM, Preston SJ, Brooks PM & Buchanan WW (1991) Is diet important in rheumatoid arthritis? *British Journal of Rheumatology* **30**, 125–135.
- Champion GD (1994) Rheumatoid arthritis: unproven remedies: alternative and complementary medicine. In *Rheumatology*, pp. 19.09–19.11 [PM Dieppe and JH Klippel, editors]. London: Mosby Year Book Ltd.
- Coleman RM, Lombard MF & Sicard RE (1992) *Fundamental Immunology*, 2nd ed., pp. 210–227. Dubuque, IA: W.C. Brown.
- Darlington LG, Jump A & Ramsey NW (1990) Dietary treatment of rheumatoid arthritis. *The Practitioner* **234**, 456–460.
- Darlington LG & Ramsey NW (1993) Review of dietary therapy for rheumatoid arthritis. *British Journal of Rheumatology* **32**, 507–514.
- De Luca P, Roithman D & Zurier RB (1995) Marine and botanical lipids as immunomodulatory and therapeutic agents in the treatment of rheumatoid arthritis. *Rheumatic Disease Clinics of North America* **21**, 759–777.
- Department of Health (1991) *Dietary Reference Values for Food, Energy and Nutrients for the United Kingdom. Report on Health and Social Subjects* no. 41. London: H.M. Stationery Office.
- Dijkmans BAC (1995) Folate supplementation and methotrexate. *British Journal of Rheumatology* **34**, 1172–1174.
- DiToro R & Polito C (1997) Nutrition in juvenile rheumatoid arthritis. *Nutrition Research* **17**, 741–758.
- Dong CH & Banks J (1975) *New Hope for the Arthritic*. New York: Ballantine Publishers.
- Endres S & von Shacky C (1996) *n*-3 Polyunsaturated fatty acids and human cytokine synthesis. *Current Opinion in Lipidology* **7**, 48–52.
- Garret SL, Kennedy LG & Calin A (1993) Patient's perceptions of disease modulation by diet in inflammatory (rheumatoid arthritis/ankylosing spondylitis) and degenerative joint disease. *British Journal of Rheumatology* **32**, Suppl. 2, 24 Abstr.
- Grimble RF (1990) Nutrition and cytokine action. *Nutrition Research Reviews* **3**, 193–210.
- Gulko PS & Mulloy AL (1996) Glucocorticoid induced osteoporosis: pathogenesis, prevention and treatment. *Clinical Experimental Rheumatology* **14**, 199–206.
- Hansen GVO, Nielsen L, Kluger E, Thyssen M, Emmertsen H, Stengaard-Pedersen K, Hansen EL, Unger B & Andersen PW (1996a) Nutritional status of Danish rheumatoid arthritis patients and effects of a diet adjusted in energy intake, fish-meal, and antioxidants. *Scandinavian Journal of Rheumatology* **25**, 325–330.
- Hansen M, Florescu A, Podenphant J, Pedersen-Zbinden B, Horslev-Petersen K, Hyldstrup L & Lorenzen I (1996b) Bone loss in rheumatoid arthritis. *Scandinavian Journal of Rheumatology* **25**, 367–376.

- Harris ED (1990) Rheumatoid arthritis. Physiology and implications for therapy. *New England Journal of Medicine* **322**, 1277–1287.
- Haughen M, Kjeldsen-Kragh J, Nordvåg BY & Førre Ø (1991) Diet and disease symptoms in rheumatic diseases—results of a questionnaire based survey. *Clinical Rheumatology* **10**, 401–407.
- Helliwell M, Coombes EJ, Moody BJ, Batstone GF & Robertson JC (1986) Nutritional status in patients with rheumatoid arthritis. *Annals of the Rheumatic Diseases* **43**, 386–390.
- Hernandez-Berain JA, Segura-Garcia C, Rodriguez-Lozano B, Bustabad S, Gantes M & Gonzalez T (1996) Under nutrition in rheumatoid arthritis patients with disability. *Scandinavian Journal of Rheumatology* **25**, 383–387.
- Hilliquin P & Menkes C-J (1994) Rheumatoid arthritis evaluation and management: early and late disease. In *Rheumatology*, pp. 13.1–13.14 [PM Dieppe and JH Klippe, editors]. London: Mosby Year Book Ltd.
- Kjeldsen-Kragh J, Haughen M, Borchgrevink CF, Laerum E, Eek M, Mowinkel P, Hovi K & Førre Ø (1991) Controlled trial of fasting and one year vegetarian diet in RA. *Lancet* **338**, 899–902.
- Koch A, Kunkel SL & Strieter RM (1995) Cytokines in rheumatoid arthritis. *Journal of Investigative Medicine* **43**, 28–38.
- Kremer JM (1996) Effects of modulation of inflammatory and immune parameters in patients with rheumatic and inflammatory disease receiving dietary supplementation of *n*-3 and *n*-6 fatty acids. *Lipids* **31**, Suppl., S243–S247.
- Kremer JM & Bigaouette J (1996) Nutrient intake of patients with rheumatoid arthritis is deficient in pyridoxine, zinc, copper and magnesium. *Journal of Rheumatology* **23**, 990–994.
- Kumar V, Cotran RS & Robbins SL (1992) Disorders of the immune system: Rheumatoid arthritis. *Basic Pathology*, 5th ed., pp. 145–149. London: W.B. Saunders & Co.
- Larkin JG (1994) Alternative medicine and rheumatoid arthritis. *Rheumatology Review* **3**, 61–71.
- Milanino R, Marrella M, Gasperini R, Pasqualicchio M & Velo G (1993) Copper and zinc body levels in inflammation: an overview of the data obtained from animal and human studies. *Agents and Actions* **39**, 195–209.
- Morgan SL, Anderson AM, Hood SM, Matthews A, Lee JY & Alarcón GS (1997) Nutrient intake patterns, body mass index, and vitamin levels in patients with rheumatoid arthritis. *Arthritis Care and Research* **10**, 9–19.
- Nelson JL & Ostenson M (1997) Pregnancy and rheumatoid arthritis. *Rheumatic Disease Clinics of North America* **23**, 195–212.
- Nononen MT (1995) Vegan diet, rich in lactobacilli ('living food'): metabolic and subjective responses in healthy subjects and in patients with rheumatoid arthritis. Dissertation, University of Kuopio, Finland.
- Oiller W & Thompson W (1992) Population genetics of rheumatoid arthritis. *Rheumatic Disease Clinics of North America* **18**, 741–759.
- Palmlblad J, Hafstrom I & Ringertz B (1991) Anti-rheumatic effects of fasting. *Rheumatic Disease Clinics of North America* **17**, 351–362.
- Palmer DG (1995) The anatomy of the rheumatoid lesion. *British Medical Bulletin* **51**, 286–295.
- Panayi GS (1993) The immunopathogenesis of rheumatoid arthritis. *British Journal of Rheumatism* **32**, Suppl. 1, 4–14.
- Panush RS, Carter RL, Katz P, Kowsari B, Longley S & Finne S (1983) Diet therapy for rheumatoid arthritis. *Arthritis and Rheumatism* **26**, 462–471.
- Parke AL, Parke DV & Jones FA (1996) Diet and nutrition in rheumatoid arthritis and other chronic inflammatory diseases. *Journal of Clinical Biochemistry and Nutrition* **20**, 1–6.
- Recht L, Helin P, Rasmussen JO, Jacobsen J, Lithman T & Bschersten BS (1990) Hand handicap and rheumatoid arthritis in a fish-eating society (the Faroe Islands). *Journal of Internal Medicine* **227**, 49–55.
- Roubenoff R, Dellarpia P, Nadeau MR, Abad LW, Muldoon BA, Selhub J & Rosenberg IH (1997) Abnormal homocysteine metabolism in rheumatoid arthritis. *Arthritis and Rheumatism* **40**, 718–722.
- Roubenoff R, Roubenoff RA & Cannon JG (1994) Rheumatoid cachexia: cytokine driven hypermetabolism accompanying reduced body cell mass in chronic inflammation. *Journal of Clinical Investigation* **93**, 2379–2386.
- Semble EL (1995) Rheumatoid arthritis: new approaches for its evaluation and management. *Archives of Physical Medicine and Rehabilitation* **76**, 190–201.
- Shapiro JA, Koepsell TD, Voight L, Dugawson CE, Kestin M & Nelson JL (1996) Diet and rheumatoid arthritis in women: Possible preventative effect of fish consumption. *Epidemiology* **7**, 256–263.
- Tarp U (1995) Selenium in rheumatoid arthritis. *Analyst* **120**, 877–881.