## Laryngology & Otology

cambridge.org/jlo

### **Main Article**

Alexander Yao takes responsibility for the integrity of the content of the paper

**Cite this article:** Yao A, Richards E, Dalton CL. Trainee-perceived benefits of a virtual temporal bone competition. *J Laryngol Otol* 2024;**138**:864–868. https://doi.org/10.1017/ S0022215124000070

Received: 22 July 2023 Revised: 22 October 2023 Accepted: 2 November 2023 First published online: 14 February 2024

#### Keywords:

Simulation training; virtual reality; temporal bone; otolaryngology; medical education

**Corresponding author:** Alexander Yao; Email: alexander.yao@nhs.net

# Trainee-perceived benefits of a virtual temporal bone competition

Alexander Yao<sup>1</sup>, Emma Richards<sup>2</sup> b and C. Lucy Dalton<sup>3</sup>

<sup>1</sup>ENT Department, Worcester Royal Hospital, Worcester, UK, <sup>2</sup>ENT Department, Princess Royal Hospital, Telford, UK and <sup>3</sup>ENT Department, Queen Elizabeth Hospital, Birmingham, UK

#### Abstract

**Objective.** To assess the perceived benefits of a novel educational approach for otolaryngology trainees: a virtual reality temporal bone simulator drilling competition.

**Methods.** Regional otolaryngology trainees participated in the competition. Drilling activities using the Voxel-Man TempoSurg simulator were scored by experts. Questionnaires that contained questions covering motivators for attending, perceived learning and enjoyment were sent to participants. Agreement with statements was measured on a 10-point Likert scale (1 = strongly disagree, 10 = strongly agree).

**Results.** Eighteen trainees participated. The most cited reason for attending was for learning and/or education (61 per cent), with most attendees (72 per cent) believing that competition encourages more reading and/or practice. Seventeen attendees (94 per cent) believed Voxel-Man TempoSurg-based simulation would help to improve intra-operative performance in mastoidectomy (mean  $7.83 \pm 1.47$ , p < 0.001) and understanding of anatomy (mean  $8.72 \pm 1.13$ , p < 0.001). All participants rated the competition as 'fun' and 83 per cent believed the competitive element added to this.

**Conclusion.** The virtual reality temporal bone competition is a novel educational approach within otolaryngology that was positively received by otolaryngology trainees.

#### Introduction

In recent years, recommendations from the Shape of Training Report<sup>1</sup> have meant further emphasis has been placed on the training of broader-based emergency-safe ENT surgeons with the knowledge and skills to treat a variety of acute ENT conditions. Of these skills, navigating the temporal bone anatomy during mastoid surgery represents a challenging yet essential part of the surgical repertoire.

Surgeons in training must develop the fine dexterity required to accurately manipulate a drill within the mastoid bone, visualised down the field of view of an operating microscope. Emergency conditions such as sub-periosteal abscesses complicating acute mastoiditis are uncommon, but are associated with high-morbidity.<sup>2</sup> Cortical mastoidect-omy involves drilling through the cortex of the mastoid, exenterating air cells, thus externalising any infection. It represents a 'gold standard' treatment for sub-periosteal abscesses, superior to simple incision and drainage.<sup>3,4</sup> The new August 2021 otolaryngology curriculum expects trainees to be competent to perform this procedure to the level of a day-one consultant by the end of phase 2 of training.<sup>5</sup>

Despite the need for experience in temporal bone drilling, opportunities to practice are limited. Over the past three decades, post-graduate surgical training within the UK has seen several major changes: Calman reforms,<sup>6</sup> European Working Time Directive legislation<sup>7</sup> and the introduction of Modernising Medical Careers.<sup>8</sup> More recently, the aerosol-generating nature of temporal bone drilling<sup>9</sup> has meant the coronavirus disease 2019 (Covid-19) pandemic further limited training opportunities in theatre for ENT trainees.<sup>10</sup> Virtual reality models such as the Voxel-Man TempoSurg simulator offer safe, accessible alternatives for learning temporal bone anatomy<sup>11</sup> and temporal bone drilling.<sup>12,13</sup> Simulation may provide more frequent access to learning opportunities than alternatives such as cadaveric temporal bone drilling, thus enhancing learning<sup>14</sup> and enabling deliberate practice.<sup>15</sup>

Interweaving competition into surgical training exercises may offer additional benefits. In laparoscopic surgery, competition has been shown to significantly increase the surgical efficiency as measured by the number of movements and instrument path length,<sup>16</sup> and has been used to achieve post-graduate fundamental proficiency standards by undergraduates.<sup>17</sup> Furthermore, gamification through competition reportedly improves the overall engagement and enjoyment of traditionally challenging areas of the undergraduate curriculum, such as anatomy.<sup>18</sup> It can also stimulate participants to consider ENT as a career choice.<sup>19</sup>

Surgical competitions have potential inherent psychosocial benefits, for example surrounding well-being, the importance of which has become particularly apparent during the Covid-19 pandemic.<sup>20</sup> In addition, they facilitate successful establishment and

© The Author(s), 2024. Published by Cambridge University Press on behalf of J.L.O. (1984) LIMITED integration of new learners into communities of practice by, for example, promoting broad participation, providing the opportunity for focus and being held in one location at one point in time.<sup>21</sup>

To our knowledge, only one previous Voxel-Man TempoSurg simulator competition, limited to core trainees, has been reported in the literature.<sup>19</sup> Our competition represents the largest temporal bone competition utilising the Voxel-Man TempoSurg simulator described in the current literature open to all training grades and equivalent non-training grades. Our study aimed to identify the trainee-perceived educational benefits of a virtual reality temporal bone drilling competition in the context of ENT training.

#### Materials and methods

#### Recruitment

All 35 ENT higher surgical trainees (ST3–ST8), 6 ENT ST1/2 run-through trainees and 6 CT1/2 ENT-themed trainees, as well as staff-grade equivalents within the West Midlands Local Education Training Board region were invited to participate in a free virtual reality temporal bone competition via email. Competition places were limited to 18 in total, with places allocated on a first-come, first-served basis.

#### Pre-competition data collection

Competition candidates were emailed a pre-competition electronic survey using Google Forms (Appendix A), to be completed before the competition date. The questions asked about prior experience with mastoid surgery, candidates' confidence performing a cortical mastoidectomy, previous experience with ENT simulators and ENT competition experience. Answers were scored on a 10-point Likert scale.

#### **Competition details**

The competition was held at the regional ENT Simulation Dry Laboratory, Queen Elizabeth Hospital, Birmingham, over two days, with participants competing on either day one or day two. Competitors were divided into junior (grades up to and including ST3) and senior (more senior than ST3) groups.

Participants were set two temporal bone drilling tasks on the Voxel-Man TempoSurg simulator lasting 30 and 90 minutes, respectively. The first was focused on drilling under time pressure, whilst the second was focused on demonstrating anatomy. Trainees were independently rated on both their surgical technique and final outcome by a judging panel consisting of three experts (consultant otologists). Raters used the previously validated modified Welling,<sup>22</sup> Stanford<sup>23</sup> and Intercollegiate Surgical Curriculum Programme<sup>5</sup> scales to calculate the overall combined competition score and identify winners (Appendix B).

After completing both drilling tasks, trainees additionally each received verbal feedback from the faculty lasting up to 10 minutes. Prizes were available for the first- and secondplace scorers in the junior and senior groups in each category. One 'Commendation' was awarded at the judges' discretion.

#### Post-competition data collection

After the competition, before prizes were announced, all participants were asked to complete a post-competition questionnaire using Google Forms (see Appendix A). Participants were asked about their experience using the simulator and of the competition in general, and how the competitive element might have affected their preparation or performance on the day, again using a 10-point Likert scale (1 = negative, 10 = positive).

#### Statistical analysis

Google Sheets was used to calculate basic summary data (mean, range and standard deviation). R was used to perform the Shapiro–Wilk test for normality and statistical tests.<sup>24</sup> The appropriate test was selected from Pearson's correlation coefficient, Spearman's rank test and the two-sample paired student's *t*-test. The one-sample Wilcoxon test or the one-sample student's *t*-test was used where the 10-point Likert scale was two-tailed ( $\mu$  was assumed to be 5.5, significance *p* < 0.025). Microsoft Excel (2023) was used to calculate the one-sample student's *t*-test.

#### **Results and analysis**

#### Participants

Eighteen trainees (12 male, 6 female) enrolled in the competition, including 3 CT2, 3 ST3, 2 junior trust grades (Senior House Officer level equivalent), 5 ST5, 3 ST6, 1 ST8 and 1 senior trust grade (Registrar level equivalent). All agreed to participate in this study.

#### Previous experience

Baseline relevant operative experience and course attendance are reported in Table 1. Seven participants had not performed any cortical mastoidectomies previously. Eight participants had not performed any mastoid explorations previously. Confidence in performing a cortical mastoidectomy was selfrated as a mean of  $4.83 \pm 2.41$  on a 10-point Likert scale (1 = extremely unconfident, 10 = extremely confident). Three participants had not attended any cadaveric bone courses previously and two had no previous experience on the virtual reality temporal bone simulator.

Four participants (22 per cent) had used ENT simulators before, including plastic models for laryngoscopy, tracheostomy, nasal packing and balloon sinus dilatation. One participant had used a sheep larynx model for tracheostomy. Four participants (22 per cent) had previously taken part in surgery-related competitions, mostly essay-writing competitions (Table 2).

#### Competition and motivation

The most cited reason for choosing to attend the competition was for learning or education, followed by a desire to compete (Figure 1).

The perceived influence of the competition as a motivator to practice or read around temporal bone surgery is described in Table 3. Thirteen participants (72 per cent) believed that a competitive element motivates more practice and/or reading compared with a non-competitive event (mean 7.06 ± 2.32, t = 2.85, p = 0.011).


Fourteen participants (78 per cent) reported undertaking some preparation for the competition. Of these, seven prepared via practice on the Voxel-Man TempoSurg simulator,

 Table 1. Summary of competitors' previous baseline experience (n = 18)

Previous experience	Mean	Median	Range
Number of cortical mastoidectomies performed	$5.28\pm6.63$	4	0–20
Number of mastoid explorations performed	3.17 ± 4.45	1	0-15
Number of cadaveric bone courses attended	$2.17 \pm 1.79$	2	0-6
Hours spent on the virtual reality temporal bone simulator	7.22 ± 4.86	9	0-16

Table 2. Summary of competitor's previous experience of surgical competitions

Type of competition	Number of candidates
Writing competition	3
Royal College of Surgeons of Edinburgh medical student competition	2
Laparoscopic surgery competition	1
Keele anatomy competition	1



**Figure 1.** The cited motivators for attending the virtual reality temporal bone simulation competition reported by candidates, organised by frequency of reason given. Each candidate could cite more than one reason for attending.

seven did reading around the topic and three watched videos of the procedure or attended theatre. Four participants did more than one type of preparation.

All participants (100 per cent) claimed they would take further action as a result of taking part in this competition. Eleven participants (61 per cent) planned to undertake practice on the simulator or in the operative setting, and 9 (50 per cent) planned to do further reading around the topic.

#### Competition and training perceptions

Participants overall agreed that 'surgical simulation improves intra-operative performance' (mean  $8.28 \pm 1.78$ , V = 169, p < 0.001). All 18 participants rated the competition as educational (mean  $8.22 \pm 1.17$ , t = 2.41, p = 0.0278). Participants rated their agreement with the statement 'the competitive element made [the event] more educational' a mean of  $6.67 \pm 2.06$  (V = 132, p = 0.0416), with 15 participants (83 per cent) agreeing with this statement.

Seventeen trainees (94 per cent) believed simulation with the Voxel-Man TempoSurg simulator improved intra-operative performance in cortical mastoidectomy, mastoid surgery and

Table 3. Influence of competition on learning activity as reported by candidates

Influence of competition on motivation to	1–10 Likert scale*	p value
Practice on the Voxel-Man TempoSurg simulator	$6.17\pm2.43$	0.261
Read around mastoid surgery	$6.56 \pm 2.66$	0.111
Read around temporal bone anatomy	$6.56 \pm 2.73$	0.118

\*1 = negative motivator; 10 = positive motivator

understanding of temporal bone anatomy (Table 4). Participants' self-rated confidence in performing cortical mastoidectomy improved from a pre-competition mean of  $4.83 \pm 2.41$  to  $5.72 \pm 2.22$  post-competition (t = 3.49, p = 0.003).

Seventeen participants (94 per cent) agreed that 'Competitions are useful within the ENT training programme' (mean 8.00 ± 1.46, t = 7.29, p < 0.001), but only 6 (33 per cent) agreed they 'should be made essential' (mean 3.83 ± 2.83, t = -2.5, p = 0.0231).

Seventeen participants (94.4 per cent) rated the competition as 'fun' and quantified this with a mean of  $8.06 \pm 1.21$  (t =8.95, p = <0.001) on a 10-point Likert scale (1 = extremely not fun, 10 = extremely fun). Thirteen participants (72 per cent) agreed that 'the competitive element made it more fun' (mean  $6.83 \pm 1.92$ , t = 2.95, p < 0.001). The most enjoyable elements of the competition were deemed to be peer or social interaction (mentioned 10 times), interaction with faculty (reported 5 times) and food (reported 3 times). Seventeen participants (94 per cent) stated they would attend another competition similar to this one (mean  $8.5 \pm 1.54$ , V = 169, p <0.001).

#### Discussion

The results from this study suggest that trainee participants perceived the virtual reality temporal bone competition to be foremost a highly valuable educational tool. Their perceptions of simulation-based learning parallel closely previously described key benefits, including the opportunity for feedback, deliberate practice, mastery learning and curriculum integration.<sup>25</sup> The most cited motivator for attending the competition in our study was for the learning potential, rather than a desire to win or compete.

All participants agreed the competition was educational, and 83 per cent agreed the competitive element made it more so. The competition encouraged preparative work in most trainees (78 per cent), including reading, simulator practice or watching videos and/or surgery, and further encouraged participants to take ongoing action after the competition. All trainees reported they would take further action as a result of the competition. However, the influence of the competition to motivate these activities did not reach statistical significance, suggesting that the external motivation provided by a virtual reality temporal bone competition alone is insufficient to bring a strong effect.

Trainees, like most adult learners, are likely to be driven more by intrinsic motivators,<sup>26</sup> thus viewing the competition more as a formative exercise. In contrast, in a larger-scale national Canadian cardiothoracic surgical competition, competitors did see the competition as a significant motivator for increased use of simulation devices.<sup>27</sup> Top performers had accrued many more hours of surgical simulation practice and studying the curriculum compared with those who did not

Table 4. Candidates' perceptions towards the benefits of the Voxel-Man TempoSurg simulator

Statement	Mean agreement on 1–10 Likert scale*	p value
Surgical simulation with the Voxel-Man TempoSurg improves intra-operative performance in cortical mastoidectomy	7.83 ± 1.47	<0.001
The simulation tasks I have performed today will improve my intra-operative performance in mastoid surgery	7.94 ± 1.39	<0.001
The simulation tasks I have performed today will improve my understanding of the anatomy of the temporal bone	8.72 ± 1.13	<0.001

\*1 = strongly disagree; 10 = strongly agree

advance in the competition. Elsewhere, the introduction of competitive game mechanics through an elimination tournament for urology residents greatly improved engagement with simulator-based training outside of the tournament.<sup>28</sup>

Virtual reality temporal bone competitions may be of particular educational benefit for more junior trainees. Eight candidates (44 per cent) had not performed any cortical mastoidectomies previously. For such trainees, an interactive virtual reality interface can help address some of the most challenging aspects of temporal bone anatomy in a safe environment. Novices were faster at identifying anatomical landmarks and showed faster improvement when evaluating the three-dimensional virtual reality temporal bone compared with more traditional cross-sectional evaluation.<sup>29</sup>

In our study, seventeen trainees (94 per cent) reported improved anatomical understanding of the temporal bone and believed this to be useful for future temporal bone surgery. This is supported by a significant post-competition improvement in self-rated confidence regarding performing cortical mastoidectomy (mean from  $4.83 \pm 2.41$  to  $5.72 \pm 2.22$ , t =3.49, p = 0.00277). We expect that the feedback of experts after the drilling exercise highlighted specific development areas in trainees' techniques. Vygotsky described the importance of this social interaction in focusing learning in the zone of proximal development.<sup>30</sup> The competition format provides opportunities for verbal guidance, affirmation and evidence of performance attainments necessary for cultivating the concept of self-efficacy highlighted by Bandura.<sup>31</sup>

Trainees perceived the social aspect of the competition format to carry great value. Interaction with peers and faculty was cited as a particular highlight, making it not only 'more fun' overall, but also more educational. The competition format aligns closely with Lave and Wenger's concept of communities of practice.<sup>21</sup> These represent forums for trainees to discuss challenges, collaborate on problems and engage with experts.<sup>32</sup> For more junior trainees, such activities help to strengthen the sense of belonging in the ENT and regional trainee community. This aspect was particularly lacking during the Covid-19 pandemic. For more senior trainees, formative assessment gives a valuable opportunity for benchmarking against peers and reflection, and guides the direction of future learning or practice.<sup>33</sup>

Competitions can improve engagement with ENT simulation training in general. Despite a variety of ENT simulators being described in the literature,<sup>34</sup> only 4 trainees (22 per cent) had reported using other types of ENT simulation models previously in their training. This surprisingly low figure may be explained by recall bias or variations in individual definitions of 'simulation models'.

In the West Midlands, trainees are expected to attend at least one regional cadaveric temporal bone course per year. Recently, optional regional virtual reality temporal bone simulator courses have started to be offered to trainees in a newly established dry laboratories to supplement their existing training. It was encouraging that prior to the competition 16 trainees (89 per cent) had already used Voxel-Man TempoSurg simulator as part of this facility, spending an estimated mean of  $7.22 \pm 4.86$  hours in total on the simulator per trainee.

In our study, most participants (94 per cent) felt competitions would be a useful addition to the ENT training programme, although most (67 per cent) felt they should not be mandated. There is much appetite for virtual reality temporal bone competitions given that almost all the trainees (94 per cent) stated they would attend a similar one in future.

We recommend surgical simulation competitions to complement traditional simulation training techniques, promote social learning opportunities and reward achievement. However, it must be recognised that competitions in isolation cannot be expected to improve trainee engagement in simulation.<sup>35</sup> Newer methods of training should incorporate suitable technologies that reflect the technology literacy of newer generations.<sup>32,36</sup> Combining competition and simulation technologies is especially suited to improve engagement with challenging topics such as anatomy and histology,<sup>18</sup> and has already been used in several surgical specialties including general surgery,<sup>35</sup> cardiothoracics<sup>27</sup> and urology.<sup>28</sup>

ENT competitions, especially those utilising virtual reality technology, such as the one reported here, have a potential role in introducing and improving operative knowledge and skill acquisition through the uptake of simulation training at a time when the Covid-19 pandemic limited many clinical opportunities.<sup>37</sup>

#### Limitations

While all West Midlands trainees were invited to take part in the competition, only 18 of the 47 trainees attended. These trainees are likely to have already formed overall positive opinions regarding simulation, competitions or both, reflecting a positive selection bias. Like any training tool or method, it must be recognised that different trainees will have different perceptions and preferences towards their learning. Individual trainees are likely to accrue different benefits from a virtual reality temporal bone competition. This competition, however, represents the largest virtual reality temporal bone competition in the published literature to date. Although only West Midlands trainees were included, we do not believe this limits the generalisability of our conclusions.

#### Conclusion

ENT trainees had an overall positive perception towards the educational benefits of a virtual reality temporal bone competition, believing the competition to be educationally valuable, fun and social. Whilst trainees have a healthy appetite for such competitions, it is generally felt they should not be mandated.

From the trainers' perspective, we encourage the use of competitions, particularly using virtual reality temporal bone

simulation model, as a supplement to traditional ENT training to engage trainees in simulation, stimulate learning, provide opportunities for social learning and networking, and recognise trainee achievements.

- Detailed knowledge of the anatomy of the temporal bone is essential for surgical management of complicated mastoiditis
- Newer methods of practice, such as virtual reality simulation, allow trainees to practice temporal bone drilling in a safe and controlled environment, complementing existing training
- The competition format can transform drilling simulation into an engaging and social learning activity, but is relatively underutilised in ENT
- ENT trainees have a positive perception of the educational benefits of virtual reality temporal bone competitions, believing them to be educational, fun and social
- Whilst trainees have a healthy appetite for such competitions, it is generally felt they should not be mandated within training programmes

**Supplementary material.** The supplementary material for this article can be found at https://doi.org/10.1017/S0022215124000070

#### Competing interests. None declared

#### References

- 1 Greenaway D. Shape of Training: Securing the Future of Excellent Patient Care. In: https://www.gmc-uk.org/-/media/documents/Shape\_of\_training\_ FINAL\_Report.pdf\_53977887.pdf
- 2 Migirov L, Yakirevitch A, Kronenberg J. Mastoid subperiosteal abscess: a review of 51 cases. Int J Pediatr Otorhinolaryngol 2005;69:1529–33
- 3 Psarommatis I, Giannakopoulos P, Theodorou E, Voudouris C, Carabinos C, Tsakanikos M. Mastoid subperiosteal abscess in children: drainage or mastoidectomy? J Laryngol Otol 2012;126:1204–8
- 4 Taylor MF, Berkowitz RG. Indications for mastoidectomy in acute mastoiditis in children. *Ann Otol Rhinol Laryngol* 2004;**113**:69–72
- 5 Davis J, Spraggs P, Murray C. Otolaryngology Curriculum. The Intercollegiate Surgical Curriculum Programme 2021. August 2021. https://www.iscp.ac.uk/ media/1106/otolaryngology-curriculum-aug-2021-approved-oct-20.pdf
- 6 Calman K, Temple J, Naysmith R, Cairncross R, Bennett S. Reforming higher specialist training in the United Kingdom: a step along the continuum of medical education. *Med Educ* 1999;33:28–33
- 7 Benes V. The European working time directive and the effects on training of surgical specialists (doctors in training). *Acta Neurochir* (*Wien*) 2006;**148**:1020–6
- 8 Neville E. Modernising medical careers. Clin Med (Lond) 2003;3:529
- 9 Mick P, Murphy R. Aerosol-generating otolaryngology procedures and the need for enhanced PPE during the COVID-19 pandemic: a literature review. J Otolaryngol Head Neck Surg 2020;49:1–10
- 10 Lion P, McClenaghan F, Hall A, Mackinnon S, Navaratnam A. ENT trainees' experience of redeployment during the coronavirus disease 2019 pandemic: a qualitative study. J Laryngol Otol 2021;135:391–5
- 11 George A, De R. Review of temporal bone dissection teaching: how it was, is and will be. J Laryngol Otol 2010;124:119–25
- 12 Arora A, Khemani S, Tolley N, Singh A, Budge J, Varela DADV et al. Face and content validation of a virtual reality temporal bone simulator. *Otolaryngol Head Neck Surg* 2012;**146**:497–503
- 13 Nash R, Sykes R, Majithia A, Arora A, Singh A, Khemani S. Objective assessment of learning curves for the Voxel-Man TempoSurg temporal bone surgery computer simulator. *J Laryngol Otol* 2012;**126**:663–9

- 14 Aussedat C, Venail F, Marx M, Boullaud L, Bakhos D. Training in temporal bone drilling. *Eur Ann Otorhinolaryngol Head Neck Dis* 2022;**139**:140–5
- 15 Ericsson KA, Krampe RT, Tesch-Römer C. The role of deliberate practice in the acquisition of expert performance. *Psychol Rev* 1993;100:363
- 16 Hashimoto DA, Gomez ED, Beyer-Berjot L, Khajuria A, Williams NN, Darzi A et al. A randomized controlled trial to assess the effects of competition on the development of laparoscopic surgical skills. J Surg Educ 2015;72:1077–84
- 17 Leraas HJ, Cox ML, Bendersky VA, Sprinkle SS, Gilmore BF, Gunasingha RM et al. Instituting a surgical skills competition increases technical performance of surgical clerkship students over time. J Surg Educ 2018;75:644–9
- 18 Janssen A, Shaw T, Goodyear P, Kerfoot BP, Bryce D. A little healthy competition: using mixed methods to pilot a team-based digital game for boosting medical student engagement with anatomy and histology content. BMC Med Educ 2015;15:1–10
- 19 Killick N, Amnolsingh R, Chu M. 1st Core Trainees Virtual Temporal Bone Competition ENT and Audiology News. Edinburgh, Scotland: Pinpoint Scotland Ltd, 2019;28:20
- 20 Ford K, Cooper L. The Surgical Skills competition. *Bull R Coll Surg Engl* 2022;**104**:14–17
- 21 Wenger E. Communities of practice and social learning systems. Organization 2000;7:225-46
- 22 Andersen SA, Cayé-Thomasen P, Sørensen MS. Mastoidectomy performance assessment of virtual simulation training using final-product analysis. *Laryngoscope* 2015;**125**:431–5
- 23 Francis HW, Niparko JK. Temporal Bone Dissection Guide. New York: Thieme, 2016
- 24 R Core Team. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing, 2021
- 25 McGaghie WC, Issenberg SB, Petrusa ER, Scalese RJ. A critical review of simulation-based medical education research: 2003–2009. *Med Educ* 2010;44:50–63
- 26 Knowles MS, Holton III EF, Swanson RA, Swanson R, Robinson PA. The adult learner: the definitive classic in adult education and human resource development. London: Routledge, 2020
- 27 Mokadam NA, Lee R, Vaporciyan AA, Walker JD, Cerfolio RJ, Hermsen JL et al. Gamification in thoracic surgical education: using competition to fuel performance. J Thorac Cardiovasc Surg 2015;150:1052–58
- 28 Kerfoot BP, Kissane N. The use of gamification to boost residents' engagement in simulation training. JAMA Surg 2014;149:1208–9
- 29 Timonen T, Dietz A, Linder P, Lehtimäki A, Löppönen H, Elomaa A-P *et al.* The effect of virtual reality on temporal bone anatomy evaluation and performance. *Eur Arch Otorhinolaryngol* 2021;**279**:4303–12
- 30 Vygotsky LS, Cole M. Mind in Society: Development of Higher Psychological Processes. Harvard: Harvard University Press, 1978
- 31 Bandura A. Social Foundations of Thought and Action. Englewood Cliffs, NJ: Prentice Hall, 1986;23–8
- 32 McCoy L, Lewis JH, Dalton D. Gamification and multimedia for medical education: a landscape review. J Am Osteopath Assoc 2016;116:22–34
- 33 Epstein RM. Assessment in medical education. N Engl J Med 2007;**356**:387–96
- 34 Pankhania R, Pelly T, Bowyer H, Shanmugathas N, Wali A. A systematic review of low-cost simulators in ENT surgery. J Laryngol Otol 2021;135:486–91
- 35 McCreery GL, El-Beheiry M, Schlachta CM. Local and national laparoscopic skill competitions: residents' opinions and impact on adoption of simulation-based training. *Surg Endosc* 2017;**31**:4711–16
- 36 Greenhalgh T. Computer assisted learning in undergraduate medical education. *BMJ* 2001;**322**:40-4
- 37 Hope C, Reilly JJ, Griffiths G, Lund J, Humes D. The impact of COVID-19 on surgical training: a systematic review. *Tech Coloproctol* 2021;25:505–20