

Original Article

*Yi-Chun Liu and Vincent Chin-Hung Chen contributed equally to this work as the joint first authors.

Cite this article: Liu Y-C, Chen VC-H, Yang Y-H, Chen Y-L, Gossop M (2021). Association of psychiatric comorbidities with the risk of transport accidents in ADHD and MPH. *Epidemiology and Psychiatric Sciences* **30**, e14, 1–9. <https://doi.org/10.1017/S2045796021000032>

Received: 27 July 2020

Revised: 16 December 2020

Accepted: 18 January 2021

Key words:

Attention-deficit/hyperactivity disorder; methylphenidate; Taiwan national insurance; transport accidents

Author for correspondence:

Yi-Lung Chen, E-mail: elong@asia.edu.tw

Association of psychiatric comorbidities with the risk of transport accidents in ADHD and MPH

Yi-Chun Liu^{1,2,3,*}, Vincent Chin-Hung Chen^{4,5,*}, Yao-Hsu Yang^{6,7},
Yi-Lung Chen^{8,9}  and Michael Gossop¹⁰

¹Department of Psychiatry, Changhua Christian Children's Hospital, Changhua, Taiwan; ²Department of Psychiatry, Changhua Christian Hospital, Changhua, Taiwan; ³Department of Eldercare, Central Taiwan University of Science and Technology, Taichung, Taiwan; ⁴School of Medicine, Chang Gung University, Taoyuan, Taiwan; ⁵Department of Psychiatry, Chiayi Chang Gung Memorial Hospital, Chiayi, Taiwan; ⁶Health Information and Epidemiology Laboratory, Chang Gung Memorial Hospital, Chiayi, Taiwan; ⁷Department of Traditional Chinese Medicine, Chang Gung Memorial Hospital, Chiayi Branch, Chiayi, Taiwan; ⁸Department of Healthcare Administration, Asia University, Taichung, Taiwan; ⁹Department of Psychology, Asia University, Taichung, Taiwan and ¹⁰National Addiction Centre, Institute of Psychiatry, King's College London, London, UK

Abstract

Aims. Although the relationship between attention-deficit/hyperactivity disorder (ADHD) and transport accidents has been shown, there is limited information on the relationship between medication and dose–response effects and transport accident risk. This study aims to determine whether young people with ADHD, including adolescents, are more prone to transport accidents than those without, and the extent to which methylphenidate (MPH) prescription in these patients reduces the risk.

Methods. We identified 114 486 patients diagnosed with ADHD from Taiwan's National Health Insurance Research Database from 1997 to 2013. Using a Cox regression model, we compared the risk of transport accidents between ADHD and non-ADHD groups and estimated the effect of MPH on accidents. Furthermore, we applied a self-control case-series analysis to compare the risk of accidents during the medication periods with the same patients' non-medication periods.

Results. Male ADHD patients had a higher risk of transport accidents than non-ADHD individuals (adjusted hazard ratio [aHR] = 1.24, [95% confidence interval (CI) 1.10–1.39]), especially for those comorbid with epilepsy, oppositional defiant disorder/conduct disorder (ODD/CD), and intellectual disabilities (ID). Female ADHD patients showed no relationship, except for comorbid with autism spectrum disorder (ASD) or ID. We found a reduced risk of transport accidents in patients with ADHD with MPH medication than those without MPH, with a plausible dose–response relationship (aHR of 0.23 to 0.07). A similar pattern was found in self-controlled case-series analysis.

Conclusions. Male patients with ADHD, especially those comorbid with epilepsy, ODD/CD, or ID, were at high risk of transport accidents. Female patients, when comorbid with ASD or ID, also exhibited a higher risk of accidents. MPH treatment lowered the accident risk with a dose–response relationship.

Introduction

As is known, attention-deficit/hyperactivity disorder (ADHD) is a common neurodevelopmental disorder characterised by persistent attention deficit, hyperactivity or impulsivity across settings (Biederman and Faraone, 2005). The onset typically occurs in childhood, and many symptoms often persist into young adulthood (Barkley *et al.*, 2002). Symptoms of ADHD can cause a wide range of functional impairment in daily life, including executive function impairment (Holst and Thorell, 2019). It is demonstrated that people with ADHD are prone to injuries and accidents, such as burns (Yeh *et al.*, 2020), fractures (Mangus *et al.*, 2004), concussions (Nelson *et al.*, 2016) and traumatic brain injuries (Rowe *et al.*, 2004; Liao *et al.*, 2018). A few studies have revealed the association between ADHD and traffic accidents (Jerome *et al.*, 2006; Barkley and Cox, 2007; Vaa, 2014). The association is possibly mediated by the core symptoms of ADHD, risky driving patterns, aggressiveness and comorbidities (e.g. substance use disorder and oppositional defiant disorder/conduct disorder [ODD/CD]) (Barkley and Cox, 2007; Vaa, 2014; Romo *et al.*, 2019). However, the above studies' results are limited because of lack of objective measures, small sample sizes, self- or parent-reported driving performance with long recall periods, referral bias and absence of adjustment for critical comorbid illnesses. One Swedish population-based cohort study by Chang *et al.* reported that patients with ADHD had an increased risk of severe traffic accidents

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article.

compared to those without ADHD (males: adjusted hazard ratio [aHR] = 1.47 [95% confidence interval (CI) 1.32–1.63]; females: aHR = 1.45 [95% CI 1.24–1.71]) (Chang *et al.*, 2014). However, their study group was restricted to adults. Adolescents tend to have more road safety problems, including risky driving behaviours, traffic violations and driving without a license (Dobhal *et al.*, 2019), especially for individuals with ADHD (Curry *et al.*, 2017; Curry *et al.*, 2019). Therefore, it is reasonable to include the adolescent population in the long-term study to explore the association between ADHD and vehicle accidents.

Transport accident is a major public health issue and one of the leading causes of physical disability and death for young people worldwide (Toroyan *et al.*, 2013). There is emerging evidence that medication used for ADHD not only alleviates core symptoms but also has a protective effect against trauma and accidental injuries (Tai *et al.*, 2013; Man *et al.*, 2015). Stimulants are considered the first choice for treating ADHD and are the most effective treatment (Cortese *et al.*, 2018). In Taiwan, methylphenidate (MPH) is the only stimulant approved for ADHD treatment. Some studies reported that MPH could improve driving performance in ADHD patients (Cox *et al.*, 2004; Barkley and Cox, 2007). However, these studies' effect sizes were small, and most of them were tested in driving simulators or specific driving tests, not in the real world. Two population-based cohort studies, one in Sweden and then another in the United States, conducted by Chang *et al.*, reported that medication treatment for ADHD, regardless of cumulative dose, significantly reduced the risk for severe traffic accidents in these patients (Chang *et al.*, 2014; Chang *et al.*, 2017). In the real world, medication discontinuation or non-adherence is common in patients with ADHD (Jensen *et al.*, 2007; Gau *et al.*, 2008). Whether the duration of medication therapy plays a moderator role in the risk of injuries remains uncertain. One report showed protective effects within 1 month (Raman *et al.*, 2013), but another reported at least 6 months (Chen *et al.*, 2017). Hence, it is vital to examine the impact of different cumulative doses of MPH on transport accident risk.

In the current study, we used a nationwide population-based cohort study to investigate the association of psychiatric comorbidities with the risk of transport accidents in ADHD and MPH. We hypothesised that young people with ADHD are prone to transport accidents than those without, and using MPH in these patients reduces the risk. We also hypothesised that the risk reduction becomes more prominent as the cumulative dose of MPH increases.

Materials and methods

Data source

Taiwan established the National Health Insurance (NHI) programme, a single-payer compulsory insurance system, on 1 March 1995. Since its launch in 1995, the coverage of the NHI programme has steadily increased. Up to 2009, the coverage rose to 99.5% of the national population in Taiwan (Ho Chan, 2010). The Bureau of NHI derived all medical claims data from NHI beneficiaries to form the National Health Insurance Research Database (NHIRD), which comprised of the records of hospital inpatient and outpatient care, dental services, ambulatory care and medication prescription. NHIRD utilised a systematic random sampling to provide a representative research sample of the national population, named Longitudinal Health Insurance

Database (LHID), which contains medical claims of 1 000 000 people from all NHI beneficiaries. There are no significant differences in sex, age or health care utilisation between this sample and all beneficiaries (NHR, 2013).

Ethics

The Ethics Institutional Review Board of Chang Gung Memorial Hospital had reviewed and approved this study.

Sample

This is a population-based retrospective cohort study. We had two research hypotheses: ADHD young patients have an elevated risk of transport accidents, and MPH can reduce the risk of transport accidents with different doses. To test the first hypothesis, we compared the transport accident incidence between ADHD cases and non-ADHD cases. For the ADHD group, we identified ADHD cases according to the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code 314 from the NHIRD. In Taiwan, clinicians used the ICD-9-CM code 314 as an insurance claim code for ADHD in the DSM-IV because of the NHI registry system (Tai and Chiu, 2009). ADHD cases enrolled in this study required at least one inpatient diagnosis or more than two outpatient diagnoses within 1 year during the observation period from 1 January 1997 to 31 December 2012, and followed up to the end of 2013. The flow chart of participant selection was presented in Fig. 1. At first, we identified 114 486 diagnoses of ADHD within the observation period after excluding those who had unknown sex status, less than 12 years old, and older than 35 years old (age was calculated as the difference from their birthdate to the end of follow-up, 31 December 2013). On the other hand, we enrolled 338 261 non-ADHD participants aged between 12 and 35 years old from the LHID 2005 database from NHIRD, which composed of 1 million beneficiaries randomly sampled in the year 2005 (NHR, 2013). We followed up these participants from 1997 to 2013.

To test the second hypothesis, we further used the 114 186 identified ADHD cases and classified them into MPH users ($n = 89\,826$) and non-users ($n = 24\,660$). Furthermore, to eliminate possible indication bias (i.e. patients with ADHD at higher risk of transport accidents may be more likely to be prescribed MPH), we performed within-person comparisons using the self-controlled case series (SCCS) model. In this model, participants with ADHD who had ever taken MPH and had a history of transport accidents were enrolled, and each participant served as his or her own control. We compared the rate of events during medication and non-medication periods.

Medication

In Taiwan, MPH is the only stimulant approved for ADHD. MPH included in our study was N06BA04 according to the Anatomical Therapeutic Chemical (ATC) code. Atomoxetine, which launched in Taiwan in 2007, is a non-stimulant approved for ADHD treatment. Before 2017, MPH had remained the first-line treatment according to the stipulation of the Bureau of NHI. Only 4% of all ADHD patients in the database were prescribed with ATX (Lee *et al.*, 2016). Hence, we confined our analysis to those with MPH prescriptions.




Fig. 1. Flow chart for participants enrolled in this cohort study. ADHD, attention-deficit/hyperactivity disorder; LHID, longitudinal national health insurance research database; NHIRD, National Health Insurance Research Database.

Transport accidents

The main outcome of our study was a transport accident. We identified the transport accidents according to the ICD-9-CM, code E810–E825. We further categorised the transport accidents into traffic accidents (ICD-9 codes: E810–E819) and non-traffic accidents (ICD-9 codes: E820–E825). Patients with ADHD were followed up for transport accidents as an outcome to the end of 2013.

Covariates

Several covariates were selected, including sex, age, and psychiatric comorbidity. Psychiatric comorbidity included autism spectrum disorder (ICD-9 code: 299), tic disorders (ICD-9 code: 307.2), epilepsy (ICD-9 code: 345), conduct disorder (ICD-9 code: 312), opposition defiant disorder (ICD-9-CM code: 313.81), anxiety disorders (ICD-9 code: 300), major depressive disorders (ICD-9 codes: 296.2, 296.3, 300.4 and 311), bipolar disorders (ICD-9 codes: 296.0, 296.1, 296.4–296.8 and 301.13) and schizophrenia (ICD-9 code: 295).

Statistical analysis

The statistical analyses were performed by using SAS Version 9.4 (SAS Institute Inc., Cary, NC, USA) and R 3.6.2 (R Foundation for Statistical Computing, Vienna, Austria). We used a chi-square (χ^2) test to compare the characteristics between the ADHD and non-ADHD groups. In response to two research hypotheses: ADHD young patients have an elevated risk of transport accidents, and MPH can reduce the risk of transport accidents with different doses, we performed two different analyses: between-subjects comparison and within-subjects comparison. There were two parts of the between-subjects comparison. First, by using a Cox proportional hazards regression model with adjustment for demographics and psychiatric comorbidities, we compared the risk of transport accidents between ADHD groups with and without each specific comorbidity and non-ADHD group. Analyses were stratified by sex and age category to examine the sex- and age-specific associations of transport accidents. Based on Korn and colleagues' suggestion, age was used as the time-scale in the survival analysis (Korn *et al.*, 1997). The results were further presented as aHRs with 95% CIs. Second, among individuals

with ADHD, we compared the risk of transport accidents between MPH users with and without each specific comorbidity and non-users. We categorised these MPH users into three subgroups based on their cumulative defined daily dose (DDD): (1) 0, (2) >0 to <180 (0–180) and (3) \geq 180, and the group of 0 of DDD served as the control group in the analysis. The WHO proposed the DDD as a unit to measure drug exposure and recommended 30 mg per day to be one DDD of MPH. In clinical practice, if the patient takes medication continuously every day, 180 DDD stands for 180 days of refill prescription. As previous studies showed, the protective effect of MPH on the risk of suicide or fractures can be found in patients with prescriptions of 180 days or longer (Chen *et al.*, 2017; Liang *et al.*, 2018). Also, most ADHD patients received medication for less than 180 days (Chen *et al.*, 2011; Garbe *et al.*, 2012). In order to determine the effect of dose on risk more precisely, we chose 180 DDD as the cut-point while categorising. We also estimated the preventable fraction: the proportion of transport accidents in patients with ADHD not taking MPH that could be prevented by MPH medication, calculated as follows: $1 - \text{adjusted hazard ratio}$

For within-patient comparisons, we compared the risk of transport accidents between the periods of medication and non-medication within MPH users by using the SCCS model (Whitaker *et al.*, 2006). Personal MPH records, including prescription date and prescribed days, were extracted throughout the entire study period (by the end of 2013). The results were presented as relative incidences (RRs) with 95% CIs. The RR estimated in the SCCS model was obtained by comparing the rate of transport accidents during the time periods of individuals having MPH with that during all other non-medication periods. The SCCS model automatically adjusts for all time-invariant factors (e.g. sex and genetic factors) and possible indication bias for the same patient before and during the follow-up. We defined the effective period of MPH exposure as 0–90 days after the prescription of the drug (Man *et al.*, 2017; Hollis *et al.*, 2019). We further divided the effective period into three 1-month effective periods: 0–30, 31–60 and 61–90 days after MPH initiation (Shin *et al.*, 2016). Finally, we merged the three effect periods into one period to evaluate the average pooled estimate of MPH use for the transport accidents risk. If patients had drug exposure continuously, we would record consecutive exposure effective periods. For example, if an individual was taking two 30-day supply of prescription successively, the effective periods will be 0–30, 0–30, 31–60 and 61–90 days.

Outcomes

Characteristics of subjects with or without ADHD

The study sample comprised of 114 486 ADHD patients and 338 261 non-ADHD participants. Table 1 lists the characteristics of ADHD patients and non-ADHD participants. Compared to the non-ADHD group, the ADHD group was more likely to be male, younger and comorbid with other psychiatric disorders ($p < 0.001$). Although the risk of transport accidents was 0.6% for the ADHD group and 1.0% for the non-ADHD, the ADHD group was at a younger age at the time of transport accidents ($p < 0.001$). Regardless of the ADHD group or the non-ADHD group, most transport accidents were traffic accidents (97.9%, 662/676 for ADHD and 98.8%, 3213/3252 for non-ADHD, respectively).

Comparisons of the transport accidents risk between ADHD and non-ADHD

The risk of transport accidents between the ADHD group with and without each specific comorbidity and the non-ADHD group was presented in Table 2. We found that male ADHD cases had a higher risk of transport accidents than the non-ADHD group, with an aHR of 1.24 and 95% CI 1.10–1.39. The risk was noticeable when male ADHD patients were comorbid with epilepsy, ODD/CD and intellectual disability (ID). In the stratification analysis, we found the risks of transport accidents were less profound in female or adult individuals with ADHD. Female individuals with ADHD were only found to have higher risks of transport accidents when comorbid with autism spectrum disorder (ASD) or ID. In addition, adult individuals with ADHD were not associated with a higher risk of transport accidents regardless of psychiatric comorbidity.

Comparisons of the effect of MPH on transport accidents risk in ADHD

Table 3 presents the dose-effect of MPH medication on transport accidents risk in patients with ADHD. Based on the Cox proportional hazard regression, MPH was associated with a reduced risk for transport accidents, and a significant dose-response relationship was observed. Compared to those without MPH use, the ADHD group with 0–180 DDDs had aHR of 0.23 (95% CI 0.19–0.26), and the group with more than 180 DDDs had aHR of 0.07 (95% CI 0.01–0.52). The preventable fractions of MPH medication were 0.77 and 0.93, indicating that 77 or 93% of transport accidents could be prevented if they took an equal dose of MPH medication (i.e. 0–180 DDD and \geq 180 DDD, respectively). Furthermore, the effect of MPH was also significant in ADHD patients comorbid with different neurodevelopment and psychiatric disorders (Table 4).

For within-comparison, we conducted a conditional Poisson regression model for the SCCS cohort study (Table 5). Although the average risk during the first 3 months after using MPH was reduced (RR = 0.85), the effect did not reach significance level (95% CI 0.57–1.25). We found a potential beneficial effect of MPH on transport accidents, which seemed to be higher during the second to third month after using MPH (RR = 0.35, 95% CI 0.11–1.11), although not statistically significant.

Discussion

Our study found that male patients with ADHD had an increased risk of transport accidents, especially for those comorbid with epilepsy, ODD/CD or ID. Female ADHD patients comorbid with ASD and ID were also at risk of accidents. For ADHD patients with or without other psychiatric disorders, MPH prescription reduced the risk of transport accidents, and the protective effect was dose-related. At doses <180 DDD and \geq 180 DDD, MPH showed a significantly reduced risk for transport accidents by 77 and 93%, respectively.

Previous studies had reported that patients with ADHD tend to be reckless or inattentive during driving, which could attribute to traffic accidents (Barkley and Cox, 2007). A population-based cohort study in Sweden conducted by Chang *et al.* showed that, regardless of gender, ADHD patients had a 42 to 47% increase in transport accident risk compared to people without ADHD (Chang *et al.*, 2014). The total follow-up time was 4 years

Table 1. Demographic characteristics of ADHD and non-ADHD youths and young adults, aged between 12 and 35 years

Characteristics	ADHD <i>n</i> = 114 486	Non-ADHD <i>N</i> = 338 261	<i>p</i> -value
Sex, <i>n</i> (%)			
Male	90 297 (78.9%)	164 860 (48.7%)	<0.001
Female	24 189 (21.1%)	173 401 (51.3%)	
Age, mean (s.d.), years	17.2 (4.30)	24.4 (6.95)	<0.001
Psychiatric comorbidity			
ASD, <i>n</i> (%)	12 770 (11.2%)	576 (0.2%)	<0.001
Tic disorders, <i>n</i> (%)	7569 (6.6%)	1386 (0.4%)	<0.001
Epilepsy, <i>n</i> (%)	6795 (5.9%)	4790 (1.4%)	<0.001
ODD/CD, <i>n</i> (%)	13 989 (12.2%)	643 (0.2%)	<0.001
Anxiety disorders, <i>n</i> (%)	37 841 (33.1%)	36 837 (10.9%)	<0.001
Major depressive disorder, <i>n</i> (%)	13 022 (11.4%)	13 911 (4.1%)	<0.001
Bipolar disorders, <i>n</i> (%)	4011 (3.5%)	2246 (0.7%)	<0.001
ID, <i>n</i> (%)	18 807 (16.4%)	2849 (0.8%)	<0.001
Schizophrenia, <i>n</i> (%)	2326 (2.0%)	1879 (0.6%)	<0.001
Any psychiatric disorder, <i>n</i> (%)	70 929 (62.0%)	46 849 (13.8%)	<0.001
Transport accidents, <i>n</i> (%)	676 (0.6%)	3252 (1.0%)	<0.001
Traffic accidents, <i>n</i> (%)	662 (0.6%)	3213 (0.9%)	<0.001
Non traffic accidents, <i>n</i> (%)	14 (>0.1%)	39 (>0.1%)	<0.001
Age of transport accidents, mean (s.d.), years	17.7 (4.3)	24.8 (7.0)	<0.001
Use of MPH, <i>n</i> (%)	89 826 (78.5)	–	<0.001

ADHD, attention-deficit/hyperactivity disorder; ASD, autism spectrum disorder; CD, conduct disorder; ID, intellectual disability; MPH, methylphenidate; ODD, oppositional defiant disorder.

Table 2. Cox proportional hazard regression model analysis for risk of transport accidents and psychiatric comorbid disorders between ADHD and non-ADHD, stratified by sex and age

Group	All aHR (95% CI) ^a	Sex		Age	
		Female ^b	Male ^b	<18 ^c	≥18 ^c
Only ADHD	1.19 (1.08–1.32)***	1.01 (0.81–1.27)	1.24 (1.10–1.39)***	1.06 (0.86–1.31)	1.02 (0.90–1.15)
ADHD plus ASD	1.09 (0.82–1.47)	2.13 (1.17–3.87)***	0.93 (0.66–1.31)	1.16 (0.84–1.59)	0.87 (0.74–1.02)
ADHD plus tic disorders	1.26 (0.89–1.79)	1.74 (0.63–4.82)	1.21 (0.83–1.76)	0.93 (0.57–1.52)	0.76 (0.50–1.15)
ADHD plus epilepsy	1.76 (1.29–2.41)***	1.74 (0.91–3.32)	1.78 (1.25–2.55)***	1.37 (0.82–2.30)	0.60 (0.32–1.09)
ADHD plus ODD/CD	1.55 (1.25–1.93)***	0.87 (0.43–1.78)	1.68 (1.34–2.12)***	1.97 (1.14–3.40)*	1.28 (0.85–1.93)
ADHD plus anxiety disorders	1.00 (0.85–1.17)	0.90 (0.65–1.24)	1.02 (0.85–1.23)	1.42 (0.96–2.10)	1.20 (0.91–1.60)
ADHD plus major depressive disorder	0.87 (0.70–1.08)	0.64 (0.42–0.97)*	1.01 (0.78–1.31)	0.91 (0.37–2.20)	0.86 (0.69–1.08)
ADHD plus bipolar disorders	1.04 (0.74–1.47)	0.80 (0.40–1.60)	1.17 (0.79–1.75)	1.10 (0.31–3.94)	1.03 (0.72–1.47)
ADHD plus ID	1.51 (1.22–1.85)***	1.57 (1.05–2.37)*	1.49 (1.17–1.90)***	1.47 (1.03–2.10)*	1.14 (0.87–1.50)
ADHD plus schizophrenia	1.03 (0.67–1.59)	1.10 (0.48–2.55)	1.03 (0.62–1.71)	2.12 (0.43–10.36)	1.01 (0.65–1.58)
ADHD plus any psychiatric disorder	1.07 (0.94–1.23)	1.31 (0.90–1.91)	1.19 (1.01–1.40)*	1.22 (0.88–1.71)	0.90 (0.77–1.06)

ADHD, attention-deficit/hyperactivity disorder; ASD, autism spectrum disorder; CD, conduct disorder; ID, intellectual disability; ODD, oppositional defiant disorder.

^aAnalysis was adjusted for age, sex and psychiatric comorbidity.

^bAnalysis was adjusted for age and psychiatric comorbidity.

^cAnalysis was adjusted for sex and psychiatric comorbidity.

The non-ADHD served as the reference group.

Table 3. Dose–response analysis of use of MPH of on transport accidents in ADHD youths using the Cox proportional hazard regression

Group	N	Transport accidents		Preventable fraction
		aHR (95% CI) ^a	p-value	
Only ADHD				
Non-user ^a	24 660	1.00		
0 DDD–180 DDD	89 172	0.23 (0.19–0.26)	<0.001	0.77
≥180 DDD	654	0.07 (0.01–0.52)	0.009	0.93

ADHD, attention-deficit/hyperactivity disorder.

^aAnalysis was adjusted for age, sex and psychiatric comorbidity.

Preventable fraction is calculated as 1 minus relative risk.

(Chang *et al.*, 2014). As in previous studies, our results supported that patients with ADHD had a higher risk of transport accidents after adjusting sex and psychiatric comorbidities (aHR = 1.19 [95% CI 1.08–1.32]). In our study, we not only extended the follow-up time to 17 years but also enrolled cases under the age of 18, the legal minimum driving age in Taiwan. Based on Curry *et al.*'s retrospective cohort study in North America, ADHD adolescents drivers had 1.36 times (95% CI 1.25–1.48) higher risk of a car accident (Curry *et al.*, 2017). Also, as in many Asian countries, motorcycles are one of the primary means of vehicles in Taiwanese daily life (World Health Organization, 2015). Because of the ease of use of motorcycles and road traffic space limitations, unlicensed teenage motorcyclists and related severe traffic crashes are not uncommon in Taiwan (Chen *et al.*, 2018; World Health Organization, 2015). As in our stratification analyses, ADHD patients who were less than 18-year-old and concurrent with ODD/CD or ID were at major risk for transport accidents, which may due to their immature impulse control or impaired cognitive function.

ADHD is highly concurrent comorbid with other mental illnesses (Jensen and Steinhausen, 2015). In our study, male ADHD patients and specifically in those with epilepsy, ODD/CD or ID had increased risk of transport accidents, increased by 78, 68 and 49%, respectively. In Taiwan, the proportion and severity of traffic accidents for men are much higher than for women (Chang *et al.*, 2020). This may partially explain why only male patients were significantly associated with accidents in our study. In our study, female ADHD patients were only found to have higher risks of accidents when being comorbid with severe neurodevelopment disorders (i.e. ASD or ID). As past studies have shown, youths with destructive behaviour disorders or mental retardation often exhibit excessive impulsive or risky behaviours. Thompson *et al.* reported that ADHD concurrent with conduct problems was prone to risky driving in a case-control study (Thompson *et al.*, 2007). Savage *et al.* performed secondary analysis from the National Longitudinal Transition Study-2 (NLTS2) and reported that adolescents with mild ID were at greater risk for engagement in risky behaviours (Savage and Bouck, 2017). In addition, some studies have reported poor driving performance of autistic patients in virtual settings, some of which may be due to impaired executive function in complex situations (Chee *et al.*, 2019; Patrick *et al.*, 2020). Epilepsy alone can also contribute to traffic accidents as the results of one Swedish nationwide cohort study (Sundelin *et al.*, 2018).

Table 4. Cox proportional hazard regression model analysis of use of MPH of on transport accidents and psychiatric comorbid disorders in ADHD youths

Group	N	Transport accidents	
		aHR (95% CI) ^a	p-value
ADHD plus ASD	9190	0.17 (0.11–0.26)	<0.001
ADHD plus tic disorders	5870	0.13 (0.07–0.23)	<0.001
ADHD plus epilepsy	5039	0.30 (0.19–0.46)	<0.001
ADHD plus ODD/CD	11 858	0.29 (0.21–0.39)	<0.001
ADHD plus anxiety disorders	31 375	0.22 (0.18–0.27)	<0.001
ADHD plus major depressive disorder	9885	0.26 (0.19–0.35)	<0.001
ADHD plus bipolar disorders	3084	0.25 (0.15–0.42)	<0.001
ADHD plus ID	14 193	0.25 (0.19–0.34)	<0.001
ADHD plus schizophrenia	1609	0.42 (0.25–0.72)	0.002
ADHD plus any psychiatric disorder	55 898	0.22 (0.19–0.27)	<0.001

ADHD, attention-deficit/hyperactivity disorder; ASD, autism spectrum disorder; CD, conduct disorder; ID, intellectual disability; ODD, oppositional defiant disorder.

Analysis was adjusted for sex and psychiatric comorbidity.

^aIndividuals with ADHD did not take MPH medication or take less than 90 DDD; served as the reference group when the effect of MPH medication was examined in ADHD plus psychiatric comorbidities.

All analyses were adjusted for age, sex and psychiatric comorbidity.

Therefore, our results highlight the importance of paying attention to transport problems in patients with ADHD, especially among those who also have the above diseases.

For comorbidities between ADHD and ASD, despite the growing body of research pointing at the frequent co-occurrence of ADHD and ASD, the previous DSM-IV-TR has not allowed a dual diagnosis of ADHD and ASD. In 2013, the DSM-5 has modified the ADHD diagnostic criteria, allowing a co-morbid diagnosis of ADHD with ASD (Leitner, 2014). To address the co-morbid diagnosis of ADHD with ASD, we did not specify that an ASD diagnosis is an exclusion criterion for ADHD as recommended in the DSM-IV. In this study, individuals with ADHD can be identified as having ASD if they have received the diagnosis of ASD, and 11.2% of individuals with ADHD also had ASD, which was slightly lower than that in a recent meta-analytical review (21%) (Hollingdale *et al.*, 2020), but close to that in a Danish nationwide study (12.4%) (Jensen and Steinhausen, 2015). We found that a higher risk of transport accidents was only found in female individuals with ADHD who were comorbid with ASD, but not in males. It is possible that females referred for clinical evaluation and diagnosed with ASD might show more obvious impairments in emotional and behavioural problems than males with ASD (Frazier *et al.*, 2014).

In our study, the prescription of MPH was associated with a decreased risk of transport accidents. Our results also revealed that as MPH dosage increasing, there was a more apparent protective effect of MPH against transport accident risk. The protective effect of MPH against injuries was shown in previous studies (Tai *et al.*, 2013; Chen *et al.*, 2017). This effect may be possibly mediated by the alleviation of core symptoms and enhancement of executive function. One Swedish nationwide cohort study

Table 5. Conditional Poisson regression model, within comparison, for SCCS study design of transport accidents in ADHD youths with use of MPH

Group	Transport accidents	
	RR (95% CI)	p-value
Only ADHD		
0–90 days	0.85 (0.57–1.25)	0.396
0–30 days	1.05 (0.66–1.68)	0.835
30–60 days	0.86 (0.45–1.65)	0.644
60–90 days	0.35 (0.11–1.11)	0.074

This analysis was conducted in ADHD youths who had a history of MPH use and transport accidents visit.

RR was calculated by conditional Poisson regression, adjusted for all time-invariant covariates that are constant within each individual during the follow-up.

SCCS analysis for ADHD was conducted using effect period for MPH medication set at 0–90 days.

conducted by Chang *et al.* reported that ADHD medication reduced the risk of serious traffic accidents in adult male patients by 58% (HR = 0.42 [95% CI 0.23–0.75]), but not in female patients (Chang *et al.*, 2014). Another similar United States population cohort by Chang *et al.* found similar risk reduction in traffic accidents in both male and female patients by 38 and 42%, respectively. However, these studies did not investigate the dose–response effect. We also compared the effect of MPH on the events when ADHD was concurrent with other psychiatric comorbidities. Our results supported the prominent protective effect of MPH in the above situations. On the other hand, we used within-patient comparisons to control the unmeasured confounders consistent over time. The results showed that MPH had a potential protective trend in transport accidents within 3 months (RR = 0.85 [95% CI 0.57–1.25]) and the second to third month, although not significant (RR = 0.35 [95% CI 0.11–1.11]). This non-significant result could be partially explained by the fact that although this is a large-scale claim data, which contains over one hundred thousand patients with ADHD, the extremely low incidence of transport accidents (i.e. 0.6%) enlarged the confidence interval.

Overall, our study used a nationwide population database cohort to investigate the association of psychiatric comorbidities with the transport accidents in ADHD young people and MPH, with particular attention to the dose–response effects of MPH. We performed two parts of the analysis: between-subjects comparison and within-subject comparisons. All registration medical claim data came from the nationally representative sample of NHL, minimising the selection and recall bias. By excluding transport accidents before ADHD diagnosis, we have precluded the reverse association between ADHD and road traffic accidents as much as possible. The advantage of between-subjects comparison was that we were able to examine the MPH effect in different dose groups. However, confounding by indication cannot be eliminated. For example, those with a severe degree of ADHD symptoms, an exhibition of risky behaviours, or comorbid with other psychiatric illnesses were more likely to be prescribed medication. Hence, we also performed within-subject comparisons to adjust for time-invariant factors.

Some limitations affect the interpretation of our findings. First, there is a lack of several potential confounding factors such as adherence, lifestyle, alcohol use and subject bodyweight in our database. Also, we cannot attribute the cause of transport

accidents. ADHD patients might also be at risk of transport accidents due to poor executive function or judgement. As to the within-subject comparison, there were still some unmeasured time-varying confounding factors (e.g. parental and school supervision) that interfere with the results.

Conclusions

Our study found that male ADHD patients had a higher risk of transport accidents, especially when comorbid with epilepsy, ODD/CD or ID. Female ADHD patients exhibited elevated transport accident risk when comorbid with ASD or ID. Treatment with MPH lowered the risk of accidents in a dose–response relationship. These results highlight the safety road issues in the ADHD young population and the importance of treating ADHD.

Data. The current study was based on the National Health Insurance Research Database, managed by the National Health Research Institutes. This database was owned by the National Health Research Institutes.

Acknowledgements. None.

Financial support. This study was supported by grants from the Chang Gung Memorial Hospital (CLRPG6G0042 and CLRPG6G0043).

Conflict of interest. None.

Ethical standards. The authors declared that all procedures contributing to this study meet the ethical standards of the pertinent national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2000.

References

- Barkley RA and Cox D (2007) A review of driving risks and impairments associated with attention-deficit/hyperactivity disorder and the effects of stimulant medication on driving performance. *Journal of Safety Research* **38**, 113–128.
- Barkley RA, Fischer M, Smallish L and Fletcher K (2002) The persistence of attention-deficit/hyperactivity disorder into young adulthood as a function of reporting source and definition of disorder. *Journal of Abnormal Psychology* **111**, 279–289.
- Biederman J and Faraone SV (2005) Attention-deficit hyperactivity disorder. *Lancet (London, England)* **366**, 237–248.
- Chang Z, Lichtenstein P, D’Onofrio BM, Sjolander A and Larsson H (2014) Serious transport accidents in adults with attention-deficit/hyperactivity disorder and the effect of medication: a population-based study. *JAMA Psychiatry* **71**, 319–325.
- Chang Z, Quinn PD, Hur K, Gibbons RD, Sjolander A, Larsson H and D’Onofrio BM (2017) Association between medication use for attention-deficit/hyperactivity disorder and risk of motor vehicle crashes. *JAMA Psychiatry* **74**, 597–603.
- Chang YH, Li CY, Lu TH, Artanti KD and Hou WH (2020) Risk of injury and mortality among driver victims involved in single-vehicle crashes in Taiwan: comparisons between vehicle types. *International Journal of Environmental Research and Public Health* **17**, 1–9.
- Chee DYT, Lee HCY, Patomella AH and Falkmer T (2019) Investigating the driving performance of drivers with and without autism spectrum disorders under complex driving conditions. *Disability and Rehabilitation* **41**, 1–8.
- Chen CY, Yeh HH, Chen KH, Chang IS, Wu EC and Lin KM (2011) Differential effects of predictors on methylphenidate initiation and discontinuation among young people with newly diagnosed attention-deficit/hyperactivity disorder. *Journal of Child and Adolescent Psychopharmacology* **21**, 265–273.
- Chen VC, Yang YH, Liao YT, Kuo TY, Liang HY, Huang KY, Huang YC, Lee Y, McIntyre RS and Lin TC (2017) The association between

- methylphenidate treatment and the risk for fracture among young ADHD patients: a nationwide population-based study in Taiwan. *PLoS ONE* **12**, e0173762.
- Chen PL, Wang MH, Linkov V and Pai CW** (2018) Motorcycle door crashes: an evaluation of crash characteristics in Taipei city, Taiwan. *PLoS ONE* **13**, e0208016.
- Cortese S, Adamo N, Del Giovane C, Mohr-Jensen C, Hayes AJ, Carucci S, Atkinson LZ, Tessari L, Banaschewski T, Coghill D, Hollis C, Simonoff E, Zuddas A, Barbui C, Purgato M, Steinhausen HC, Shokraneh F, Xia J and Cipriani A** (2018) Comparative efficacy and tolerability of medications for attention-deficit hyperactivity disorder in children, adolescents, and adults: a systematic review and network meta-analysis. *The Lancet. Psychiatry* **5**, 727–738.
- Cox DJ, Humphrey JW, Merkel RL, Penberthy JK and Kovatchev B** (2004) Controlled-release methylphenidate improves attention during on-road driving by adolescents with attention-deficit/hyperactivity disorder. *The Journal of the American Board of Family Practice* **17**, 235–239.
- Curry AE, Metzger KB, Pfeiffer MR, Elliott MR, Winston FK and Power TJ** (2017) Motor vehicle crash risk among adolescents and young adults with attention-deficit/hyperactivity disorder. *JAMA Pediatrics* **171**, 756–763.
- Curry AE, Yerys BE, Metzger KB, Carey ME and Power TJ** (2019) Traffic crashes, violations, and suspensions among young drivers with ADHD. *Pediatrics* **143**, e20182305.
- Dobhal P, Dobhal A, Kashyap A and Bhadoria AS** (2019) Assessment of road traffic behavior using youth risk behavior survey questionnaire among school-going adolescents of Jaipur city, Rajasthan: an observational analysis. *Journal of Family Medicine and Primary Care* **8**, 3595–3599.
- Frazier TW, Georgiades S, Bishop SL and Hardan AY** (2014) Behavioral and cognitive characteristics of females and males with autism in the Simons simplex collection. *Journal of the American Academy of Child & Adolescent Psychiatry* **53**, 329–340, e323.
- Garbe E, Mikolajczyk RT, Banaschewski T, Petermann U, Petermann F, Kraut AA and Langner I** (2012) Drug treatment patterns of attention-deficit/hyperactivity disorder in children and adolescents in Germany: results from a large population-based cohort study. *Journal of Child and Adolescent Psychopharmacology* **22**, 452–458.
- Gau SS, Chen SJ, Chou WJ, Cheng H, Tang CS, Chang HL, Tzang RF, Wu YY, Huang YF, Chou MC, Liang HY, Hsu YC, Lu HH and Huang YS** (2008) National survey of adherence, efficacy, and side effects of methylphenidate in children with attention-deficit/hyperactivity disorder in Taiwan. *Journal of Clinical Psychiatry* **69**, 131–140.
- Ho Chan WS** (2010) Taiwan's healthcare report 2010. *The EPMA Journal* **1**, 563–585.
- Hollingdale J, Woodhouse E, Young S, Fridman A and Mandy W** (2020) Autistic spectrum disorder symptoms in children and adolescents with attention-deficit/hyperactivity disorder: a meta-analytical review. *Psychological Medicine* **50**, 2240–2253.
- Hollis C, Chen Q, Chang Z, Quinn PD, Viktorin A, Lichtenstein P, D'Onofrio B, Landen M and Larsson H** (2019) Methylphenidate and the risk of psychosis in adolescents and young adults: a population-based cohort study. *The Lancet. Psychiatry* **6**, 651–658.
- Holst Y and Thorell LB** (2019) Functional impairments among adults with ADHD: a comparison with adults with other psychiatric disorders and links to executive deficits. *Applied Neuropsychology. Adult*, 1–13.
- Jensen CM and Steinhausen HC** (2015) Comorbid mental disorders in children and adolescents with attention-deficit/hyperactivity disorder in a large nationwide study. *Attention Deficit and Hyperactivity Disorders* **7**, 27–38.
- Jensen PS, Arnold LE, Swanson JM, Vitiello B, Abikoff HB, Greenhill LL, Hechtman L, Hinshaw SP, Pelham WE, Wells KC, Conners CK, Elliott GR, Epstein JN, Hoza B, March JS, Molina BSG, Newcorn JH, Severe JB, Wigal T, Gibbons RD and Hur K** (2007) 3-year follow-up of the NIMH MTA study. *Journal of the American Academy of Child and Adolescent Psychiatry* **46**, 989–1002.
- Jerome L, Habinski L and Segal A** (2006) Attention-deficit/hyperactivity disorder (ADHD) and driving risk: a review of the literature and a methodological critique. *Current Psychiatry Reports* **8**, 416–426.
- Korn EL, Graubard BI and Midthune D** (1997) Time-to-event analysis of longitudinal follow-up of a survey: choice of the time-scale. *American Journal of Epidemiology* **145**, 72–80.
- Lee MJ, Yang KC, Shyu YC, Yuan SS, Yang CJ, Lee SY, Lee TL and Wang LJ** (2016) Attention-deficit hyperactivity disorder, its treatment with medication and the probability of developing a depressive disorder: a nationwide population-based study in Taiwan. *Journal of Affective Disorders* **189**, 110–117.
- Leitner Y** (2014) The co-occurrence of autism and attention deficit hyperactivity disorder in children – what do we know? *Frontiers in Human Neuroscience* **8**, 268.
- Liang SH, Yang YH, Kuo TY, Liao YT, Lin TC, Lee Y, McIntyre RS, Kelsen BA, Wang TN and Chen VC** (2018) Suicide risk reduction in youths with attention-deficit/hyperactivity disorder prescribed methylphenidate: a Taiwan nationwide population-based cohort study. *Research in Developmental Disabilities* **72**, 96–105.
- Liao Y-T, Yang Y-H, Kuo T-Y, Liang H-Y, Huang K-Y, Wang T-N, Lee Y, McIntyre RS and Chen VC-H** (2018) Dosage of methylphenidate and traumatic brain injury in ADHD: a population-based study in Taiwan. *European Child & Adolescent Psychiatry* **27**, 279–288.
- Man KK, Chan EW, Coghill D, Douglas I, Ip P, Leung LP, Tsui MS, Wong WH and Wong IC** (2015) Methylphenidate and the risk of trauma. *Pediatrics* **135**, 40–48.
- Man KKC, Coghill D, Chan EW, Lau WCY, Hollis C, Liddle E, Banaschewski T, McCarthy S, Neubert A, Sayal K, Ip P, Schuemie MJ, Sturkenboom M, Sonuga-Barke E, Buitelaar J, Carucci S, Zuddas A, Kovshoff H, Garas P, Nagy P, Inglis SK, Konrad K, Hage A, Rosenthal E and Wong ICK** (2017) Association of risk of suicide attempts with methylphenidate treatment. *JAMA Psychiatry* **74**, 1048–1055.
- Mangus RS, Bergman D, Zieger M and Coleman JJ** (2004) Burn injuries in children with attention-deficit/hyperactivity disorder. *Burns: Journal of the International Society for Burn Injuries* **30**, 148–150.
- National Health Insurance Research Database** (2013) National Health Insurance Research Database. Available at https://nhird.nhri.org.tw/en/Data_Subsets.html.
- Nelson LD, Guskiewicz KM, Marshall SW, Hammeke T, Barr W, Randolph C and McCrema MA** (2016) Multiple self-reported concussions are more prevalent in athletes with ADHD and learning disability. *Clinical Journal of Sport Medicine* **26**, 120.
- Patrick KE, Schultheis MT, Agate FT, McCurdy MD, Daly BP, Tarazi RA, Chute DL and Hurewitz F** (2020) Executive function 'drives' differences in simulated driving performance between young adults with and without autism spectrum disorder. *Child Neuropsychology: A Journal on Normal and Abnormal Development in Childhood and Adolescence* **26**, 649–665.
- Raman SR, Marshall SW, Haynes K, Gaynes BN, Naftel AJ and Sturmer T** (2013) Stimulant treatment and injury among children with attention deficit hyperactivity disorder: an application of the self-controlled case series study design. *Injury Prevention: Journal of the International Society for Child and Adolescent Injury Prevention* **19**, 164–170.
- Romo L, Julien Sweerts S, Ordonneau P, Blot E and Gicquel L** (2019) Road accidents in young adults with ADHD: which factors can explain the occurrence of injuries in drivers with ADHD and how to prevent it? *Applied Neuropsychology. Adult*, 1–6.
- Rowe R, Maughan B and Goodman R** (2004) Childhood psychiatric disorder and unintentional injury: findings from a national cohort study. *Journal of Pediatric Psychology* **29**, 119–130.
- Savage MN and Bouck EC** (2017) Predictors of risky behavior and offending for adolescents with mild intellectual disability. *Intellectual and Developmental Disabilities* **55**, 154–166.
- Shin JY, Roughead EE, Park BJ and Pratt NL** (2016) Cardiovascular safety of methylphenidate among children and young people with attention-deficit/hyperactivity disorder (ADHD): nationwide self controlled case series study. *BMJ* **353**, i2550.
- Sundelin HEK, Chang Z, Larsson H, Lichtenstein P, Almqvist C, Tomson T and Ludvigsson JF** (2018) Epilepsy, antiepileptic drugs, and serious transport accidents: a nationwide cohort study. *Neurology* **90**, e1111–e1118.

- Tai YM and Chiu HW** (2009) Comorbidity study of ADHD: applying association rule mining (ARM) to national health insurance database of Taiwan. *International Journal of Medical Informatics* **78**, e75–e83.
- Tai YM, Gau SS and Gau CS** (2013) Injury-proneness of youth with attention-deficit hyperactivity disorder: a national clinical data analysis in Taiwan. *Research in Developmental Disabilities* **34**, 1100–1108.
- Thompson AL, Molina BS, Pelham Jr W and Gnagy EM** (2007). Risky driving in adolescents and young adults with childhood ADHD. *Journal of Pediatric Psychology* **32**, 745–759.
- Toroyan T, Peden MM and Iaych K** (2013) WHO Launches second global status report on road safety. *Injury Prevention: Journal of the International Society for Child and Adolescent Injury Prevention* **19**, 150.
- Vaa T** (2014) ADHD And relative risk of accidents in road traffic: a meta-analysis. *Accident; Analysis and Prevention* **62**, 415–425.
- Whitaker HJ, Farrington CP, Spiessens B and Musonda P** (2006) Tutorial in biostatistics: the self-controlled case series method. *Statistics in Medicine* **25**, 1768–1797.
- World Health Organization** (2015) *Global Status Report on Road Safety 2015*. World Health Organization.
- Yeh JY, Hou TY, Tseng WT, Chen VC, Yang YH, Kuo TY, Weng JC, Lee CT, Chen YL and Lee MJ** (2020) Association between attention deficit hyperactivity disorder and risk of burn injury: a propensity-matched cohort study. *Neuropsychiatric Disease and Treatment* **16**, 1249–1255.