

Atomic and Molecular Gas in M17 SW

J. P. Pérez-Beaupuits¹, J. Stutzki², R. Güsten¹, V. Ossenkopf², and H. Wiesemeyer¹

¹Max-Planck-Institut für Radioastronomie,
 Auf dem Hügel 69, 53121 Bonn, Germany
 email: jp@mpifr.de

²I. Physikalisches Institut der Universität zu Köln,
 Zùlpicher Straße 77, 50937 Köln, Germany

Abstract. We probe the spatial distribution of the [C II] 158 μm fine-structure emission and its association with neutral and molecular gas in a $5'.7 \times 3'.7$ ($\sim 3.3 \times 2.1 \text{ pc}^2$) region of the M17 SW nebula. Comparison of velocity-resolved [C II] emission maps with other atomic and molecular tracers is possible for the first time with the dual band receiver GREAT on board the SOFIA airborne telescope. We detected [C II] emission in a much broader velocity range than the CO lines (Pérez-Beaupuits *et al.* 2012). Only [C II] narrow channel maps at intermediate velocities (between 10 and 24 km s^{-1}) show correlations with other molecular gas components, supporting a clumpy cloud scenario. At lower ($< 10 \text{ km s}^{-1}$) and higher ($> 24 \text{ km s}^{-1}$) velocities instead, we see more than 60% of the region mapped in [C II] that is not associated with other tracers of star-forming material, the so called “CO-dark” gas. Interaction with winds and outflows lead to substantial excitation of [C II] emitting gas, so that ablation and shock-interaction have to be taken into account to model the observed [C II] emission.

Keywords. ISM: structure — ISM: atoms — ISM: clouds — ISM: individual (M17)




Figure 1. Channel maps of the [C II] 158 μm emission (gray scale - K km s^{-1}) integrated in 1 km s^{-1} channel wide in the velocity range 3.5 km s^{-1} to 35.5 km s^{-1} . The contours show the corresponding channel maps of $^{12}\text{CO } J=2-1$ (left panel), and $\tau(\text{HI})$ (right panel) in 20% steps of the peak integrated values (bottom right in the maps). All maps but $\tau(\text{HI})$ were smoothed to a resolution of $\sim 21''$. This is an extended version of Fig.3 by Pérez-Beaupuits *et al.* (2012).

Reference

Pérez-Beaupuits, J. P., Wiesemeyer, H., Ossenkopf, V., Stutzki, J., *et al.* 2012, *A&A*, 542, L13