

# Cycle-Level Intersection Theory for Toric Varieties

Hugh Thomas

*Abstract.* This paper addresses the problem of constructing a cycle-level intersection theory for toric varieties. We show that by making one global choice, we can determine a cycle representative for the intersection of an equivariant Cartier divisor with an invariant cycle on a toric variety. For a toric variety defined by a fan in  $N$ , the choice consists of giving an inner product or a complete flag for  $M_{\mathbb{Q}} = \mathbb{Q} \otimes \text{Hom}(N, \mathbb{Z})$ , or more generally giving for each cone  $\sigma$  in the fan a linear subspace of  $M_{\mathbb{Q}}$  complementary to  $\sigma^{\perp}$ , satisfying certain compatibility conditions. We show that these intersection cycles have properties analogous to the usual intersections modulo rational equivalence. If  $X$  is simplicial (for instance, if  $X$  is non-singular), we obtain a commutative ring structure to the invariant cycles of  $X$  with rational coefficients. This ring structure determines cycles representing certain characteristic classes of the toric variety. We also discuss how to define intersection cycles that require no choices, at the expense of increasing the size of the coefficient field.

## 1 Introduction

The basic building block of intersection theory is the intersection of a Cartier divisor and a cycle, which is defined to be a cycle modulo rational equivalence. In this paper, we restrict our attention to toric varieties, and we ask whether there is a way to define the intersection of an equivariant Cartier divisor and an invariant cycle as a cycle, rather than a cycle modulo rational equivalence. This cannot be done naturally; a choice must be made. The goal of this paper is to show that a simple initial choice allows us to define an action of equivariant Cartier divisors on invariant cycles in a consistent manner. From this, a cycle-level intersection theory can be developed. For simplicial toric varieties, this gives rise to an intersection product on the level of cycles, analogous to the usual intersection product on cycles modulo rational equivalence.

### 1.1 Overview

First, we describe the properties which our action possesses. Let  $X$  be a toric variety,  $D$  and  $E$  equivariant Cartier divisors on  $X$ ,  $z$  an invariant cycle of  $X$  with coefficients in  $\mathbb{Q}$ , and  $\bar{z}$  the image of  $z$  in the Chow group of cycles modulo rational equivalence. Let  $D \cdot$  denote either our action of equivariant Cartier divisors on cycles, or the usual action of Cartier divisors on cycles modulo rational equivalence, depending on context. Then the action we define has the following properties:

---

Received by the editors January 8, 2003; revised August 2, 2003.  
AMS subject classification: Primary 14M25; Secondary 14C17.  
Keywords: toric varieties, intersection theory.  
©Canadian Mathematical Society 2004.

- (1)  $\overline{D \cdot z} = D \cdot \bar{z}$ , i.e., when we pass to rational equivalence, our intersection cycle agrees with the usual intersection defined on cycles modulo rational equivalence.
- (2) If  $D$  and  $z$  intersect properly, then there is a naturally well-defined cycle-level intersection, namely  $[D|_z]$ , the cycle corresponding to the restriction of  $D$  to  $z$ . In this case,  $D \cdot z = [D|_z]$ .
- (3)  $D \cdot (E \cdot z) = E \cdot (D \cdot z)$ .

Now, we briefly describe the choice that must be made in order to define our action. Let  $N$  be a free abelian group of rank  $n$ . Let  $\Sigma$  be a fan in  $N_{\mathbb{Q}} = N \otimes \mathbb{Q}$ , that is, a collection of strongly convex rational polyhedral cones in  $N_{\mathbb{Q}}$ , such that any face of a cone  $\sigma$  in  $\Sigma$  is again in  $\Sigma$  and the intersection of any two cones in  $\Sigma$  is a face of both and is again in  $\Sigma$ . (For further details about toric varieties, see Section 1.2 and [4].) Let  $M = \text{Hom}(N, \mathbb{Z})$ . Each  $\sigma \in \Sigma$  defines a subspace  $\sigma^{\perp}$  of  $M_{\mathbb{Q}} = \mathbb{Q} \otimes M$ . To define our action, we need a function  $\Psi$  associating to each cone  $\sigma$  of  $\Sigma$  a subspace  $\Psi(\sigma)$  with the following two properties:

- (1) For each  $\sigma \in \Sigma$ ,  $\Psi(\sigma)$  and  $\sigma^{\perp}$  are complementary subspaces of  $M_{\mathbb{Q}}$ .
- (2) If  $\sigma \in \Sigma$  and  $\tau$  is a face of  $\sigma$ , then  $\Psi(\sigma)$  contains  $\Psi(\tau)$ .

There are two easy ways to define a map  $\Psi$  having these properties, starting with either an inner product on  $M_{\mathbb{Q}}$  or a complete flag in  $M_{\mathbb{Q}}$  which is generic in a suitable sense with respect to  $\Sigma$ .

Section 2 is concerned with constructing our action and showing that it has the properties we have already described. We also prove an analogue of the projection formula from intersection theory, as follows. Consider the situation where we have two toric varieties and a proper equivariant map between them  $f: X' \rightarrow X$ . In this case, if we choose  $\Psi$  for  $X$  and  $\Psi'$  for  $X'$  in compatible ways, then for  $z$  an invariant cycle on  $X'$  and  $D$  an equivariant Cartier divisor on  $X$ , we have:

$$f_*(f^*(D) \cdot z) = D \cdot f_*(z).$$

In Section 3, we consider the case where  $\Sigma$  is simplicial. In this case we obtain a commutative ring structure on the invariant cycles of  $X$  with rational coefficients. We work out the details for one realization of projective space as a toric variety. In Section 4, we discuss techniques for performing computations using our action. In the final section, we take a quick look at two further topics. First, we show how we can produce actions of equivariant Cartier divisors on invariant cycles independent of any choice, at the expense of enlarging the coefficient field. Secondly, we consider how one can go about constructing cycles which represent characteristic classes of a toric variety, and the Todd class in particular.

### 1.2 Notation

We begin by recalling necessary background about toric varieties. For further explanation and proofs, see [4].

Let  $N$  be a free abelian group of rank  $n$ , and  $M = \text{Hom}(N, \mathbb{Z})$  its dual. We will denote  $\mathbb{Q} \otimes N$  by  $N_{\mathbb{Q}}$ . In general, a subscript  $\mathbb{Q}$  will mean “tensor by  $\mathbb{Q}$ ”, and similarly for other fields.

Let  $\Sigma$  be a fan in  $N$ , and  $X = X(\Sigma)$  the associated toric variety. (To define  $X(\Sigma)$  we have to choose a ground field, but this choice is of no importance for our purposes. In our examples, the ground field will be the complex numbers.) If  $\tau$  and  $\sigma$  are two cones of  $\Sigma$ ,  $\sigma \succ \tau$  means that  $\tau$  is a proper face of  $\sigma$ , and  $\sigma \rightarrow \tau$  means that  $\tau$  is a maximal proper face of  $\sigma$ , also known as a facet. We shall say that  $\Sigma$  is affine if it consists of a single cone and its faces. For general  $\Sigma$ , and  $\sigma \in \Sigma$ , we write  $U_\sigma$  for the toric variety associated to the affine fan consisting of  $\sigma$  and its faces.  $U_\sigma$  is canonically an open subset of  $X$ ; the collection of all the  $U_\sigma$  covers  $X$ .

For  $\sigma \in \Sigma$ ,  $N_\sigma$  is the subgroup generated by  $\sigma \cap N$ , and  $N(\sigma) = N/N_\sigma$ .  $M(\sigma)$  is the dual of  $N(\sigma)$ , so naturally a sublattice of  $M$ ; in fact,  $M(\sigma) = M \cap \sigma^\perp$ .  $M_\sigma = M/M(\sigma)$  is dual to  $N_\sigma$ . For  $\sigma \prec \tau$ , we set  $M(\sigma)_\tau = M(\sigma)/M(\tau)$ , and  $N(\sigma)_\tau = N_\tau/N_\sigma$ . These two are also dual. All the duality pairings mentioned in this paragraph are denoted  $\langle \cdot, \cdot \rangle$ .

Let  $D \in \text{Div}_T(X)$ , the equivariant Cartier divisors on  $X$  (from now on, we shall omit the word equivariant). Define  $m_\sigma \in M_\sigma$  by requiring that  $\chi^{m_\sigma}$  be a local equation for  $D$  on  $U_\sigma$ . (Frequently in the study of toric varieties it is preferred to take the local equations for  $-D$ . To follow that convention would introduce a number of negative signs, and so we have chosen not to.) By an abuse of terminology, we will refer to the collection of  $m_\sigma$  associated to a divisor  $D$  as its local equations. The  $m_\sigma$  satisfy certain agreement conditions: if  $\tau \prec \sigma$ , the image of  $m_\sigma$  under the natural map from  $M_\sigma$  to  $M_\tau$  is  $m_\tau$ . Conversely, given a collection of  $m_\sigma$  satisfying these conditions, they determine a Cartier divisor.

We shall also want to consider  $\mathbb{Q}$ -Cartier divisors, by which we mean rational multiples of Cartier divisors. More precisely, we will think of a  $\mathbb{Q}$ -Cartier divisor as a collection of  $m_\sigma \in \mathbb{Q} \otimes M_\sigma$ , satisfying the same agreement conditions as above for Cartier divisors.

Given  $\sigma \in \Sigma$  a cone of dimension  $n - k$ , there is an associated  $k$ -dimensional subvariety of  $X$ , which is denoted  $V(\sigma)$ .  $V(\sigma)$  is also a toric variety, and is given by a fan in  $N(\sigma)$ . Its fan is  $\text{Star}(\sigma) = \{(\tau + N_\sigma)/N_\sigma \mid \tau \in \Sigma, \tau \succeq \sigma\}$ .

The group of invariant  $k$ -cycles, which we will denote  $Z_k(X)$ , is the free abelian group on the set of  $[V(\sigma)]$ , where  $\sigma$  ranges over the cones of dimension  $n - k$  in  $\Sigma$ . From now on, we omit the word ‘‘invariant’’: all our cycles are understood to be torus invariant. Let  $\rho_1, \dots, \rho_r$  be the rays of  $\Sigma$ . Denote  $V(\rho_i)$  by  $D_i$ .

Denote by  $A_k(X)$  the  $k$ -th Chow group of  $X$ . It has a presentation as the quotient of  $Z_k(X)$  by the invariant  $k$ -cycles which are rationally equivalent to zero. (See [5] for more details.)

There is a map from Cartier divisors to  $Z_{n-1}(X)$ . If  $D$  is a Cartier divisor, the associated element of  $Z_{n-1}(X)$  is  $[D] = \sum_{i=1}^r \langle m_{\rho_i}, v_i \rangle D_i$ , where  $v_i$  is the first lattice point of  $N$  along  $\rho_i$  and  $m_{\rho_i}$  is the local equation for  $D$  on  $U_{\rho_i}$ . The same definition gives a map from  $\mathbb{Q}$ -Cartier divisors to  $Z_{n-1}(X)_\mathbb{Q}$  which extends the previous map.

We shall also be interested in this map from  $\mathbb{Q}$ -Cartier divisors to invariant cycles when the divisors are on  $V(\sigma)$  rather than  $X$ . Let  $D$  be a  $\mathbb{Q}$ -Cartier divisor on  $V(\sigma)$  and let  $\tau$  be a maximal proper face of  $\sigma$ . Then the coefficient of  $[V(\tau)]$  in  $[D]$  is  $\langle m_\tau, n_{\tau,\sigma} \rangle$ , where  $n_{\tau,\sigma}$  is the generator of the (1-dimensional) image of  $\tau$  in  $N(\sigma)$ , *i.e.*, the first lattice point of  $N(\sigma)$  along the ray in  $N(\sigma)$  corresponding to  $\tau$ .

### 1.3 Preliminary Linear Algebra

Let  $<$  denote the relation of being a vector subspace. We establish two lemmas.

**Lemma 1.1** *Let  $V \geq A \geq B$ . Let  $A'$  and  $B'$  be complementary subspaces to  $A$  and  $B$  in  $V$  such that  $B' \geq A'$ . Then*

- (i) *There is a canonical isomorphism between  $A/B$  and  $B'/A'$ .*
- (ii) *There is a canonical map from  $V/B$  to  $A/B$ .*

**Proof** This follows from the facts that  $V$  is canonically isomorphic to  $B \oplus (A \cap B') \oplus A'$ , while  $A$  is canonically isomorphic to  $B \oplus (A \cap B')$ . ■

**Lemma 1.2** *Let  $V \geq A \geq B \geq C$ . Let  $A'$ ,  $B'$ , and  $C'$  be complementary subspaces to  $A$ ,  $B$ , and  $C$ , such that  $V \geq C' \geq B' \geq A'$ . Then the following diagram commutes, where the horizontal arrows are projection, and the vertical arrows are the maps from Lemma 1.1.*

$$\begin{array}{ccc}
 V/C & \longrightarrow & V/B \\
 \downarrow & & \downarrow \\
 A/C & \longrightarrow & A/B
 \end{array}$$

**Proof** The proof is similar to that of the previous lemma.  $V$  is canonically isomorphic to  $C \oplus (B \cap C') \oplus (A \cap B') \oplus A'$ , and similarly for  $A$  and  $B$ . ■

## 2 The Action and Its Properties

**Convention** We shall frequently need to refer to sub- and quotient groups of  $M$  tensored by  $\mathbb{Q}$ , and very seldom to the groups themselves. Therefore, we shall drop the  $\mathbb{Q} \otimes$  from our notation: whenever we refer to  $M, M(\sigma), M_\tau, M(\sigma)_\tau$ , etc., the  $\mathbb{Q} \otimes$  is to be understood. Note that this does not apply to sub- and quotient groups of  $N$ .

### 2.1 Definition of the Action

To define the action, we fix a map  $\Psi$  taking cones of  $\Sigma$  to linear subspaces of  $M$  of the same dimension, satisfying the following properties

- (i)  $\Psi(\sigma)$  and  $M(\sigma)$  span  $M$  (or equivalently,  $\Psi(\sigma) \cap (M(\sigma)) = \{0\}$ ).
- (ii) If  $\sigma \succ \tau$ ,  $\Psi(\sigma) > \Psi(\tau)$ .

If  $\Psi$  satisfies these two properties, we call  $\Psi$  a choice of complements for  $\Sigma$ .

We immediately give possible constructions of  $\Psi$ .

**Example 1 (Inner Product)** Fix an inner product on  $M$ . Define  $\Psi(\sigma)$  to be the orthogonal complement of  $\sigma^\perp$  in  $M$ .

**Example 2 (Generic Flag)** Fix a complete flag in  $M$ ,  $0 = F_0 < F_1 < \dots < F_n = M$ , which is generic with respect to  $\Sigma$  in the sense that if  $\sigma$  is  $k$ -dimensional, then  $\sigma^\perp \cap F_k = \{0\}$ . Define  $\Psi(\sigma) = F_k$  for all  $\sigma$  of dimension  $k$ .

Let  $D$  be a  $\mathbb{Q}$ -Cartier divisor and  $\sigma$  a cone of dimension  $n - k$  in  $\Sigma$ . We proceed to define the action of  $D$  on  $[V(\sigma)]$ .

The decomposition  $M = M(\sigma) \oplus \Psi(\sigma)$  determines a projection  $\pi_\sigma : M \rightarrow M(\sigma)$ . Also, for any  $\tau \succ \sigma$ , writing  $M_\tau = M/M(\tau)$ ,  $M(\sigma)_\tau = M(\sigma)/M(\tau)$ , we can apply Lemma 1.1 to get a map, denoted the same way,  $\pi_\sigma : M_\tau \rightarrow M(\sigma)_\tau$ . (Observe that in the case of the inner product action, the map  $\pi_\sigma$  is orthogonal projection from  $M$  to  $M(\sigma)$ .)

For  $\tau \in \Sigma$ , let the local equation of  $D$  on  $\tau$  be  $m_\tau$ . For  $\tau \succ \sigma$ , define  $\overline{m}_\tau \in M(\sigma)_\tau$  by  $\overline{m}_\tau = \pi_\sigma(m_\tau)$ . We verify that these  $\overline{m}_\tau$  form local equations for a  $\mathbb{Q}$ -Cartier divisor on the fan of  $V(\sigma)$  in  $M(\sigma)$ . Suppose  $\tau \succ \gamma \succ \sigma$ . We need to show that  $\overline{m}_\tau$  maps to  $\overline{m}_\gamma$  under the natural map from  $M(\sigma)_\tau$  to  $M(\sigma)_\gamma$ . But this follows immediately from Lemma 1.2 and the fact that  $m_\tau$  goes to  $m_\gamma$  under the map from  $M_\tau$  to  $M_\gamma$ . Thus, the  $\overline{m}_\tau$  form local equations for a  $\mathbb{Q}$ -Cartier divisor, which we denote  $D_\sigma$ .

So, define  $D \cdot [V(\sigma)] = [D_\sigma]$ . Explicitly, this definition is:

$$D \cdot [V(\sigma)] = \sum_{\tau \rightarrow \sigma} \langle \pi_\sigma(m_\tau), n_{\tau,\sigma} \rangle [V(\tau)].$$

For any  $\mathbb{Q}$ -Cartier divisor and any  $k$ , this gives us a map  $Z_k(X)_\mathbb{Q} \rightarrow Z_{k-1}(X)_\mathbb{Q}$ , as desired.

**Example 3 ( $\mathbb{C}^n$  with the standard inner product)** Let  $e_1, \dots, e_n$  be a basis of  $N$ , and  $e_1^*, \dots, e_n^*$  the dual basis of  $M$ . Put the usual inner product on  $M$  so that the  $e_i^*$  form an orthonormal basis. Let us consider the example where  $\Sigma$  is the fan consisting of the positive orthant  $\sigma$  and its faces, so  $X(\Sigma) \cong \mathbb{C}^n$ . Let  $\rho_i$  be the ray in the positive  $e_i$  direction, and  $D_i = V(\rho_i)$ . Then  $D_i \cdot V(\tau) = 0$  if  $\rho_i \prec \tau$ , and  $D_i \cdot V(\tau) = V(\tau + \rho_i)$  otherwise.

### 2.2 Elementary properties

Given  $D$  a  $\mathbb{Q}$ -Cartier divisor, we say the  $D$  intersects  $V(\sigma)$  properly if when  $D$  is written as a sum of codimension one irreducible subvarieties, no subvarieties containing  $[V(\sigma)]$  occur with non-zero coefficients. Equivalently,  $D$  intersects  $[V(\sigma)]$  properly if it restricts to a  $\mathbb{Q}$ -Cartier divisor on  $V(\sigma)$ . If  $m_\sigma$  is the local equation of  $D$  on  $\sigma$ ,  $D$  intersects  $V(\sigma)$  properly iff  $m_\sigma = 0$ .

We now examine  $D \cdot V(\sigma)$  when  $D$  intersects  $V(\sigma)$  properly.

**Proposition 2.1** *Let  $D$  be a  $\mathbb{Q}$ -Cartier divisor, and suppose it intersects  $[V(\sigma)]$  properly. Then  $D_\sigma$  is the restriction of  $D$  to  $V(\sigma)$ , so  $D \cdot [V(\sigma)]$  is the cycle corresponding to the restriction of  $D$  to  $V(\sigma)$ .*

**Proof** Since  $m_\sigma = 0$ , the image of  $m_\tau$  in  $M_\sigma$  is 0, so  $m_\tau \in M(\sigma)_\tau$ , and thus  $\pi_\sigma(m_\tau) = m_\tau$ . Thus, the local equations for  $D_\sigma$  are just the local equations for  $D$ . Thus we see that  $D_\sigma$  is the restriction of  $D$  to  $V(\sigma)$ , and from the definition of the action it follows that  $D \cdot [V(\sigma)]$  is just the cycle associated to the restriction of  $D$  to  $V(\sigma)$ . ■

If  $X$  is a general algebraic variety, and  $D$  a Cartier divisor on  $X$ ,  $D$  induces a map from  $A_k(X)$  to  $A_{k-1}(X)$ , also denoted  $D \cdot$ . (See [3] for details.) If  $z \in Z_k(X)_\mathbb{Q}$ , denote by  $\bar{z}$  its image in  $A_k(X)_\mathbb{Q}$ . We now show that our action agrees with the usual action of Cartier divisors on Chow groups once we pass to rational equivalence.

**Proposition 2.2** *If  $z \in Z_k(X)_\mathbb{Q}$ , the image of  $D \cdot z$  in  $A_{k-1}(X)_\mathbb{Q}$  equals  $D \cdot \bar{z}$ .*

**Proof**  $D \cdot \overline{[V(\sigma)]}$  is defined by translating  $D$  by a principal divisor so that it intersects  $V(\sigma)$  properly, then taking the class modulo rational equivalence of the intersection. Observe that  $D_\sigma$  is the restriction to  $V(\sigma)$  of a translate of  $D$  by a principal divisor, so  $\overline{[D_\sigma]} = D \cdot \overline{[V(\sigma)]}$ , as desired. ■

### 2.3 All Computations Can Be Done on Affine Varieties

Recall that  $X$  is covered by affine open sets  $U_\sigma$  with  $\sigma \in \Sigma$ , where  $U_\sigma$  is the toric variety corresponding to the fan consisting of  $\sigma$  and all its faces. This covering is useful because it allows us in some cases to localize the questions we are interested in to  $U_\sigma$  and deal with them there, where we don't need to worry about the global structure of the fan.

We recall some useful facts about the covering of  $X$  by the  $U_\sigma$ . If  $\tau \prec \sigma$ , then we can perform the construction of  $V(\tau)$  in the toric variety  $U_\sigma$ . Let us denote this subvariety of  $U_\sigma$  by  $V(\tau)_\sigma$ . Then  $V(\tau)_\sigma = V(\tau) \cap U_\sigma$  is a dense open subset of  $V(\tau)$ . (See [4] for more details.)

**Lemma 2.1** *Let  $\tau \rightarrow \sigma$ . Let  $D$  be a  $\mathbb{Q}$ -Cartier divisor on  $X$ . Then the coefficient of  $[V(\tau)]$  in  $D \cdot [V(\sigma)]$  equals the coefficient of  $[V(\tau)_\tau]$  in  $D|_{U_\tau} \cdot [V(\sigma)_\tau]$ .*

**Proof** Let  $m_\tau$  be the local equation of  $D$  on  $\tau$ . The local equation of the restriction of  $D$  to  $U_\tau$  is also  $m_\tau$ . Thus, the coefficients in question both equal  $\langle \pi_\sigma(m_\tau), n_{\tau,\sigma} \rangle$ . ■

The analogous result for the effect of a composition of the action of multiple Cartier divisors is also true. For simplicity, if  $D$  and  $E$  are  $\mathbb{Q}$ -Cartier divisors and  $z$  is a cycle, instead of writing  $D \cdot (E \cdot z)$ , we write  $D \cdot E \cdot z$ .

**Corollary** *Let  $\tau \succ \sigma$ . Let  $E_1, \dots, E_s$  be  $\mathbb{Q}$ -Cartier divisors on  $X$ . Then the coefficient of  $[V(\tau)]$  in  $E_1 \cdots E_s \cdot [V(\sigma)]$  equals the coefficient of  $[V(\tau)_\tau]$  in  $E_1|_{U_\tau} \cdots E_s|_{U_\tau} \cdot [V(\sigma)_\tau]$ .*

**Proof** This follows by a repeated application of Lemma 2.1. ■

In fact, computing the coefficient of a cycle  $[V(\tau)]$  as above can be reduced to computing the coefficient of a cycle  $[V(\tau')]$  where  $\tau'$  is of maximal dimension in a lattice  $N'$ .

Let  $\tau$  be a cone of  $\Sigma$ . Let  $N' = N_\tau$ , and let  $\Sigma'$  be the fan in  $N'$  consisting of  $\tau$  and its faces. For  $\gamma \preceq \tau$ , let  $\gamma'$  denote the corresponding cone of  $\Sigma'$ . Let  $E_1, \dots, E_s$  be  $\mathbb{Q}$ -Cartier divisors on  $X$ , and for  $1 \leq i \leq s$  let  $E'_i$  be the  $\mathbb{Q}$ -Cartier divisor on  $X(\Sigma')$  having the same local equations on  $\gamma'$  as  $E_i$  does on  $\gamma$ , for all  $\gamma \preceq \tau$ . Then we have the following result:

**Lemma 2.2** *The coefficient of  $[V(\tau)]$  in  $E_1 \cdots E_s \cdot [V(\sigma)]$  equals the coefficient of  $[V(\tau')]$  in  $E'_1 \cdots E'_s \cdot [V(\sigma')]$ .*

**Proof** As in the proof of Lemma 2.1, the explicit expression for the two coefficients is the same. ■

### 2.4 Commutativity

We now prove the following commutativity result:

**Theorem 2.1** *Let  $D$  and  $E$  be  $\mathbb{Q}$ -Cartier divisors on  $X$ , and  $\sigma$  a cone of  $\Sigma$  of dimension  $n - k$ . Then  $D \cdot E \cdot [V(\sigma)] = E \cdot D \cdot [V(\sigma)]$ .*

**Proof**  $D \cdot E \cdot [V(\sigma)]$  is a linear combination of  $[V(\tau)]$  where  $\tau$  ranges over cones of dimension  $n - k + 2$ . More explicitly, if the local equations for  $E$  are  $e_\delta$ ,

$$E \cdot V(\sigma) = \sum_{\delta \rightarrow \sigma} \langle \pi_\sigma(e_\delta), n_{\delta,\sigma} \rangle [V(\delta)]$$

and, if the local equations for  $D$  are  $d_\tau$ ,

$$D \cdot E \cdot [V(\sigma)] = \sum_{\tau \rightarrow \delta} \sum_{\delta \rightarrow \sigma} \langle \pi_\delta(d_\tau), n_{\tau,\delta} \rangle \langle \pi_\sigma(e_\delta), n_{\delta,\sigma} \rangle [V(\tau)].$$

Suppose we wish to compute the coefficient of some particular  $[V(\tau)]$  in the above expression. From convex geometry, we know that there are exactly two cones  $\delta$  satisfying  $\tau \rightarrow \delta \rightarrow \sigma$ . Call these cones  $\gamma$  and  $\beta$ . Then the coefficient of  $[V(\tau)]$  in  $D \cdot E \cdot [V(\sigma)]$  is

$$(1) \quad \langle \pi_\gamma(d_\tau), n_{\tau,\gamma} \rangle \langle \pi_\sigma(e_\gamma), n_{\gamma,\sigma} \rangle + \langle \pi_\beta(d_\tau), n_{\tau,\beta} \rangle \langle \pi_\sigma(e_\beta), n_{\beta,\sigma} \rangle.$$

To prove that  $D \cdot E \cdot [V(\sigma)] = E \cdot D \cdot [V(\sigma)]$ , we need to check that the coefficient of  $[V(\tau)]$  in the two expressions is the same for all  $\tau$ . Thus, to show commutativity, it suffices to show that the expression (1) is the same after we change all the  $d$ 's to  $e$ 's and vice versa.

Our first goal is to modify (1) so that all the pairings in it are between the same pair of spaces,  $M(\sigma)_\tau$  and  $\mathbb{Q} \otimes N(\sigma)_\tau$ . By replacing  $\pi_\sigma(e_\gamma)$  and  $\pi_\sigma(e_\beta)$  by  $\pi_\sigma(e_\tau)$ ,




Figure 1: An illustration of  $N(\sigma)_\tau$  and  $M(\sigma)_\tau$

which we may do by the compatibility conditions for local equations, we have put the second and fourth pairings of (1) into the desired form.

Figure 1 shows an example where  $\Psi$  is induced from the standard inner product.

We move on to consider the first and third pairings. Both  $n_{\tau,\gamma}$  and  $n_{\tau,\beta}$  live in quotients of  $N(\sigma)_\tau$ , so we choose liftings of them to  $\mathbb{Q} \otimes N(\sigma)_\tau$ : define  $\tilde{n}_{\tau,\gamma} \in \mathbb{Q} \otimes N(\sigma)_\tau$  to be the lifting of  $n_{\tau,\gamma}$  which lies in  $\beta$ . Similarly, define  $\tilde{n}_{\tau,\beta}$  to be the lifting of  $n_{\tau,\beta}$  which lies in  $\gamma$ . (See Figure 2.) Thus we can write the coefficient of  $[V(\tau)]$  in  $D \cdot E \cdot [V(\sigma)]$  as

$$\langle \pi_\sigma(e_\tau), n_{\gamma,\sigma} \rangle \langle \pi_\gamma(d_\tau), \tilde{n}_{\tau,\gamma} \rangle + \langle \pi_\sigma(e_\tau), n_{\beta,\sigma} \rangle \langle \pi_\beta(d_\tau), \tilde{n}_{\tau,\beta} \rangle$$

where all the pairings are between  $M(\sigma)_\tau$  and  $\mathbb{Q} \otimes N(\sigma)_\tau$ .

We would now like to analyze  $\tilde{n}_{\tau,\gamma}$  and  $\tilde{n}_{\tau,\beta}$  further. It is clear by construction that there are constants  $c$  and  $c'$  such that  $c\tilde{n}_{\tau,\gamma} = n_{\beta,\sigma}$  and  $c'\tilde{n}_{\tau,\beta} = n_{\gamma,\sigma}$ . We need to show that  $c = c'$ . We do this by the following lemma:

**Lemma 2.3** *Let  $L$  be the sublattice of  $N(\sigma)_\tau$  generated by  $n_{\gamma,\sigma}$  and  $n_{\beta,\sigma}$ . The following equalities hold:*

$$\begin{aligned} [N(\sigma)_\tau : L] \tilde{n}_{\tau,\gamma} &= n_{\beta,\sigma} \\ [N(\sigma)_\tau : L] \tilde{n}_{\tau,\beta} &= n_{\gamma,\sigma}. \end{aligned}$$

**Proof** We prove the first equality.

Let  $W$  denote the fundamental domain of  $L$  whose vertices are  $0$ ,  $n_{\gamma,\sigma}$ ,  $n_{\beta,\sigma}$ , and  $n_{\gamma,\sigma} + n_{\beta,\sigma}$ , and we consider  $W$  not to include any points from its left or top sides; in particular the unique point of  $L$  contained in  $W$  is  $n_{\beta,\sigma}$ . (See Figure 2.) The number






Figure 2: An illustration of  $N(\sigma)_\tau$  showing  $W$

of points of  $N(\sigma)_\tau$  in  $W$  is then  $[N(\sigma)_\tau : L]$ . I will proceed to show that the number of points is also  $c$ .

Consider the projection  $\xi: N(\sigma)_\tau \rightarrow N(\gamma)_\tau = N(\sigma)_\tau / \mathbb{Z} \cdot n_{\gamma,\sigma} = \mathbb{Z} \cdot n_{\tau,\gamma}$ . Since  $N(\gamma)_\tau$  is a one-dimensional lattice, we can put a linear order on it by saying that  $n_{\tau,\gamma}$  designates the positive direction. Now  $\xi$  defines a bijection between  $W \cap N(\sigma)_\tau$  and  $\{x \in N(\gamma)_\tau \mid 0 < x \leq \xi(n_{\beta,\sigma})\}$ . The size of the first set is  $[N(\sigma)_\tau : L]$ , while the size of the second set is  $\xi(n_{\beta,\sigma}) / n_{\tau,\gamma} = c$ . This completes the proof of the lemma. ■

Thus the coefficient of  $[V(\tau)]$  in  $D \cdot E \cdot [V(\sigma)]$  is

$$\frac{1}{[N(\sigma)_\tau : L]} (\langle \pi_\sigma(e_\tau), n_{\gamma,\sigma} \rangle \langle \pi_\gamma(d_\tau), n_{\beta,\sigma} \rangle + \langle \pi_\sigma(e_\tau), n_{\beta,\sigma} \rangle \langle \pi_\beta(d_\tau), n_{\gamma,\sigma} \rangle).$$

Coordinatize  $M(\sigma)_\tau$  by  $\langle \cdot, n_{\gamma,\sigma} \rangle, \langle \cdot, n_{\beta,\sigma} \rangle$ . Let  $\pi_\sigma(d_\tau) = (d_1, d_2)$ ,  $\pi_\sigma(e_\tau) = (e_1, e_2)$ . There are constants  $a$  and  $b$  such that  $\pi_\gamma((x, y)) = (0, ax+y)$ , while  $\pi_\beta((x, y)) = (x + by, 0)$ . Then the expression above simplifies to:

$$\frac{1}{[N(\sigma)_\tau : L]} (ad_1e_1 + d_1e_2 + d_2e_1 + bd_2e_2)$$

which is unchanged if  $d$ 's and  $e$ 's are interchanged, as desired. ■

### 2.5 Proper Maps and Push Forwards

Let  $N'$  be a free abelian group of rank  $n'$ , and  $N$  a free abelian group of rank  $n$ . Let  $\Sigma'$  and  $\Sigma$  be fans in  $N'$  and  $N$  respectively. The set of fans in lattices is made into a category by defining the morphisms from  $(N', \Sigma')$  to  $(N, \Sigma)$  to be the group homomorphisms from  $N'$  to  $N$  such that the image of every cone in  $\Sigma'$  is contained in some cone in  $\Sigma$ . Such a map of fans induces an equivariant morphism of toric varieties. From now on, whenever we refer to a map between toric varieties, we mean one which is induced from a map of fans.

A proper map of toric varieties has a characterization in terms of the map of fans: given a map  $\phi$  from  $(N', \Sigma')$  to  $(N, \Sigma)$ , the corresponding map of toric varieties is proper iff  $\phi^{-1}(|\Sigma|) = |\Sigma'|$ . If  $\phi$  induces a proper map of toric varieties, we say that  $\phi$  is a proper map.

In general, given a proper map  $f: X' \rightarrow X$  of algebraic varieties, there is a push forward map  $f_*$  on cycles. If  $Z$  is a  $k$ -dimensional subvariety of  $X'$ , let  $W$  be the closure of the image of  $Z$ . If the dimension of  $W$  is less than that of  $Z$ ,  $f_*([Z]) = 0$ . If the dimensions are the same,  $f_*([Z]) = [k(Z):k(W)][W]$ , where  $k(\cdot)$  denotes the function field. (See [3] for details.)

In the case of toric varieties and invariant subvarieties, this has a translation to lattices. Again, suppose  $f$  is a proper map from  $X'$  to  $X$ , toric varieties given by fans  $\Sigma'$  in  $N'$  and  $\Sigma$  in  $N$ , and  $\phi$  is the corresponding map from  $(N', \Sigma')$  to  $(N, \Sigma)$ . Suppose for convenience that  $\Sigma$  is full-dimensional in  $N$ , that is, that  $\Sigma$  is not contained in any proper subspace of  $N$ . Let  $\sigma'$  be a cone in  $\Sigma'$ . By the definition of a map,  $\phi(\sigma')$  is contained in some cone of  $\Sigma$ . Taking the intersection of all cones of  $\Sigma$  containing  $\phi(\sigma')$ , we find a minimal cone  $\sigma \in \Sigma$  containing  $\phi(\sigma')$ . Then if the codimension of  $\sigma$  is less than the codimension of  $\sigma'$ ,  $f_*([V(\sigma')]) = 0$ . If the codimensions are equal,  $f_*([V(\sigma')]) = [M'(\sigma'):\phi^*(M(\sigma))][V(\sigma)]$ , where  $\phi^*$  denotes the induced map from  $M$  to  $M'$ , and  $[\cdot : \cdot]$  denotes the index of the second lattice in the first. (Here we mean the actual lattices, without tensoring by  $\mathbb{Q}$ .)

In the general context of a proper map  $f: X' \rightarrow X$  of algebraic varieties, there are two important facts about  $f_*$ . Firstly, it respects rational equivalence, so it passes to Chow groups, and secondly, given  $\bar{z}$  an element of a Chow group of  $X'$  and a Cartier divisor  $D$  on  $X$ ,  $f_*(f^*(D) \cdot \bar{z}) = D \cdot f_*(\bar{z})$ .

We proceed to prove an analogous identity for our action of  $\mathbb{Q}$ -Cartier divisors on invariant cycles.

**Theorem 2.2** *Let  $f: X' \rightarrow X$  be a proper map of toric varieties, with corresponding fans  $\Sigma'$  in  $N'$  and  $\Sigma$  in  $N$  and corresponding map  $\phi: N' \rightarrow N$ . Let  $\Psi'$  and  $\Psi$  be choices of complements for  $\Sigma'$  and  $\Sigma$ , such that if  $\phi(\sigma') \subset \sigma$  with  $\sigma'$  and  $\sigma$  of the same codimension, then  $\phi^*(\Psi(\sigma)) \subset \Psi(\sigma')$ . Let  $D$  be a  $\mathbb{Q}$ -Cartier divisor on  $X$ . Let  $\sigma' \in \Sigma'$ . Then*

$$f_*(f^*(D) \cdot [V(\sigma')]) = D \cdot f_*([V(\sigma')]).$$

**Proof** For  $\gamma' \in \Sigma'$  let  $c(\gamma')$  denote the smallest cone in  $\Sigma$  containing  $\phi(\gamma')$ . Let  $P$  denote the set of  $\gamma' \in \Sigma'$  such that the codimension of  $\gamma'$  equals the codimension of  $c(\gamma')$ . (Note that  $\gamma' \in P$  iff  $f_*([V(\gamma')]) \neq 0$ .)

Let the dimension of  $\sigma'$  be  $n' - k$ . The following lemma describes the possible arrangements of  $\sigma'$ , the cones over  $\sigma'$ , and smallest cones in  $\Sigma$  containing them.

**Lemma 2.4** *Let  $\Sigma$  in  $N$  and  $\Sigma'$  in  $N'$  be two fans, and let  $\phi$  be a proper map from  $N'$  to  $N$ . Let  $\sigma'$  be a codimension- $k$  cone of  $\Sigma'$ . Then one of the three following situations must occur:*

- (i)  $\sigma' \notin P$ , and for all  $\tau' \rightarrow \sigma'$ ,  $\tau' \notin P$ .
- (ii)  $\sigma' \notin P$ , there are two cones  $\tau', \gamma' \in P$  such that  $\tau' \rightarrow \sigma'$ ,  $\gamma' \rightarrow \sigma'$ ,  $c(\sigma') = c(\tau') = c(\gamma')$ , and for any other  $\delta' \rightarrow \sigma'$ ,  $\delta' \notin P$ .
- (iii)  $\sigma' \in P$ . For each  $\tau \rightarrow c(\sigma')$ , there is exactly one  $\tau' \rightarrow \sigma'$  such that  $c(\tau') = \tau$ . Any other  $\gamma' \rightarrow \sigma'$  (i.e., one which doesn't correspond to some  $\tau \rightarrow c(\sigma')$ ) is not contained in  $P$ .

The three situations are shown in Figure 3. In order to represent a three-dimensional situation, the diagrams are of a 2-dimensional affine slice of the fans in question. The map  $\phi$  is the identity in each case.

**Proof** First, let us suppose  $\sigma' \notin P$ . Then if we are not in (i), there is some  $\tau' \rightarrow \sigma'$ , such that  $\tau' \in P$ . Since  $\sigma' \notin P$  but  $\tau' \in P$ ,  $c(\sigma') = c(\tau')$ . Let us denote  $c(\tau')$  by  $\tau$ .

The image of  $\sigma'$  is not contained in the boundary of  $\tau$ , since  $c(\sigma') = \tau$ . Thus, we can pick a point  $x$  in the interior of  $\sigma'$  such that  $\phi(x)$  is in the interior of  $\tau$ .

Now consider a ray in  $N_{\mathbb{Q}}$  from a point in the interior of  $\tau'$ , passing through  $x$ , and continuing beyond. Let  $y$  be a point just past  $x$  on this ray.  $\phi(y) \in \tau$ , so by properness,  $y$  is in some cone of  $\Sigma'$ .

Now we need a lemma:

**Lemma 2.5** *Let  $\phi$  be a proper map from  $\Sigma'$  to  $\Sigma$ . Let  $\tau$  be a cone of  $\Sigma$ , of codimension  $s$ . Then the inverse image of  $\tau$  is a union of cones of  $\Sigma'$  all of codimension  $s$ .*

**Proof** Suppose some  $y$  in the inverse image of  $\tau$  is not contained in any codimension- $s$  cone of  $\Sigma'$ . By properness, it is contained in some cone of  $\Sigma'$ . Let  $\gamma'$  be the smallest cone of  $\Sigma'$  containing  $y$ , and let the codimension of  $\gamma'$  be  $t < s$ .


Pick a small line segment  $\ell'$  contained in  $\gamma'$  with midpoint  $y$ , such that the only point of  $\ell'$  contained in  $\phi^{-1}(\tau)$  is  $y$ . Then  $\phi(\ell')$  is also a line segment, which we denote  $\ell$ . The endpoints of  $\ell$  are in  $c(\gamma')$  but not in  $\tau$ , and  $\phi(y) \in \tau$  is a convex combination of the endpoints. This contradicts the assumption that  $\Sigma$  is a fan. ■

From Lemma 2.5 it follows that  $y$  must be contained in some cone  $\gamma'$  of codimension  $k - 1$ . By construction,  $\gamma'$  has  $\sigma$  in its boundary. Also by Lemma 2.5,  $c(\gamma') = \tau$ , and so  $\gamma' \in P$ .


Now suppose there was some other cone  $\delta' \rightarrow \sigma'$ , such that  $\delta' \in P$ . Now  $c(\delta') \supset c(\sigma') = \tau$ , so  $\delta' \in P$  implies that  $c(\delta') = \tau$ .

Thus  $\tau'$ ,  $\gamma'$ , and the putative  $\delta'$  are all contained in the affine span of  $\phi^{-1}(\tau)$ , which is a dimension  $n' - k + 1$  linear subspace of  $N'$ . Now  $\sigma'$  is a cone of codimension one in that subspace, so it can be contained in at most two larger cones in the

Case (i)



Case (ii)



Case (iii)




Figure 3: The three cases of Lemma 2.4. The cones  $\sigma'$  and  $\sigma$  are shown in grey.

subspace, which means that no cone  $\delta'$  can exist. Thus, we have established all the conditions of (ii).

Now suppose  $\sigma' \in P$ . Let the cones of dimension one higher containing  $c(\sigma')$  be  $\tau_1, \dots, \tau_k$ . I claim that for each such  $\tau_i$  there is a corresponding cone  $\tau_i'$  of dimension  $n' - k + 1$  containing  $\sigma'$ , such that the image of  $\tau_i'$  is contained in  $\tau_i$ .

By Lemma 2.5, the inverse image of  $\tau_i$  is a union of cones of codimension  $k - 1$ . Pick a point in the interior of  $\sigma'$ . It is in the inverse image of  $\tau_i$ , so there is some cone  $\tau_i'$  of dimension  $n' - k + 1$  which contains it, and therefore all of  $\sigma'$ .

Since  $\sigma'$  is a full-dimensional region in the boundary of the inverse image of  $\tau_i$ , and it fully lies in some  $\tau_i'$ , it cannot lie in any other  $(n' - k + 1)$ -dimensional cone in the inverse image of  $\tau_i$ .

Thus, the situation is that  $\sigma'$  is contained in cones  $\tau_i'$  as above and also possibly certain other cones, whose images are contained in no  $\tau_i$ , and which are therefore not contained in  $P$ . Thus we have established the conditions for (iii). ■

Now, we use Lemma 2.4 to prove the theorem.

In case (i),  $f^*(D) \cdot [V(\sigma')]$  is a linear combination of  $[V(\tau)]$  with  $\tau \rightarrow \sigma$ , but since all such  $\tau$  are not in  $P$ ,  $f_*(f^*(D) \cdot [V(\sigma')]) = 0$ , and since  $\sigma \notin P$ ,  $f_*([V(\sigma)]) = 0$ , so  $D \cdot f_*([V(\sigma)]) = 0$ . So both sides of the desired identity are zero.

Now we consider case (ii). Again,  $D \cdot f_*([V(\sigma)]) = 0$ . If the local equation for  $D$  on  $\tau$  is  $m$ , the local equations for  $f^*(D)$  on  $\tau'$  and  $\gamma'$  are both  $\phi^*(m)$ . So the coefficients of  $[V(\tau')]$  and  $[V(\gamma')]$  are  $\langle \pi_{\sigma'}(\phi^*(m)), n_{\tau',\sigma'} \rangle$  and  $\langle \pi_{\sigma'}(\phi^*(m)), n_{\gamma',\sigma'} \rangle$  respectively. Observe that  $N_{\tau'} = N_{\gamma'}$ . Now  $n_{\tau',\sigma'}$  is a generator of  $N_{\tau'}/N_{\sigma'}$ , while  $n_{\gamma',\sigma'}$  is a generator of  $N_{\gamma'}/N_{\sigma'}$ , but  $n_{\tau',\sigma'}$  and  $n_{\gamma',\sigma'}$  are oriented in opposite directions. So  $n_{\tau',\sigma'} = -n_{\gamma',\sigma'}$ , and therefore  $f_*(f^*(D) \cdot [V(\sigma')]) = 0$ .

Now we consider case (iii). We wish to show that the coefficients of  $[V(\tau_i)]$  in  $f_*(f^*(D) \cdot [V(\sigma)])$  and  $D \cdot f_*([V(\sigma')])$  are the same for all  $i$ . For simplicity of notation, fix an  $i$ , and let  $\tau = \tau_i, \tau' = \tau'_i$ .

Let  $m$  be the local equation for  $D$  on  $\tau$ . Then  $\phi^*(m)$  is the local equation for  $f^*(D)$  on  $\tau'$ . The coefficient of  $[V(\tau')]$  in  $f^*(D) \cdot [V(\sigma)]$  is  $\langle \pi_{\sigma'}(\phi^*(m)), n_{\tau',\sigma'} \rangle$ . We can write (in a unique way)  $m = \pi_\sigma(m) + u$  for some  $u \in \Psi(\sigma)$ . Applying  $\phi^*$  to this, we get  $\phi^*(m) = \phi^*(\pi_\sigma(m)) + \phi^*(u)$ . Now  $u \in \Psi(\sigma)$ , so  $\phi^*(u) \in \Psi'(\sigma')$  by the assumed agreement condition between  $\Psi$  and  $\Psi'$ . And  $\phi^*(\pi_\sigma(m)) \in M'(\sigma')$ . So  $\pi_{\sigma'}(\phi^*(m)) = \phi^*(\pi_\sigma(m))$ . So the coefficient of  $[V(\tau')]$  in  $f^*(D) \cdot [V(\sigma)]$  is  $\langle \phi^*(\pi_\sigma(m)), n_{\tau',\sigma'} \rangle = \langle \pi_\sigma(m), \phi(n_{\tau',\sigma'}) \rangle$ . When we push this forward, we find the coefficient of  $[V(\tau)]$  in  $f_*(f^*(D) \cdot [V(\sigma)])$  to be

$$[M(\tau') : \phi^*(M(\tau))] \langle \pi_\sigma(m), \phi(n_{\tau',\sigma'}) \rangle.$$

On the other hand, the coefficient of  $[V(\tau)]$  in  $D \cdot f_*([V(\sigma')])$  is

$$[M(\sigma') : \phi^*(M(\sigma))] \langle \pi_\sigma(m), n_{\tau,\sigma} \rangle.$$

We wish to show that these two expressions are equal. Observe that  $\phi$  induces a map from  $N'(\sigma')_{\tau'}$  to  $N(\sigma)_\tau$ , which implies that  $\phi(n_{\tau',\sigma'})$  is in  $N(\sigma)_\tau$ , and thus an integer multiple of  $n_{\tau,\sigma}$ . Further, it is a positive integer multiple, because they point in the same direction.

The lattice  $M(\sigma')$  is generated by  $M(\tau')$  and one other element, say  $u$ . The lattice  $\phi^*(M(\sigma))$  is generated by  $\phi^*(M(\tau))$  and one other element, say  $tu$  with  $t > 0$ .

$$\langle u, n_{\tau',\sigma'} \rangle = 1, \text{ so } \langle tu, n_{\tau',\sigma'} \rangle = t. \text{ But } \langle \phi(tu), n_{\tau,\sigma} \rangle = 1. \text{ So } \phi(n_{\tau',\sigma'}) = tn_{\tau,\sigma}.$$

But it is also clear that  $t = [M(\sigma') : \phi^*(M(\sigma))] / [M(\tau') : \phi^*(M(\tau))]$ . This establishes the identity in the third case, which completes the proof of the theorem. ■

### 3 Ring Structure on $Z_*(X)_\mathbb{Q}$ for Simplicial $\Sigma$

In this section, we consider the case where  $\Sigma$  is simplicial, *i.e.*, for all  $k$ , the  $k$ -dimensional cones of  $\Sigma$  have  $k$  extremal rays. Given this condition, the map from  $\mathbb{Q}$ -Cartier divisors to  $Z_{n-1}(X)_\mathbb{Q}$  is an isomorphism. Thus, we can use  $D_i$  to refer both to the invariant cycle and the associated  $\mathbb{Q}$ -Cartier divisor. In this case, as the title of the section advertises, we get a ring structure on  $Z_*(X)_\mathbb{Q} = \bigoplus_{k=0}^n Z_k(X)_\mathbb{Q}$ .

### 3.1 Definition of the Ring Structure

The definition of the ring structure proceeds in a somewhat unusual order, as in [2]. First, observe that by Theorem 2.1 (Commutativity),  $Z_*(X)_{\mathbb{Q}}$  is a module over  $\mathbb{Q}[Y_1, \dots, Y_r]$ , the polynomial ring in  $r$  variables ( $r$  being as always the number of rays of  $\Sigma$ ), where  $Y_i$  acts on  $Z_*(X)_{\mathbb{Q}}$  by  $D_i$ . (The simplicialness of  $\Sigma$  is used here to give us that the  $D_i$  are  $\mathbb{Q}$ -Cartier.)

The essential observation is that  $Z_*(X)_{\mathbb{Q}}$  is actually generated as a module over  $\mathbb{Q}[Y_1, \dots, Y_r]$  by  $[X]$ . Let  $\tau$  be a cone of dimension  $k$ , and renumber the rays of  $\Sigma$  so that  $\tau$  contains exactly  $\rho_1, \dots, \rho_k$ . Now consider  $D_{\rho_1} \cdots D_{\rho_k} \cdot [X]$ . Since the  $D_i$  meet properly, this is just their geometric intersection  $[V(\tau)]$ , up to multiplication by a non-zero rational. Thus  $[V(\tau)]$  is in the image of  $\mathbb{Q}[Y_1, \dots, Y_n] \cdot [X]$ , as desired. Since  $Z_*(X)_{\mathbb{Q}}$  is a cyclic module over  $\mathbb{Q}[Y_1, \dots, Y_r]$ ,  $Z_*(X)_{\mathbb{Q}} \cong \mathbb{Q}[Y_1, \dots, Y_r]/I$  for some ideal  $I$ . This puts a ring structure on  $Z_*(X)_{\mathbb{Q}}$ .

Naturally, we would like to give explicit generators for  $I$ . We begin the process in this subsection. Let  $\alpha$  denote the ring homomorphism  $\mathbb{Q}[Y_1, \dots, Y_n] \rightarrow Z_*(X)_{\mathbb{Q}}$ .  $I$  is the kernel of  $\alpha$ .

Define the ideal

$$I_{\Sigma} = \langle Y_{i_1} \cdots Y_{i_s} \mid \rho_{i_1}, \dots, \rho_{i_s} \text{ are not all contained in some cone of } \Sigma \rangle.$$

This is a Stanley-Reisner ideal, see [9]. For now, we prove the following lemma:

**Lemma 3.1**  $I_{\Sigma}$  is contained in  $I$ .

**Proof** This is a triviality. Let  $Y_{i_1} \cdots Y_{i_s}$  be a generator of  $I_{\Sigma}$ . Its image under  $\alpha$  is  $D_{i_1} \cdots D_{i_s}$ , which must consist of a linear combination of  $[V(\tau)]$  with  $\tau$  containing all the  $\rho_{i_j}$  — but there are no such  $\tau$  by assumption. So its image is zero, and it is in  $I$ . ■

### 3.2 A Presentation for $Z_*(X)_{\mathbb{Q}}$ Defined by an Inner Product Action

We can say more about  $I$  in the case where  $\Psi$  is induced from an inner product on  $M$ .

The inner product on  $M$  determines an isomorphism  $\omega: N_{\mathbb{Q}} \rightarrow M$  by requiring that  $\langle \cdot, \omega(v) \rangle_M = \langle \cdot, v \rangle$ , where the first set of angle brackets refer to the inner product in  $M$ , and the second to the duality pairing.

Fix  $j, 1 \leq j \leq r$ . Let  $E_j$  be the principal  $\mathbb{Q}$ -Cartier divisor whose local equation on each maximal  $\sigma$  is  $\omega(v_j)$ . I claim that  $E_j \cdot D_j = 0$ . We compute  $E_j \cdot D_j$  by viewing  $E_j$  as a  $\mathbb{Q}$ -Cartier divisor, and  $D_j$  as a cycle. By the definition of  $\omega$ ,  $\omega(v_j)$  is perpendicular in  $M$  to  $v_j^{\perp} = \rho_j^{\perp}$ . Thus,  $\pi_{\rho_j}(\omega(v_j)) = 0$ , so the divisor on  $D_j$  obtained from  $E_j$  is zero, so  $E_j \cdot D_j = 0$ .

We wish to convert this into a statement about  $I$ , the kernel of the map  $\alpha: \mathbb{Q}[Y_1, \dots, Y_r] \rightarrow Z_*(X)_{\mathbb{Q}}$ . To do this, we must write  $E_j$  as a sum of  $D_i$ . It is easy to see that

$$E_j = \sum_{i=1}^r \langle \omega(v_j), v_i \rangle D_i.$$

So, if we let  $J$  be the ideal in  $\mathbb{Q}[Y_1, \dots, Y_n]$  generated by  $\sum_{j=1}^r \langle \omega(v_j), v_i \rangle Y_i Y_j$  for  $1 \leq i \leq r$ , then  $J$  is contained in  $I$ . (The argument above, largely due to William Fulton, is a simplification of my original argument.)

Now we can give a presentation of  $Z_*(X)_{\mathbb{Q}}$ .

**Theorem 3.1** *When  $\Psi$  is induced from an inner product on  $M$ ,  $I = I_{\Sigma} + J$ , i.e.,*

$$Z_*(X)_{\mathbb{Q}} = \mathbb{Q}[Y_1, \dots, Y_n]/(I_{\Sigma} + J).$$

**Proof** As we already remarked, we know that  $I_{\Sigma} + J$  is contained in  $I$ . To show the other containment, we proceed as in [2].

For  $\sigma \in \Sigma$ , denote by  $Y_{\sigma}$  the product of those  $Y_i$  such that  $\rho_i$  is a ray of  $\sigma$ . The collection of all the  $Y_{\sigma}$  form a  $\mathbb{Q}$ -basis for  $\mathbb{Q}[Y_1, \dots, Y_n]/I$ . So, given any monomial in  $\mathbb{Q}[Y_1, \dots, Y_n]$ , it suffices to show that it is congruent to a linear combination of  $Y_{\sigma} \pmod{I_{\Sigma} + J}$ .

Let  $T$  be a monomial in  $\mathbb{Q}[Y_1, \dots, Y_n]$ . Let  $|T|$  denote the set of  $i$  such that  $Y_i$  appears in  $T$ . Let  $|\sigma|$  denote the set of  $i$  such that  $\rho_i$  is contained in  $\sigma$ . Now, we divide into two cases. First suppose that  $|T|$  does not equal  $|\sigma|$  for any  $\sigma \in \Sigma$ . Then  $T$  is divisible by one of the generators of  $I_{\Sigma}$ , so it is congruent to 0 mod  $I_{\Sigma}$ , and we are done.

Now suppose that  $|T| = |\sigma|$  for some  $\sigma$ . We first prove as a lemma the desired reducibility for certain values of  $T$ , and then use that to prove the general result.

**Lemma 3.2** *For  $i \in |\sigma|$ ,  $Y_i Y_{\sigma}$  is congruent to a linear combination of  $Y_{\tau}$  mod  $I_{\Sigma} + J$ .*

**Proof** Renumbering, we may assume that the  $\rho_j$  which are in  $\sigma$  are numbered 1 through  $k$ . Then, for  $1 \leq j \leq k$ ,

$$\sum_{i=1}^r \langle \omega(v_j), v_i \rangle Y_i Y_{\sigma} \in J.$$

Equivalently, for  $1 \leq j \leq k$

$$(2) \quad \sum_{1 \leq i \leq k} \langle \omega(v_j), v_i \rangle Y_i Y_{\sigma} \equiv - \sum_{k+1 \leq i \leq r} \langle \omega(v_j), v_i \rangle Y_i Y_{\sigma} \pmod{J}.$$

Observe that the monomials on the right-hand side are square-free.

We express (2) in matrix notation. Let  $A$  be the  $k \times k$  matrix whose entries are  $a_{ji} = \langle \omega(v_j), v_i \rangle$ . Let  $W$  be the  $k \times 1$  matrix  $w_{i1} = Y_i Y_{\sigma}$ . Let  $B$  be the  $k \times (n - k)$  matrix  $b_{ji} = \langle \omega(v_j), v_{i+k} \rangle$ . Let  $U$  be the  $(n - k) \times 1$  matrix  $u_{i1} = Y_{i+k} Y_{\sigma}$ .

Then (2) can be rewritten as

$$AW \equiv -BU \pmod{J}.$$

As in [2], the critical fact is that the matrix  $A$  is invertible. We prove this as follows. Extend the inner product on  $M$  to an inner product on  $M_{\mathbb{R}} = \mathbb{R} \otimes M$ . Pick an

orthonormal basis for  $M_{\mathbb{R}}$  and let  $V$  be the  $k \times n$  matrix whose  $i$ -th row is  $\omega(v_i)$  written out in that basis. Then  $A = VV^t$ . Since  $V$  has rank  $k$ , by the Cauchy-Binet formula  $A$  has non-zero determinant, and is therefore invertible.

So we can write

$$W = -A^{-1}BU \pmod J,$$

which gives us an expression for  $Y_i Y_\sigma$  ( $1 \leq i \leq k$ ) as a linear combination of square-free monomials. This suffices because every square-free monomial is either in  $I_\Sigma$  or equals  $Y_\tau$  for some  $\tau \in \Sigma$ . So for  $1 \leq i \leq k$ ,  $Y_i Y_\sigma$  is congruent to a linear combination of  $Y_\tau \pmod{I_\Sigma + J}$ , which proves the lemma. ■

Returning to the general case, let  $T$  be a monomial in  $\mathbb{Q}[Y_1, \dots, Y_n]$  with  $|T| = \sigma$ , and let  $T = \prod_{i \in |\sigma|} Y_i^{a_i}$ . Define its excess exponent to be  $e(T) = \sum_{i \in |\sigma|} (a_i - 1)$ . So a monomial is square-free iff its excess exponent is zero. Now, using Lemma 3.2, it is easy to see that we can express  $T$  as a sum of terms with smaller excess exponents, and this suffices by induction. ■

### 3.3 A Hard Lefschetz-Type Theorem for $Z_*(X)_{\mathbb{Q}}$ Defined by an Inner Product Action

Let  $\Sigma$  be simplicial. In this subsection we establish a hard Lefschetz-type theorem for  $Z_*(X)_{\mathbb{Q}}$ , where  $\Psi$  is induced from a “generic” inner product, an expression which we immediately explain. By fixing a basis for  $M$ , we can identify the set of inner products on  $M$  with the set  $P$  of  $n \times n$  symmetric matrices with coefficients in  $\mathbb{Q}$  and all eigenvalues positive. The set of  $n \times n$  symmetric matrices can be thought of as a finite vector space over  $\mathbb{Q}$ , and as such has both a Hausdorff and a Zariski topology.  $P$  is easily seen to be an open set in the Hausdorff topology. We say that a property holds for a generic inner product if there is a non-empty Zariski-open subset  $S$  of the symmetric matrices such that any matrix in  $P \cap S$  induces an inner product on  $M$  which has the desired property. (In particular, because of the nature of the Zariski topology, this guarantees that there is a non-empty Hausdorff open subset of  $P$  such that any matrix in the subset induces an inner product with the desired property.)

The statement of the theorem is as follows:

**Theorem 3.2** *Let  $\Sigma$  be a simplicial fan. Let  $\xi = \sum_{i=1}^r a_i D_i$ , with all the  $a_i$  non-zero. Then, for a generic inner product on  $M$ , the map from  $Z_{n-i}(X)_{\mathbb{Q}}$  to  $Z_i(X)_{\mathbb{Q}}$  induced by multiplication by  $\xi^{n-2i}$  is an injection for  $0 \leq i \leq n/2$ .*

**Proof** Usually a hard Lefschetz theorem would say that  $\xi^{n-2i}$  should induce an isomorphism from  $Z_{n-i}(X)_{\mathbb{Q}}$  to  $Z_i(X)_{\mathbb{Q}}$  but we cannot hope for that because the spaces usually have different dimensions. If they have the same dimensions (as for instance in the affine case) then injection and isomorphism are of course the same thing.

First, we prove a very special case.

**Lemma 3.3** *Let  $\Sigma$  be affine simplicial, let  $\xi = \sum_{i=1}^n a_i D_i$ , with all the  $a_i$  non-zero, and let the inner product be such that the  $v_i$  are all orthogonal to one another. Then the map induced by multiplication by  $\xi^{n-2i}$  from  $Z_{n-i}(X)_{\mathbb{Q}}$  to  $Z_i(X)_{\mathbb{Q}}$  is an isomorphism.*



**Proof** The choice of inner product means that

$$Z_*(X)_{\mathbb{Q}} \cong \mathbb{Q}[Y_1, \dots, Y_n] / \langle Y_1^2, \dots, Y_n^2 \rangle,$$

where the isomorphism takes  $D_i$  to  $Y_i$ .

Let  $W_i = a_i Y_i$ .  $\mathbb{Q}[Y_1, \dots, Y_n] / \langle Y_1^2, \dots, Y_n^2 \rangle \cong \mathbb{Q}[W_1, \dots, W_n] / \langle W_1^2, \dots, W_n^2 \rangle$ . For  $K \subset \{1, \dots, n\}$ , let  $W_K$  be the product of the  $W_i$  with  $i \in K$ . Then the  $W_K$  form a basis for  $Z_*(X)_{\mathbb{Q}}$  in which multiplication is particularly simple:  $W_K W_L = W_{K \cup L}$  if  $K \cap L = \emptyset$  and 0 otherwise. In this basis  $\xi = W_1 + \dots + W_n$ .

To prove the lemma, it suffices to show that for any  $i \leq n/2$ , and any set  $K$  of size  $n - i$ ,  $W_K$  is a multiple of  $\xi^{n-2i}$ . A quick proof, due to Matt Frank, is as follows. We can assume  $K = \{1, \dots, n - i\}$ . Let  $\zeta = W_{n-i+1} + \dots + W_n$ . Then

$$W_K = \frac{1}{(n-i)!} (\xi - \zeta)^{n-i} = \frac{1}{(n-i)!} \sum_{j=0}^{n-i} \binom{n-i}{j} \xi^{n-i-j} \zeta^j.$$

But for  $j > i$ ,  $\zeta^j = 0$ . So

$$W_K = \frac{1}{(n-i)!} \sum_{j=0}^i \binom{n-i}{j} \xi^{n-i-j} \zeta^j = \frac{\xi^{n-2i}}{(n-i)!} \sum_{j=0}^i \binom{n-i}{j} \xi^{i-j} \zeta^j.$$

This proves the lemma. ■

Keep  $\Sigma$  fixed and affine, and consider the set of inner products such that for  $0 \leq i \leq n/2$ ,  $\xi^{n-2i}$  induces an isomorphism from  $Z_{n-i}(X)_{\mathbb{Q}}$  to  $Z_i(X)_{\mathbb{Q}}$ . The conditions which define this subset amount to the non-vanishing of certain determinants, and are thus Zariski-open conditions on the set of symmetric matrices. The lemma proves that the set is non-empty. So there is a non-empty Zariski-open subset  $S$  of the symmetric matrices such that any element of  $S \cap P$  induces an inner product with the desired property. Thus, a generic inner product on  $M$  has the property that  $\xi^{n-2i}$  induces an isomorphism between  $Z_{n-i}(X)_{\mathbb{Q}}$  and  $Z_i(X)_{\mathbb{Q}}$ .

Now we return to the proof of the theorem as originally stated, and let  $\Sigma$  be any simplicial fan. The condition that  $\xi^{n-2i}$  induces an injection can be checked locally on affine cones, and thus the set of inner products with the desired property is an intersection of non-empty Zariski-open sets, and hence is itself non-empty and Zariski-open. ■

### 3.4 Example: Projective Space

In this subsection, we work out the details of the ring structure on the invariant cycles for a particular realization of projective space as a toric variety, and we compute a cycle representing its Todd class.

**Convention** For the duration of this example, we rescind our convention that  $M$  means  $\mathbb{Q} \otimes M$  — we will do our tensoring explicitly.

Let  $N' = \mathbb{Z}^{n+1}$  with basis vectors  $e_1, \dots, e_{n+1}$ . Let  $N = \{\sum_{i=1}^{n+1} a_i e_i \mid \sum a_i = 0\}$ . Let  $v_i = e_i - e_{i+1}$ , interpreted cyclically, so that  $v_{n+1} = e_{n+1} - e_1$ . Define a fan  $\Sigma$  in  $N$  consisting of all cones  $\sigma$  generated by a proper subset of  $\{v_1, \dots, v_{n+1}\}$ . Then  $v_i$  is the first lattice point along the ray generated by  $v_i$ , which we call  $\rho_i$ . We denote by  $D_i$  the Cartier divisor corresponding to  $\rho_i$ . Let  $W$  be the toric variety associated to  $\Sigma$ . Then  $W$  is isomorphic to  $\mathbb{P}^n$ .

Let  $M'$  be dual to  $N'$ , and  $e_i^*$  its dual basis. Let  $M$  be dual to  $N$ . Then  $M = M'/\mathbb{Z}(e_1^* + \dots + e_n^*)$ . Let  $\bar{e}_i^*$  denote the image of  $e_i^*$  in  $M$ .

We put the standard inner product on  $M'_\mathbb{Q}$ , and identifying  $M_\mathbb{Q}$  with the orthogonal complement of  $e_1^* + \dots + e_{n+1}^*$ , we induce an inner product on  $M_\mathbb{Q}$ . By the results of Section 3.2, we see that the ring on the invariant cycles is as follows:

$$Z_*(W)_\mathbb{Q} \cong \mathbb{Q}[Y_1, \dots, Y_{n+1}] / (\langle Y_i(Y_i - \frac{1}{2}Y_{i-1} - \frac{1}{2}Y_{i+1}) \mid 1 \leq i \leq n+1 \rangle + \langle Y_1 Y_2 \dots Y_{n+1} \rangle).$$

This can also be seen by direct calculation of  $D_i^2$  in  $Z_*(W)_\mathbb{Q}$ ; the reader is encouraged to try this out, to get the flavour of the computation of intersection cycles.

$\text{Td}(X)$ , the Todd class of  $X$ , is an element of  $A_*(X)_\mathbb{Q}$ . If  $X$  is a non-singular toric variety, it satisfies the following formula (see [4]):

$$\text{Td}(X) = \prod_{i=1}^r \frac{D_i}{1 - \exp(-D_i)},$$

where the computation takes place in the Chow ring of  $X$ . By interpreting this as taking place in the ring  $Z_*(X)_\mathbb{Q}$ , we may obtain a cycle-level Todd class for any non-singular variety. For a non-singular variety  $X$ , let  $t(X)$  denote the cycle-level Todd class obtained in this way.

Suppose that we have an inner product on  $N_\mathbb{Q}$ . For  $\sigma \in \Sigma$ , let  $B_\sigma$  be the ball of radius 1 about the origin in  $N_\sigma$ . Let  $\text{vol}$  denote Lebesgue measure on  $N_\sigma \otimes \mathbb{R}$ . Then we say that the fraction of the linear span of  $\sigma$  which it occupies is

$$\text{vol}(\sigma \cap B_\sigma) / \text{vol}(B_\sigma).$$

**Theorem 3.3** *Let  $W$  be the above realization of  $\mathbb{P}^n$  as a toric variety, with fan  $\Sigma$ . For  $\sigma$  any cone in  $\Sigma$ , the coefficient of  $[V(\sigma)]$  in  $t(W)$  is the fraction of the linear span of  $\sigma$  which is contained in  $\sigma$ .*

**Proof** In [2], a ring on the cycles of  $W$  is constructed in which  $D_i^2 = D_i \cdot D_{i+1}$  (with indices interpreted cyclically), and it is shown there that the cycle-level Todd class in this ring has the property described in the statement of the theorem. The essential details of the proof in [2] go through for our ring without any changes.

Define the formal power series:

$$\Phi(x) = \frac{x}{1 - e^{-x}}$$

$$Y(x) = -\log(1 - x) = x + \frac{x^2}{2} + \frac{x^3}{3} + \dots$$

Observe that  $x\Phi(Y(x)) = Y(x)$ .

We consider the set  $\{1, \dots, n + 1\}$  arranged around a circle. If  $S$  is a subset of  $\{1, \dots, n + 1\}$ , we call the components of  $S$  its maximal subsets of consecutive integers (understood cyclically). Define  $q(S) = \prod \frac{1}{|T|+1}$  where the product is taken over all components  $T$  of  $S$ . Let  $D_S = \prod_{i \in S} D_i$ .

**Lemma 3.4** *In  $Z_*(W)_{\mathbb{Q}}$ ,*

$$t(W) = \prod_{i=1}^{n+1} \Phi(D_i) = \sum_{S \subsetneq \{1, \dots, n+1\}} q(S)D_S.$$

**Proof** We think of the monomials  $D_S$  for  $S \subsetneq \{1, \dots, n + 1\}$  as a  $\mathbb{Q}$ -basis for the ring  $Z_*(W)_{\mathbb{Q}}$ . Let  $r(S)$  be the coefficient of  $D_S$  in the above product. Note that since the monomials produced by multiplying a monomial by  $D_i$  all include  $D_i$ ,  $r(S)$  is unchanged if we omit all the  $\Phi(D_i)$  for  $i \notin S$ . Also, it is clear that  $r(S) = \prod r(T)$  as  $T$  ranges over the components of  $S$ . Since this is also true of  $q(S)$ , it suffices to consider  $S$  connected, so by symmetry we may assume that  $S = \{1, \dots, k\}$ . By the same argument as above, we may assume that  $n = k$ .

Now consider the map from  $Z_*(W)_{\mathbb{Q}}$  to  $\mathbb{Q}[x]$  which takes all the  $D_i$  to  $x$ . Then the sum of  $q(S)$  over all subsets  $S$  of size  $n$  of  $\{1, \dots, n + 1\}$  is the coefficient of  $x^n$  in  $\Phi(x)^{n+1}$ .

Given a power series  $f(x)$ , denote the  $x^n$  coefficient of  $f(x)$  by  $[x^n]f(x)$ . Then by the Lagrange Inversion Formula (see [12]),

$$(n + 1)[x^{n+1}]Y(x) = [x^n]\Phi(x)^{n+1}.$$

So  $[x^n]\Phi(x)^{n+1} = 1$ . So the sum over all the subsets  $S$  of size  $n$  of  $\{1, \dots, n + 1\}$  is 1. By symmetry, then, for any such  $S$ ,  $r(S) = \frac{1}{n+1}$ , so  $r(S) = q(S)$  as desired. ■

Thus, for a cone  $\sigma \in \Sigma$ , define  $S$  to be the indices of the rays of  $\Sigma$  contained in  $\sigma$ . Then the coefficient of  $[V(\sigma)]$  in  $t(W)$  is  $q(S)$ , and, as stated in [2], this is the fraction of the linear span of  $\sigma$  which is contained in  $\sigma$ , which proves the theorem. ■

## 4 Computation

In this section, we address the matter of doing computations using our action.

### 4.1 Computing $D^n$

In this subsection, we compute the coefficient of  $[V(\sigma)]$  in  $D^n$ , where  $\sigma$  is an  $n$ -dimensional cone of  $\Sigma$ , and express it as the sum of plus or minus the volume of a large number of simplices. The result is somewhat unwieldy, but gives the basic geometrical picture which we shall exploit in the remainder of this section.

We begin with some very elementary statements about volumes. Given a simplex  $S$  with vertices  $u_0, \dots, u_n$  in  $M$ , consider the matrix  $A$  whose  $i, j$ -th entry is given by

$\langle u_j - u_0, v_i \rangle$ , where the  $\{v_j\}$  are a basis of  $N$ . Then the lattice volume of  $S$ , written  $\text{vol}(S)$  or  $\text{vol}(u_0, u_1, \dots, u_n)$  is  $\frac{1}{n!} |\det(A)|$ . This is a well-defined element of  $\mathbb{Q}$ , and doesn't depend on the order of the vertices. In general, when we refer to volume, we mean volume normalized in this way with respect to the lattice. Note that this is only possible if the affine span of the set whose volume we wish to take is rational, *i.e.*, is parallel to some sublattice of the group  $M$  (not meaning  $\mathbb{Q} \otimes M$  here), so that there is a well-defined lattice in the affine span of the set.

**Lemma 4.1** *Let  $\rho$  be a rational ray in  $N$ ,  $v$  the first lattice point along it. Suppose  $u_0, \dots, u_{n-1} \in \rho^\perp$ . Then*

$$\text{vol}(u_0, u_1, \dots, u_n) = \frac{1}{n} \langle u_n, v \rangle \text{vol}(u_0, \dots, u_{n-1}).$$

**Corollary** *Let  $\rho$  be a rational ray in  $N$ ,  $v$  the first lattice point along it. Let  $P$  be a pyramid in  $M$  over a polytope  $Q$  in  $\rho^\perp$ , with apex  $u_n$ . Then*

$$\text{vol}(P) = \frac{1}{n} \langle u_n, v \rangle \text{vol}(Q).$$

Let  $D$  be a  $\mathbb{Q}$ -Cartier divisor on  $X$ , a general  $n$ -dimensional toric variety. Let  $\sigma$  be an  $n$ -dimensional cone in  $\Sigma$ , the fan corresponding to  $X$ . We will compute the coefficient of  $[V(\sigma)]$  in  $D^n$ . Let the local equation for  $D$  on  $\sigma$  be  $m \in M$ .

Clearly, the coefficient of  $[V(\sigma)]$  in  $D^n$  is

$$\sum_{0=\gamma_0 \prec \gamma_1 \prec \dots \prec \gamma_n=\sigma} \prod_{1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle$$

Now we analyze the contribution from each term in the sum.

**Lemma 4.2** *Fix a chain  $0 = \gamma_0 \prec \gamma_1 \prec \dots \prec \gamma_n = \sigma$ . The absolute value of the contribution from this chain to the coefficient of  $[V(\sigma)]$ ,*

$$\left| \prod_{1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle \right|$$

*is  $n!$  times the lattice volume of the simplex with vertices  $\pi_{\gamma_0}(m), \pi_{\gamma_1}(m), \dots, \pi_{\gamma_n}(m)$ .*

**Proof** The proof is by induction on  $n$ . For  $n = 1$  it is clear. Suppose it is true for  $n - 1$ . Then  $\left| \prod_{1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle \right|$  is  $(n - 1)!$  times the lattice volume of the simplex with vertices  $\pi_{\gamma_1}(m), \dots, \pi_{\gamma_n}(m)$ .

We apply Lemma 4.1, with  $\gamma_1$  for  $\rho$ . Observe that  $v$  is  $n_{\gamma_1, \gamma_0}$ . The result follows. ■

Suppose that  $X$  is a proper toric variety (which is equivalent to saying that the support of  $\Sigma$  is all of  $N_{\mathbb{Q}}$ ), and  $D$  a  $\mathbb{Q}$ -Cartier divisor on it with local equations  $m_\sigma$ . Let  $S$  be the set of all  $-m_\sigma \in M$ , for  $\sigma$  a maximal cone. Form the convex hull of

$S$ , and call it  $P$ . If there is a 1–1 inclusion-reversing map  $F$  from the cones of  $\Sigma$  to the faces of  $P$ , which extends the map from the maximal cones to  $S$ , we say that the divisor  $D$  corresponds to the polytope  $P$ .

In this case, we see that if we add together the simplices of Lemma 4.2, taking care to cancel overlaps with opposite signs, we obtain  $P$ , from which we may recover the well-known fact that for a Cartier divisor corresponding to  $P$ ,  $\text{deg}(D^n) = n! \text{vol}(P)$ .

### 4.2 Computation for the Generic Flag Action

We can obtain better results when we specialize to the generic flag action. Our goal is the following theorem:

**Theorem 4.1** *Let  $\sigma$  be a  $k$ -dimensional simplicial cone in  $\Sigma$ , let  $E_1, \dots, E_s$  be  $\mathbb{Q}$ -Cartier divisors with local equations  $m_i$  on  $\sigma$ , and let  $q_k(x_1, \dots, x_s)$  be a homogeneous polynomial of degree  $k$ . Let  $w$  be a non-zero vector in  $N_\sigma$  in the direction perpendicular to  $L_{k-1}$ . Let  $t_1, \dots, t_k$  be chosen in  $M_\sigma$  to lie on the extreme rays of the image of  $\sigma$  in  $M_\sigma$ , so that the lattice volume of the simplex with vertices  $0, t_1, \dots, t_k$  is  $1/k!$ .*

*Then the coefficient of  $[V(\sigma)]$  in  $q_k(E_1, \dots, E_s)$  is*

$$\frac{q_k(\langle m_1, w \rangle, \dots, \langle m_s, w \rangle)}{\langle t_1, w \rangle \cdots \langle t_k, w \rangle}.$$

We begin by investigating the coefficient of  $[V(\sigma)]$  in  $D^n$  for  $\Psi$  given by a generic flag in  $M$ , under the assumption that  $\sigma$  is simplicial.

Let  $0 = L_0 \subset L_1 \subset \dots \subset L_n = M$  be a complete generic flag in  $M$ , and  $\Psi$  the corresponding choice of complements. Let  $\sigma$  be an  $n$ -dimensional simplicial cone. Let  $D$  be a  $\mathbb{Q}$ -Cartier divisor, and let its local equation on  $\sigma$  be  $m$ . Let the facets of  $\sigma$  be  $\tau_1, \dots, \tau_n$ , and the extreme rays of  $\sigma$  be  $\rho_1, \dots, \rho_n$ , with  $\rho_i$  the unique extreme ray not contained in  $\tau_i$ . Let  $\sigma^\vee$  be the dual cone to  $\sigma$  — by definition this is the cone consisting of elements of  $M$  whose duality pairing with all the elements of  $\sigma$  is non-negative. For  $\gamma \prec \sigma$ , let  $\gamma^* = \gamma^\perp \cap \sigma^\vee$ . The  $\tau_i^*$  are the extreme rays of the cone  $\sigma^\vee$ , and the  $\rho_i^*$  are the facets of  $\sigma^\vee$ .

A simplex in  $M$  determines  $n + 1$  hyperplanes, the affine spans of its facets. Conversely, hyperplanes  $H_1, \dots, H_{n+1}$  in general position determine a simplex, as follows. Let  $v_i$  be the intersection of all the hyperplanes excluding  $H_i$ . Then we say that the collection of hyperplanes determines the simplex which is the convex hull of the  $v_i$ .

We prove the following proposition:

**Proposition 4.1** *Let  $\sigma$  be an  $n$ -dimensional simplicial cone. The coefficient of  $[V(\sigma)]$  in  $D^n$  is plus or minus  $n!$  times the volume of the simplex corresponding to the affine spans of the  $\rho_i^*$  and the hyperplane  $H$  passing through  $m$  parallel to  $L_{n-1}$ . It is positive or negative as the number of  $\tau_i^*$  which  $H$  does not intersect is even or odd.*

**Proof** The proof is by induction on  $k$ . The induction claim is that if  $\delta$  is an  $n - k$ -dimensional face of  $\sigma$  (so, by renumbering, we may assume that  $\delta$  is the intersection of  $\tau_1, \dots, \tau_k$ ), then

$$\frac{1}{k!} \sum_{\delta=\gamma_{n-k} \leftarrow \dots \leftarrow \gamma_n = \sigma} \prod_{n-k+1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle$$

is plus or minus the volume of the simplex in  $\delta^\perp$  corresponding to the hyperplanes  $\delta^\perp \cap \rho_1^*, \dots, \delta^\perp \cap \rho_k^*, \delta^\perp \cap H$ , signed positive or negative as the number of  $\tau_i^*$  for  $1 \leq i \leq k$  that  $\delta^\perp \cap H$  misses is even or odd.

The case  $k = 0$  is true by convention, and  $k = n$  is the desired result. Assume the statement holds for  $k - 1$ . Then

$$\begin{aligned} (3) \quad & \frac{1}{k!} \sum_{\delta=\gamma_{n-k} \leftarrow \dots \leftarrow \gamma_n = \sigma} \prod_{n-k+1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle \\ &= \frac{1}{k} \sum_{i=1}^k \left( \frac{\langle \pi_\delta(m), n_{\delta+\rho_i, \delta} \rangle}{(k-1)!} \sum_{\substack{\delta+\rho_i= \\ \gamma_{n-k+1} \leftarrow \dots \leftarrow \gamma_n = \sigma}} \prod_{j=n-k+2}^n \langle \pi_{\gamma_{j-1}}(m), n_{\gamma_j, \gamma_{j-1}} \rangle \right) \end{aligned}$$

Let  $S$  be the simplex in  $\delta^\perp$  corresponding to the hyperplanes  $\delta^\perp \cap \rho_1^*, \dots, \delta^\perp \cap \rho_k^*$  and  $\delta^\perp \cap H$ . Let  $F_i$  be the face of  $S$  corresponding to the hyperplane  $\delta^\perp \cap \rho_i^*$ . Then by the induction hypothesis, for each  $i, 1 \leq i \leq k$ ,

$$\frac{1}{(k-1)!} \left| \sum_{\substack{\delta+\rho_i= \\ \gamma_{n-k+1} \leftarrow \dots \leftarrow \gamma_n = \sigma}} \prod_{n-k+2 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle \right| = \text{vol}(F_i).$$

I claim that  $\pi_\delta(m) \in H \cap \delta^\perp$ . By construction it is in  $\delta^\perp$ . To show it is in  $H$ , first observe that by the definition of  $H, m \in H$ . But now by construction,  $\pi_\delta(m) - m \in \Psi(\delta) = L_{n-k}$ .  $L_{n-k} \subset L_{n-1}$  except when  $k = 0$ , but then  $\pi_\delta(m) - m = 0$ . So  $\pi_\delta(m) - m \in L_{n-1}$ . So  $\pi_\delta(m) \in H$ , and hence in  $H \cap \delta^\perp$ .

Let  $C_i$  be the pyramid over  $F_i$  with apex  $\pi_\delta(m)$ . Then

$$\text{vol}(C_i) = \frac{1}{k} \left| \langle \pi_\delta(m), n_{\delta+\rho_i, \delta} \rangle \right| \text{vol}(F_i).$$

So (3) is a sum of plus or minus the volumes of the  $C_i$ . Let  $s \in \{1, -1\}$  be positive if the number of  $\tau_i^*$  with  $1 \leq i \leq k$  missed by  $H$  is even, and negative if the number missed is odd. Let  $s_i$  denote the sign such that

$$\frac{1}{k!} \langle \pi_\delta(m), n_{\delta+\rho_i, \delta} \rangle \sum_{\substack{\delta+\rho_i= \\ \gamma_{n-k+1} \leftarrow \dots \leftarrow \gamma_n = \sigma}} \prod_{n-k+1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle = s_i \text{vol}(C_i).$$

Observe that  $\langle \pi_\delta(m), n_{\delta+\rho_i, \delta} \rangle$  is positive if  $\pi_\delta(m)$  is on the positive side of  $\delta^\perp \cap \rho_i^*$ , that is, the same side as  $\tau_i^*$ . So  $s_i$  is positive if  $H$  hits  $\tau_i^*$  and  $\pi_\delta(m)$  is on the  $\tau_i^*$  side of  $\delta^\perp \cap \rho_i^*$  or if  $H$  misses  $\tau_i^*$  and  $\pi_\delta(m)$  is on the  $-\tau_i^*$  side of  $\delta^\perp \cap \rho_i^*$ . But this amounts to saying that  $s_i$  is positive iff  $\pi_\delta(m)$  is on the same side of  $\delta^\perp \cap \rho_i^*$  as the intersection of  $H$  with the linear span of  $\tau_i^*$ .

Now there are two cases. The first case is if  $\pi_d(m)$  is in  $S$ . Then for all  $i$ ,  $ss_i = 1$ . Then the cones  $C_i$  form a dissection of  $S$ , that is to say, their union is  $S$ , and they overlap only on boundaries. So

$$\begin{aligned} \frac{1}{k!} \sum_{\delta=\gamma_{n-k} \leftarrow \gamma_{n-k+1} \leftarrow \dots \leftarrow \gamma_n = \sigma} \prod_{n-k+1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle &= \sum_{i=1}^k s_i \text{vol}(C_i) \\ &= s \text{vol}(S). \end{aligned}$$

The second case is if  $\pi_\delta(m)$  is not in  $S$ . Then let  $Q$  be the convex hull of  $S$  and  $\pi_\delta(m)$ . We see that the  $C_i$  with  $ss_i$  positive form a dissection of  $Q$ , while the  $C_i$  with  $ss_i$  negative form a dissection of  $Q \setminus S$ . So

$$\begin{aligned} \frac{1}{n!} \sum_{\delta=\gamma_{n-k} \leftarrow \gamma_{n-k+1} \leftarrow \dots \leftarrow \gamma_n = \sigma} \prod_{n-k+1 \leq i \leq n} \langle \pi_{\gamma_{i-1}}(m), n_{\gamma_i, \gamma_{i-1}} \rangle \\ &= \sum_{i=1}^k s_i \text{vol}(C_i) \\ &= s(\text{vol}(Q) - \text{vol}(Q \setminus S)) \\ &= s(\text{vol}(S)). \end{aligned}$$

This finishes the induction step, and the proof of the proposition. ■

We next reformulate Proposition 4.1 algebraically.

**Proposition 4.2** *Let  $\sigma$  be an  $n$ -dimensional simplicial cone in  $\Sigma$ ,  $\tau_1, \dots, \tau_n$  the facets of  $\sigma$ , and  $D$  a  $\mathbb{Q}$ -Cartier divisor, with local equation  $m$  on  $\sigma$ . Let  $w \in N_{\mathbb{Q}}$  be a non-zero vector perpendicular to  $L_{n-1}$ . For  $1 \leq i \leq n$ , pick  $t_i \in \tau_i^*$ , such that the lattice volume of the simplex with vertices  $0, t_1, \dots, t_n$  is  $\frac{1}{n!}$ .*

*Then the coefficient of  $[V(\sigma)]$  in  $D^n$  is*

$$\frac{\langle m, w \rangle^n}{\langle t_1, w \rangle \cdots \langle t_n, w \rangle}.$$

**Proof** Consider how much we would need to stretch the simplex with vertices located at  $0, t_1, \dots, t_n$  along each of the  $\tau_i^*$  in order for it to coincide with the simplex from the statement of Proposition 4.1, which we denote  $S$ . The scaling factor along  $\tau_i^*$  is  $\langle m, w \rangle / \langle t_i, w \rangle$ . Thus, the volume of  $S$  is:

$$\text{vol}(S) = \frac{1}{n!} \left| \frac{\langle m, w \rangle^n}{\langle t_1, w \rangle \cdots \langle t_n, w \rangle} \right|.$$

Now, observe that the sign on the expression inside the absolute value signs is exactly that dictated by Proposition 4.1, which proves the proposition. ■

From this, we derive a corollary which holds for products of different Cartier divisors:

**Corollary** *Let  $\sigma$  be an  $n$ -dimensional simplicial cone in  $\Sigma$  and let  $E_1, \dots, E_n$  be  $\mathbb{Q}$ -Cartier divisors, with local equations  $m_1, \dots, m_n$  on  $\sigma$ . Pick  $w$  and  $t_1, \dots, t_n$  as in the theorem. Then the coefficient of  $[V(\sigma)]$  in  $E_1 \cdots E_n$  is*

$$\frac{\langle m_1, w \rangle \cdots \langle m_n, w \rangle}{\langle t_1, w \rangle \cdots \langle t_n, w \rangle}.$$

**Proof** First, by a straightforward algebraic identity,

$$E_1 \cdots E_n = \frac{1}{n!} \sum_{j=0}^n \sum_{\substack{S \subset \{1, \dots, n\} \\ |S|=j}} (-1)^{n-j} \left( \sum_{i \in S} E_i \right)^n.$$

Thus, the coefficient of  $[V(\sigma)]$  in  $E_1 \cdots E_n$  is

$$\begin{aligned} & \frac{1}{\langle t_1, w \rangle \cdots \langle t_n, w \rangle} \sum_{j=0}^n \sum_{\substack{S \subset \{1, \dots, n\} \\ |S|=j}} (-1)^{n-j} \left( \sum_{i \in S} \langle m_i, w \rangle \right)^n \\ &= \frac{\langle m_1, w \rangle \cdots \langle m_n, w \rangle}{\langle t_1, w \rangle \cdots \langle t_n, w \rangle}. \end{aligned}$$

This proves the corollary. ■

Theorem 4.1 now follows by the techniques of Section 2.3 for reducing the computation of the coefficient of an arbitrary cone to that of a cone of full dimension.

Theorem 4.1 is only directly applicable to computing the coefficient of  $[V(\sigma)]$  in a product of  $\mathbb{Q}$ -Cartier divisors when  $\sigma$  is simplicial. However, it could still be applied to computing the coefficient of  $[V(\sigma)]$  with  $\sigma$  non-simplicial, as follows.

Suppose  $X$  is a non-simplicial toric variety. Find a simplicial toric variety  $X'$  and a proper birational map  $f: X' \rightarrow X$ . (This amounts to choosing a simplicial subdivision of  $\Sigma$ .) Let  $E_1, \dots, E_k$  be  $\mathbb{Q}$ -Cartier divisors on  $X$ . Then by Theorem 2.2,

$$E_1 \cdots E_k \cdot [X] = f_*(f^*(E_1) \cdots f^*(E_k) \cdot [X']),$$

and  $f^*(E_1) \cdots f^*(E_k) \cdot [X']$  can be computed using Theorem 4.1.

## 5 Extensions and Applications

### 5.1 Canonical Constructions

From either the inner product action or the generic flag action there is a way to obtain an action of Cartier divisors on cycles which does not depend on any choice, but one has to allow the cycles to take coefficients in a much larger field. The construction we give is based on a construction in [6].



**Convention** For this section we revoke the convention of tacitly tensoring  $M$  and its sub- and quotient lattices by  $\mathbb{Q}$ .

First, we discuss the canonical inner product action. Let  $B(M)$  be the set of symmetric bilinear (not necessarily non-degenerate) forms on  $M_{\mathbb{Q}}$ .  $B(M)$  can be thought of as a variety defined over the rationals. Let  $\mathbb{K}$  be its rational function field. Let  $B^+(M)$  denote the positive definite forms in  $B(M)$ . Now, we wish to define a canonical action of Cartier divisors on cycles with coefficients in  $\mathbb{K}$ .

For  $D \in \text{Div}(X)$  and  $\sigma$  an  $n - k$  dimensional cone in  $\Sigma$ , we want to define  $D \cdot_{\text{IP}} [V(\sigma)]$  in  $Z_{k-1}(X)_{\mathbb{K}}$ , which we can think of as a rational function from  $B(M)$  to  $Z_{k-1}(X)_{\mathbb{Q}}$ . Since  $B^+(M)$  is Zariski dense in  $B(M)$ , to define a rational function on  $B(M)$  it suffices to define it on  $B^+(M)$ . So, for  $b \in B^+(M)$ , let  $(D \cdot_{\text{IP}} [V(\sigma)])(b) = D \cdot_b [V(\sigma)]$ , where  $\cdot_b$  denotes the inner product action relative to the inner product given by  $b$ . One easily sees that this is a rational function, and thus we can let this constitute the definition of  $D \cdot_{\text{IP}} [V(\sigma)]$ . Extending linearly, we obtain an action of  $\text{Div}(X)$  on  $Z_*(X)_{\mathbb{K}}$ .

This action has all the properties we proved for the inner product action (commutativity, agreement with usual intersection modulo rational equivalence, etc.) because all these properties hold for all  $b \in B^+(M)$ , which is Zariski dense in  $B(M)$ .

An interesting result of this construction is that it allows us to define an action of Cartier divisors on cycles with respect to a symmetric bilinear form which is not positive definite: simply evaluate the canonical action at that bilinear form. Note that this will not necessarily be well-defined, however, since the coefficients in question may blow up there.

The canonical flag action is defined in much the same way. The set  $\text{Fl}(M)$  of complete flags in  $M_{\mathbb{Q}}$  is a variety, and we let  $\mathbb{F}$  be its function field. For  $\mathcal{F} \in \text{Fl}(M)$ , we define

$$(D \cdot_{\text{Fl}} [V(\sigma)])(\mathcal{F}) = D \cdot_{\mathcal{F}} [V(\sigma)].$$

This will only be well-defined for  $\mathcal{F}$  generic with respect to  $\sigma$ , but this is an open set in  $\text{Fl}(M)$ , and therefore dense, so the same arguments go through. Thus, we have defined an action of Cartier divisors on cycles with coefficients in  $\mathbb{F}$ , and, as above, it has all the properties we are used to.

As an example, we state the result about the canonical flag action corresponding to Theorem 4.1.

**Theorem 5.1** *Let  $\sigma$  be a  $k$ -dimensional cone.  $E_1, \dots, E_s$  Cartier divisors on  $X$ , and  $q_k(x_1, \dots, x_s)$  a homogeneous polynomial of degree  $k$ . Let the local equation for  $E_i$  on  $\sigma$  be  $m_i \in M_{\sigma}$ . Let  $t_1, \dots, t_k$  be as in the statement of Theorem 4.1. Let  $w: \text{Fl}(M) \rightarrow \mathbb{Q} \otimes N_{\sigma}$ , where if  $\mathcal{F} = F_0 \leq \dots \leq F_n$ , then  $w(\mathcal{F})$  is a non-zero vector in  $\mathbb{Q} \otimes N_{\sigma}$  perpendicular to  $F_{k-1}$ .*

*Then the coefficient of  $[V(\sigma)]$  in  $q_k(E_1, \dots, E_s)$  is*

$$\frac{q_k(\langle m_1, w \rangle, \dots, \langle m_s, w \rangle)}{\langle t_1, w \rangle \cdots \langle t_k, w \rangle}.$$

## 5.2 Characteristic Classes of Toric Varieties

Let  $X$  be a simplicial toric variety. Suppose we have a characteristic class of  $X$  which is written as a polynomial  $p(D_1, \dots, D_r)$  interpreted in  $A_*(X)$ . (For example, the  $j$ -th Chern class of  $X$  can be written as the  $j$ -th symmetric polynomial on the  $D_i$ , and one can use this to obtain a polynomial in the  $D_i$  representing any class written as a polynomial in the Chern classes.) If we pick  $\Psi$  and interpret  $p(D_1, \dots, D_r)$  in  $Z_*(X)_{\mathbb{Q}}$ , we determine a cycle which represents the characteristic class in question. If  $\Psi$  was chosen using the generic flag action or the canonical flag action, we can compute this cycle quite explicitly using Theorem 4.1 or Theorem 5.1, respectively. We recover in this case the cycle-level characteristic classes obtained by Morelli in [6] using the Baum-Bott residue formula, but our result is somewhat more general: the results of [6] apply only when  $X$  is non-singular and compact.

If  $X$  is non-singular, one can use this approach to compute a cycle-level Todd class, as we have already seen in Section 3.4. (Determining cycle-level Todd classes is a matter of some interest, because of connections to counting lattice points in polytopes, see [1, 4, 6].)

For singular  $X$ ,  $\text{Td}(X)$  is computed by taking a resolution of singularities  $X' \rightarrow X$ , computing the Todd class of  $X'$ , and pushing it forward to  $X$ . This procedure can be done for the cycle-level Todd class of a toric variety as well, but *a priori* this introduces a dependence on the choice of resolution. In fact, under reasonable conditions, the resulting cycle-level Todd class does not depend on the choice of resolution, but the proof of this relies on results from [7] and is somewhat complicated, see [10, 8].

There is more work to be done when it comes to understanding cycle-level Todd classes of toric varieties. One would like to know, for instance, for what toric varieties Theorem 3.3 holds. Examples in [6] show that it does not hold for all toric varieties. On the other hand there is reason to think that it may hold for toric varieties arising from partially ordered sets, see [11].

**Acknowledgements** This paper is based on my thesis [10], written at the University of Chicago under the direction of William Fulton. I would like to thank him for his help in shaping its substance and its form. I would also like to thank Matt Frank and Ian Robertson for useful conversations, and an anonymous referee for his helpful comments.

## References

- [1] V. I. Danilov, *The geometry of toric varieties*. Russian Math. Surveys **33**(1978), 97–154; Uspekhi Mat. Nauk **33** (1978), 85–134.
- [2] P. Diaconis and W. Fulton, *A growth model, a game, an algebra, Lagrange inversion, and characteristic classes*. Rend. Sem. Math. Univ. Politec. Torino **49**(1991), 95–119.
- [3] W. Fulton, *Intersection theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete 3, Springer-Verlag, Berlin, 1984.
- [4] ———, *Introduction to toric varieties*. Annals of Mathematics Studies 131, Princeton University Press, Princeton, NJ, 1993.
- [5] W. Fulton and B. Sturmfels, *Intersection theory on toric varieties*. Topology **36**(1997), 335–353.
- [6] R. Morelli, *Pick's theorem and the Todd class of a toric variety*. Adv. Math. **100**(1993), 183–231.
- [7] J. Pommersheim, *Products of Cycles and the Todd Class of a Toric Variety*. J. Amer. Math. Soc. **9**(1996), 813–826.

- [8] J. Pommersheim and H. Thomas, *Cycles representing the Todd class of a toric variety*. J. Amer. Math. Soc. (to appear).
- [9] R. Stanley, *Combinatorics and commutative algebra*. Second edition, Progress in Mathematics 41, Birkhäuser, Boston, MA, 1996.
- [10] H. Thomas, *An action of equivariant Cartier divisors on invariant cycles for toric varieties*. Ph.D. Thesis, University of Chicago, 2000.
- [11] ———, *Order-preserving maps from a poset to a chain, the order polytope, and the Todd class of the associated toric variety*. European J. Combin. **24**(2003), 809–814.
- [12] H. Wilf, *generatingfunctionology*. Academic Press, Boston, MA, 1990.

*The Fields Institute*  
222 College St.  
Toronto, ON  
M5T 3J1  
e-mail: [hthomas@fields.utoronto.ca](mailto:hthomas@fields.utoronto.ca)