



Eating frequency and weight status in Portuguese children aged 3–9 years: results from the cross-sectional National Food, Nutrition and Physical Activity Survey 2015–2016

Sofia Vilela^{1,2,*}, Daniela Correia¹, Milton Severo^{1,2}, Andreia Oliveira^{1,2}, Duarte Torres^{1,3}, Carla Lopes^{1,2} and the IAN-AF Consortium†

¹EPIUnit, Instituto de Saúde Pública, Universidade do Porto, Rua das Taipas n° 135, 4050-600 Porto, Portugal:

²Departamento de Ciências da Saúde Pública e Forenses, e Educação Médica, Faculdade de Medicina, Universidade do Porto, Porto, Portugal: ³Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Porto, Portugal

Submitted 18 May 2018: Final revision received 17 January 2019: Accepted 7 February 2019: First published online 21 May 2019

Abstract

Objective: To evaluate daily eating frequency (main meals and snacks) in relation to weight status in children aged 3–9 years, representative of the Portuguese population.

Design: Cross-sectional study. Dietary intake was estimated as the mean of two non-consecutive days of food diaries, followed by face-to-face interviews. Weight and height were measured by trained observers. Eating occasions (EO) were defined by the children's caregiver; an EO was considered separate if the time of consumption was different from other EO and it provided at least 209 kJ (50 kcal). Main meals defined as 'breakfast', 'lunch' and 'dinner' could be selected only once per day. The remaining EO were considered snacks. The association between eating frequency and overweight/obesity was evaluated through logistic regressions weighted for the population distribution.

Setting: National Food, Nutrition and Physical Activity Survey of the Portuguese population, 2015–2016.

Participants: Portuguese children aged 3–9 years with complete dietary data and anthropometric measurements (*n* 517).

Results: Overall, the number of daily EO ranged from 3.5 to 11, and on average children had 5.7 daily EO. After adjustment for child's sex, age and total energy intake, and considering only plausible energy intake reporters, having < 3 snacks/d was positively associated with being overweight/obese (OR = 1.98; 95 % CI 1.00, 3.90), compared with having ≥ 3 snacks/d.

Conclusions: Lower daily frequency of EO was associated with increased odds of being overweight or obese in children. A higher eating frequency, maintaining the same energy intake, seems to contribute to a healthy body weight in children.

Keywords
Eating frequency
Snacks
Children
Obesity
Dietary surveys

The WHO estimates that over 340 million children and adolescents aged 5–19 years are classified as overweight or obese⁽¹⁾. Among pre-school children (< 5 years of age), the global prevalence of excessive weight (overweight and obesity) increased from 4.2% in 1990 to 6.7% in 2010. This trend is expected to reach 9.1%, representing 60 million children, in 2020⁽²⁾. Childhood obesity is a complex disease with several adverse sequelae, including psychological, social and health consequences^(3,4). Poor self-esteem, depression and eating disorders have been

described more extensively in obese children than in their normal-weight peers^(4–6). Furthermore, obese children are more susceptible to CVD risk factors such as hypertension⁽⁷⁾, dyslipidaemia⁽⁷⁾, chronic inflammation⁽⁸⁾, endothelial dysfunction^(9,10) and insulin resistance⁽¹¹⁾. Research has suggested that early childhood may be a particularly sensitive period for the development of obesity later in life⁽¹²⁾, thus early-life prevention of overweight and obesity should be valued.

There is evidence that meal pattern characteristics, such as irregularity of meals or 'skipping breakfast', could negatively influence diet quality^(13,14) and cardiometabolic health outcomes including obesity^(15,16). On the topic of eating frequency, increasing the number of daily meals

† Members of the IAN-AF Consortium are listed in the Appendix.

*Corresponding author: Email sofia.vilela@ispup.up.pt

might lead to a higher exposure to energy-dense foods, such as fast foods and soft drinks, and large-portion-size foods, resulting in increased hunger, excess energy intake and ultimately unhealthy body weight gain^(17–19). Nevertheless, a recent systematic review⁽²⁰⁾ of observational studies among adults included fourteen original studies reporting an inverse association between eating frequency and body weight or body composition, and seven studies reporting a positive association. Among children and adolescents, a previous meta-analysis⁽²¹⁾ described an overall inverse association between eating frequency and the likelihood of being overweight or obese. However, when stratified by sex, the effect remained significant only in boys. The presence of publication bias and heterogeneity of studies might in part explain these differences, and further research was advised. The inconsistency in definitions of eating frequency might also impact the associations with body weight and hamper data interpretation and comparison between studies⁽²²⁾.

There is no general agreement on the most appropriate definition of eating occasion, particularly the differentiation between what constitutes a main meal or a snack⁽²³⁾. The literature describes a variety of approaches to defining eating occasions (main meals and snacks), using, for example, the time of consumption ('time-of-day'), food groups or nutritional profile ('food-based classification') or identification by the participant ('participant-identified')⁽²³⁾. These approaches can be contradictory or complement one another depending on the context^(24,25).

A lack of public health recommendations on eating frequency or temporal distribution is common among European authorities⁽²⁶⁾. There is a lack of a universal definition of what constitutes an eating occasion, and for the differentiation between a main meal and a meal snack-type⁽²³⁾. The characterization of meal patterns among children using European harmonized methodology to collect dietary data and a national representative sample will support the improvement of the research field and aid the development of specific dietary guidelines.

The present study aimed to characterize eating frequency (daily number of eating occasions, main meals and snacks) in relation to weight status in Portuguese children aged 3–9 years, using data from a national dietary survey.

Materials and methods

Participants and design

The present study was conducted within the National Food, Nutrition and Physical Activity Survey of the Portuguese population, 2015–2016 (IAN-AF 2015–2016)⁽²⁷⁾.

The IAN-AF 2015–2016 aimed to collect nationwide and regional data on dietary habits and physical activity and to assess their relationship with health determinants, namely socio-economic factors. A representative sample of the Portuguese population, aged between 3 months and

84 years old, was selected from the National Health Registry by multistage sampling in each of the seven geographical regions (NUTS II), and weighted according to sex and age groups. Data were collected by trained interviewers using computer-assisted personal interviewing, in two interviews (8–15 d apart) during 12 months⁽²⁷⁾, distributed over the four seasons and including all days of the week, and following European standard guidelines⁽²⁸⁾. A total of 5811 participants completed two interviews and 6553 completed only the first one. Participants' characteristics were compared with participants who refused to participate but accepted to answer a short questionnaire by telephone. Non-participants were older (> 65 years: 22 *v.* 13%) and less educated (> 12 years: 19 *v.* 27%). Regarding variables representing the main research areas of the survey, the differences between participants and non-participants were considered of small magnitude: fruit and vegetable consumption (≥ 5 portions/d: 18.6 *v.* 18.1%), practice of regular structured physical activity (33 *v.* 39%) and obesity (12.4 *v.* 12.7%)⁽²⁷⁾. As the children younger than 3 years and adolescents had different methods of collection of dietary data⁽²⁷⁾, we decided to include only children aged 3–9 years. From the total children (< 10 years, *n* 1514), 1329 had complete data in the two face-to-face interviews. Our analysis included children 3–9 years of age (*n* 604) who had completed two interviews (*n* 521). Children's characteristics were also compared with children who were not allowed to participate. Non-participants were similar for consumption of fruits and vegetables (≥ 5 portions/d: 31.2 *v.* 31.6%) and for practice of regular structured physical activity (50.4 *v.* 51.2%); however, non-participants had a slightly higher prevalence of obesity (5.0 *v.* 2.4%). Four children did not have data on measured weight and height and were excluded, achieving a final sample of 517 children.

Dietary intake

For children under 10 years, dietary intake was obtained by two, non-consecutive, one-day food diaries followed by a face-to-face interview allowing the parent or another main caregiver to add details related to food description and quantification. The time between interviews was between 8 and 15 d and the days of reporting were selected (including all days of the week), but participants were able to change them, according to their own availability for the interview. During the face-to-face interview with a trained researcher, the food diaries completed by the caregiver in the previous day were included in the 'eAT24' module (Electronic Assessment Tool for 24-hours recall) which allowed the assessment of dietary data by an automated multiple-pass method for 24 h⁽²⁷⁾. All foods, including beverages and dietary supplements consumed during a 24 h period, were recorded per food consumption occasion (and quantified and described as eaten). The place and time of consumption were also recorded for each food consumption occasion. For food and recipe quantification several methods were used, such as standard unit method



(for foods consumed in distinguishable units with more or less standard weights), household measures and food photographs (a digital colour food picture book⁽²⁹⁾ was developed, based on a previously validated one⁽³⁰⁾, including 176 food photo series with six portions of each food/recipe item and eleven household measures photo series). The total children's energy intake was obtained through conversion of foods into nutrients by the software, using by default the Portuguese food composition table⁽³¹⁾.

Eating frequency

The 'eAT24' comprises twelve predefined food consumption occasions to standardize the reporting of events (intake of foods and/or drinks) during the 24 h period: 'before breakfast', 'breakfast', 'during morning', 'before lunch', 'lunch', 'after lunch', 'during afternoon', 'before dinner', 'dinner', 'after dinner', 'during evening/at night' and 'liquids intake'. All eating occasions reported in the food diaries were recorded and considered as separate intake occasions if the time of eating was different from another eating occasion; events consisting of water only (zero-energy foods) were excluded. To exclude negligible amounts of energy intake, eating occasions providing less than 209 kJ (50 kcal) were also excluded⁽²³⁾. Main meals defined as 'breakfast', 'lunch' and 'dinner' were unique and could be selected only once per day. The remaining eating occasions were considered snacks, and they could be selected more than once in case of having a snack on several occasions during the day.

Daily eating frequency describes the total number of separate eating occasions during a 24 h period, as the result of the average of the two reported days.

Anthropometrics

Children's weight and height were measured by trained observers, according to standard procedures⁽³²⁾. Weight was measured in the same conditions, to the nearest tenth of a kilogram, using a digital scale (SECA® 813, Hamburg, Germany). Height was measured to the nearest centimetre, with children in a standing position with light clothing and barefoot, using a portable wall stadiometer (SECA® 213). The child's age- and sex-specific BMI Z-score was calculated according to the WHO criteria⁽³³⁾. The BMI Z-score was categorized as underweight (< -2), normal weight (≥ -2 to ≤ 1) or overweight/obese (> 1), according to the WHO cut-off points⁽³³⁾.

Covariables

For socio-economic characterization, mother's and father's education and maternal age were considered. Parental education was defined as the maximum level of education of any of the parents and categorized as 'no formal education or primary education' (less than 6 years), 'secondary or post-secondary education' (between 6 and 12 years plus post-secondary education not including tertiary education) and 'tertiary education' (university degree, master and PhD).

Parents also reported if their child had regular practice of physical exercise (reported as non-practitioners *v.* practitioners) and the daily minutes spent watching television or playing video games during both weekdays and weekend days. This variable was combined; the number of minutes were averaged between weekdays and weekend days, converted into hours (categorized as watching television TV or playing video games for < 2 h/d, 2–3 h/d or ≥ 3 h/d) and used as an indicator of a more sedentary lifestyle⁽³⁴⁾. For children aged 6 years or older, physical activity was also assessed by an activity diary adapted from a model previously proposed⁽³⁵⁾, including two consecutive days during the week and two days on the weekend. This information was used only to classify dietary report of children aged 6 years or older into plausible reports or not. The diary consisted of a grid dividing each day (24 h) into 15 min periods. Children registered the primary activity performed in each 15 min interval, over the four days, according to previous written instructions. Each activity refers to a specific level of energy expenditure and was converted into metabolic ratios of expended energy. Individual daily energy expenditure was obtained as the mean expenditure of the 4 d diaries⁽²⁷⁾.

Statistical analysis

To evaluate the extent of misreporting of energy intake, the ratio of proxy-reported energy intake to predicted BMR was used to classify the dietary report into under-reports, plausible reports and over-reports, based on Goldberg cut-offs⁽³⁶⁾. The BMR was estimated using the equations published by Schofield⁽³⁷⁾, considering age, sex, body weight and height. The physical activity level was defined according to Torun⁽³⁸⁾ for children aged less than 6 years. For the remaining children, the individual physical activity level obtained by the diaries was used.

The characteristics of Portuguese children aged 3–9 years were examined according to BMI (non-overweight *v.* overweight/obese). Data were presented as prevalence estimates weighted by the inverse of its sampling probability for the Portuguese population distribution, according to the complex sampling design, considering in each region (NUTS II Statistical Geographic Units of Portugal) a cluster effect for the Primary Health Care Unit selected and stratification by sex and age groups⁽²⁷⁾. Mixed weighting was used to correct proportions of the sample in order to coincide to the population proportions and to expand to the population size. The respective 95 % CI were also obtained. Prevalence estimates according to socio-economic variables, eating occasions (main meals and snacks), energy intake, physical activity and sedentary habits were compared using second-order Rao–Scott corrections to the Pearson χ^2 test for categorical variables and by generalized linear models with Gauss family and identity link function for continuous variables. The number of eating occasions per day was categorized as ≤ 5 , > 5 – < 7 and ≥ 7 and the number of snacks per day was categorized as < 3 and ≥ 3 .



The probability of a child having a specific food consumption occasion was calculated using two trials binomial regression with logit link function.

To evaluate the association between eating frequency and overweight and obesity, weighted logistic regressions were used, obtaining crude and adjusted OR and respective 95% CI. The significance of coefficients was estimated using the Wald test. The following potential confounders were tested: children's sex and age, maternal age and education, maximum level of parents' education, NUTS II, child's practice of sports and sedentary lifestyle. To select which of the above variables to include in the models we used the change-in-estimate criterion with a cut-off point of 10% and/or the statistical significance of the variable for the model. Only children's sex and age (as a quadratic term) were significant to the models and included as confounder variables (model 2). To assess the effect of eating frequency on obesity independently of increasing energy intake, we also adjusted for the child's energy intake averaged from the two days (model 3). Interactions between children's sex, sport and maternal education were also tested. Tests for interactions were not significant and therefore we present results for all children together.

Statistical analyses were performed using R software version 3.4.1 for Windows and library 'survey'⁽³⁹⁾. A significance level of 5% was assumed.

Results

Table 1 presents characteristics of the children, for the total sample and according to weight status (no overweight *v.* overweight or obese). Only three children were classified as underweight and were added to the group of normal-weight children. Compared with normal-weight children, overweight/obese children were older on average and the majority were girls (55 *v.* 46%). The number of daily eating occasions ranged from 3.5 to 11, and on average children had 5.7 (95% CI 5.5, 5.8) eating occasions per day. In the overweight/obese children, the mean number of daily eating occasions (5.5 (95% CI 5.2, 5.9)) was lower than in the normal-weight children (5.8 (95% CI 5.6, 5.9)), although the difference was not statistically significant. The number of daily main meals was similar in both groups, but normal-weight children (mean = 2.8 (95% CI 2.7, 2.9)) had on average more snacks than the overweight children (mean = 2.6 (95% CI 2.2, 2.9)), although the difference was not statistically significant. The numbers of daily eating occasions and snacks were categorized; for the inclusion of these categorical variables in the regression models, the 'reference category' was considered as the mean of daily frequency: the middle category in the overall eating occasions (> 5–< 7) and the second category in the snacks frequency (≥ 3). The majority of overweight children had < 3 snacks/d, compared with normal-weight

children (66.9 (95% CI 54.4, 79.4)% *v.* 48.9 (95% CI 42.1, 55.7)%; $P=0.024$). On average, overweight children had a higher daily energy intake than normal-weight children (7590 (95% CI 7084, 8100) kJ (1814 (95% CI 1693, 1936) kcal) *v.* 6958 (95% CI 6724, 7196) kJ (1663 (95% CI 1607, 1720) kcal)). Normal-weight children had younger mothers and higher level of parents' education than did the overweight/obese children. However, normal-weight children practised sports less frequently and watched television/played video games less than the overweight/obese group. Regarding the misreporting evaluation, 93% of the sample had plausible reports.

Table 2 shows the percentage of children having eating occasions during the two recall days by food consumption occasion. For the three main meals, the percentage of children who consumed meals during both days was 95, 97 and 96% for breakfast, lunch and dinner, respectively. About 90% of the children did not have a meal before breakfast, before or after lunch, and before or after dinner. These results were similar across normal-weight and overweight/obese children. The average time of day that each food consumption occasion occurred is shown in Fig. 1. Usually, children had breakfast between 07.30 and 09.30 hours, lunch was reported as early as 12.00 hours until 13.30 hours, and dinner between 19.30 and 20.45 hours. The period of time when usually meals occurred did not seem to be different between underweight/normal-weight and overweight/obese children.

Energy intake increased significantly with increasing eating occasions and snacks in all children and in normal-weight children. The overweight and obese children had an average energy intake higher than the non-overweight children (Table 3).

Table 4 presents the results for the association between eating frequency and weight status. Considering the total sample, and after adjustment for child's sex, age and total energy intake, having ≤ 5 eating occasions/d was positively associated with being overweight or obese (OR = 1.93; 95% CI 1.00, 3.73). Having < 3 snacks/d, compared with having ≥ 3 snacks/d, was also positively associated with overweight or obesity (OR = 2.17; 95% CI 1.13, 4.16; Table 4). Due to the low number of misreporting reporters, we decided to perform the regression analysis excluding these children ($n=33$). Including only plausible reporters for energy intake, the association remained significant only for snack frequency and overweight/obesity (OR = 1.98; 95% CI 1.00, 3.90, < 3 snacks/d *v.* ≥ 3 snacks/d).

Discussion

Portuguese children aged 3–9 years had on average six daily eating occasions: three main meals and three snacks.

**Table 1** Characteristics of the 3–9-year-old Portuguese children, for the total sample and according to BMI category*, weighted for the Portuguese population distribution

	All children (n 517)		Normal weight (n 375)		Overweight/obese (n 142)		P value
	n, mean or %	% or 95 % CI	n, mean or %	% or 95 % CI	n, mean or %	% or 95 % CI	
Estimated population, n and %	630 731	100.0	436 241	69.2	194 338	30.8	
Age (years), mean and 95 % CI	5.8	5.5, 6.0	5.4	5.1, 5.7	6.5	6.3, 6.8	<0.001
Sex (girls), % and 95 % CI	49.1	45.7, 52.5	46.4	41.5, 51.3	55.1	46.7, 63.5	0.136
Eating occasions (frequency per day), mean and 95 % CI	5.7	5.5, 5.8	5.8	5.6, 5.9	5.5	5.2, 5.9	0.205
Main meals (frequency per day), mean and 95 % CI	2.9	2.9, 3.0	2.9	2.9, 3.0	2.9	2.9, 3.0	0.720
Snacks (frequency per day), mean and 95 % CI	2.7	2.6, 2.9	2.8	2.7, 2.9	2.6	2.2, 2.9	0.215
Eating occasions per day, % and 95 % CI							
≤5	35.0	29.2, 41.0	31.4	24.9, 37.9	43.6	30.6, 56.5	
>5–<7	49.9	43.4, 56.4	53.8	45.9, 61.8	40.9	29.4, 52.4	
≥7	15.0	10.6, 19.4	14.8	9.7, 19.8	15.5	7.0, 24.1	0.166
Snacks per day, % and 95 % CI							
<3	54.4	48.4, 60.5	48.9	42.1, 55.7	66.9	54.4, 79.4	
≥3	45.6	39.5, 51.6	51.1	44.3, 57.9	33.1	20.6, 45.6	0.024
Energy intake (kJ/d), mean and 95 % CI	7155	6925, 7381	6958	6724, 7196	7590	7084, 8100	0.031
Energy intake (kcal/d), mean and 95 % CI	1710	1655, 1764	1663	1607, 1720	1814	1693, 1936	0.031
Mother's age (years), mean and 95 % CI	37.2	36.5, 37.9	36.8	36.1, 37.5	38.0	36.8, 39.4	0.091
Level maximum of parents' education, % and 95 % CI							
No formal education or primary education	8.1	4.8, 11.4	7.7	3.8, 11.5	9.2	3.0, 15.4	
Secondary or post-secondary education (but non-tertiary education)	47.4	49.9, 53.8	46.7	39.5, 53.9	48.8	37.9, 59.7	
Tertiary education	44.5	37.0, 52.0	45.6	37.9, 53.3	42.0	29.5, 54.5	0.814
Physical activity practice, % and 95 % CI							
Yes	61.1	53.5, 68.6	59.0	51.2, 66.8	65.8	53.4, 78.2	0.299
Watching television/playing video games, % and 95 % CI							
<2 h/d	47.1	40.1, 54.0	50.0	42.1, 57.8	40.4	28.4, 52.4	
>2–<3 h/d	28.6	22.7, 34.6	27.4	20.4, 34.4	31.5	20.2, 42.8	
≥3 h/d	24.3	18.1, 30.4	22.6	15.8, 29.4	28.1	15.4, 40.8	0.448
Misreporting							
Plausible	93.4	90.2, 96.6	93.4	89.8, 97.1	93.3	86.6, 100.0	
Under-report	1.4	0.0, 3.4	0.3	0.0, 0.7	4.0	0.0, 10.0	
Over-report	5.2	2.5, 7.9	6.3	2.6, 9.9	2.7	0.0, 6.0	0.013

P values are determined by generalized linear models with Gauss family and identity link function for continuous variables and by second-order Rao–Scott corrections to the Pearson χ^2 test for categorical variables.

*BMI categories defined according to the WHO criteria⁽³³⁾.

The distribution of the meal schedule throughout the day had a higher variation for 'before breakfast', 'breakfast', 'during afternoon' and 'before dinner'. Overweight or obese children had on average fewer eating occasions than normal-weight children. Among these children, having five or fewer eating occasions, and particularly having fewer than three daily snacks, increased the odds of having excessive weight, after adjustment for potential confounders. Taking into consideration misreporting, the positive association with body weight was found only for having fewer than three daily snacks.

We opted for not analysing main meals separated from snacks, as the variability was low and the majority of children had the three main meals. So our analyses included all eating occasions and then separated by daily snacks.

Several pathways might explain the association between a lower eating frequency and higher body weight.

Theoretically, eating frequency could modify metabolism, increasing satiety and improving glucose and insulin metabolism^(40,41). Controlled feeding studies in adults⁽⁴⁰⁾ showed that reduced eating frequency had a negative impact on appetite control, increasing perceived appetite and reducing perceived satiety. In a randomized crossover trial among normal-weight women, meal regularity was associated with a greater thermic effect of food and lower glucose response, which might benefit weight management and metabolic health⁽⁴²⁾. Moreover, increased eating frequency has been associated with higher levels of physical activity and improved composition of snacks and breakfast, in terms of fruit and vegetables, fat, fibre and carbohydrates, in healthy-weight children aged 9–10 years⁽²⁵⁾.

Obesity is a complex disorder and some remaining uncontrolled confounding may not be ruled out. However, we tested several potential confounder

Table 2 Percentage of the 3–9-year-old Portuguese children having eating occasions by specific food consumption occasions (FCO), for the total sample and according to BMI category*, weighted for the Portuguese population distribution

	Before breakfast	Breakfast	During morning	Before lunch	Lunch	After lunch	During afternoon	Before dinner	Dinner	After dinner	During evening
All children											
Probability of having the FCO											
None of the days (%)	92.81	0.00	20.92	95.28	0.00	96.40	0.03	87.81	0.12	93.06	52.99
One of the days (%)	4.94	4.87	29.74	4.69	2.58	3.52	6.67	10.62	4.11	6.57	30.87
Both days (%)	2.25	95.13	49.35	0.03	97.42	0.08	93.30	1.57	95.76	0.38	16.14
Normal weight											
Probability of having the FCO											
None of the days (%)	92.58	0.00	18.01	94.61	0.00	97.31	0.05	84.11	0.18	91.64	53.51
One of the days (%)	4.35	5.78	32.54	5.34	1.61	2.58	5.12	13.78	3.25	7.88	28.81
Both days (%)	3.07	94.22	49.45	0.05	98.39	0.11	94.83	2.11	96.57	0.48	17.68
Overweight/obese											
Probability of having the FCO											
None of the days (%)	93.34	0.00	27.44	96.76	0.00	94.37	0.00	96.09	0.00	96.23	51.83
One of the days (%)	6.25	2.83	23.46	3.24	4.75	5.63	10.14	3.54	6.05	3.63	35.49
Both days (%)	0.41	97.17	49.11	0.00	95.25	0.00	89.86	0.37	93.95	0.14	12.68

*BMI categories defined according to the WHO criteria⁽³³⁾.




Fig. 1 Distribution of time of consumption (mean, with standard deviation represented by horizontal bars), in hours, for each of the food consumption occasions among 3–9-year-old Portuguese children, weighted for the Portuguese population distribution

Table 3 Mean energy intake across categories of eating occasions and snack events among 3–9-year-old Portuguese children, for the total sample and according to BMI category*, weighted for the Portuguese population distribution

	All children		Normal weight		Overweight/obese	
	Energy intake (kJ/d)	Energy intake (kcal/d)	Energy intake (kJ/d)	Energy intake (kcal/d)	Energy intake (kJ/d)	Energy intake (kcal/d)
	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI
Eating occasions, per day						
≤5	6506	6104, 6904	1555	1459, 1650	6184	5820, 6544
>5–<7	7351	7084, 7619	1757	1693, 1821	7159	6849, 7464
≥7	8029	7418, 8640	1919	1773, 2065	7891	7209, 8577
P value	<0.001		<0.001		<0.001	
Snacks, per day						
<3	6845	6489, 7205	1636	1551, 1722	6498	6188, 6807
≥3	7523	7213, 7837	1798	1724, 1873	7401	7012, 7791
P value	0.011		0.011		0.002	

P values were calculated with the likelihood ratio test with Rao–Scott corrections and the F test.

*BMI categories defined according to the WHO criteria⁽³³⁾.

**Table 4** Association between eating frequency (daily eating occasions and daily snacks) and overweight and obesity, in a national representative sample of children aged 3–9 years of age, weighted for the Portuguese population distribution, for the total sample (*n* 517) and excluding misreporters (*n* 484)

	OR for overweight/obesity (95 % CI)											
	All children (<i>n</i> 517)						Only plausible reporters (<i>n</i> 484)					
	Model 1 (unadjusted)		Model 2		Model 3		Model 1 (unadjusted)		Model 2		Model 3	
	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI
Eating occasions, per day												
≤5	1.83	0.95, 3.51	1.78	0.90, 3.52	1.93	1.00, 3.73	1.73	0.89, 3.35	1.64	0.83, 3.24	1.91	0.97, 3.77
>5–<7	Reference		Reference		Reference		Reference		Reference		Reference	
≥7	1.38	0.63, 3.04	1.48	0.68, 3.22	1.40	0.61, 3.23	1.66	0.73, 3.79	1.88	0.84, 4.22	1.40	0.67, 4.05
Snacks, per day												
<3	2.11	1.11, 4.04	2.00	1.05, 3.80	2.17	1.13, 4.16	1.88	0.95, 3.69	1.72	0.88, 3.35	1.98	1.00, 3.90
≥3	Reference		Reference		Reference		Reference		Reference		Reference	

Model 1, unadjusted; model 2, adjusted for child's sex and age (quadratic term); model 3, adjusted for child's sex and age (quadratic term) and daily children's energy intake (kcal).

Bold entries denote statistical significance.

variables, namely children's sex and age, maternal age and education, parental education, geographic region, child's sports practice and sedentary lifestyle. Only children's sex and age remained significant in the models and were included as confounder variables, as well as children's energy intake (averaged from the two days). Interactions between children's sex, sport and maternal education were also tested but no modifying effect was found.

Irregular consumption of energy intake at breakfast and between meals was previously associated with a higher cardiometabolic risk in adults, including an increased waist circumference and BMI, in a cross-sectional study⁽¹⁶⁾; another cross-sectional study showed that in children larger variability in eating frequency was already associated with a disruption in total and LDL cholesterol concentrations⁽⁴³⁾. Although we did not assess these effects (since we evaluated dietary intake of only two days) in the present study, we were successful in finding a significant association between eating frequency, namely snacks frequency, and BMI in children.

A previous meta-analysis⁽²¹⁾, that included ten cross-sectional studies and one case–control study, among children and adolescents, described an overall inverse association between eating frequency and the likelihood of being overweight or obese. However, when stratified by sex, the effect remained significant only in boys. In the present study, we tested an interaction effect between sex and eating frequency in the association with obesity; because we did not find a statistically significant effect we decided to present the results for the entire sample, and not stratified by sex.

More recently, a cross-sectional study⁽⁴⁴⁾ failed to find any association between eating frequency and BMI in UK children aged 4–10 years, while another cross-sectional study⁽²⁵⁾ found an inverse association in healthy-weight children aged 9–10 years, but not in centrally obese

children. In a longitudinal study⁽⁴⁵⁾ in very young children, the daily eating frequency was not associated with the current or subsequent change in BMI. The authors highlighted that these null associations might be due to an improvement of internal energy self-regulation by young children, in comparison with older children⁽⁴⁵⁾.

Methodological differences between studies might explain the discrepancy of results regarding the association between eating frequency and weight. In the present study, the weight and height of the children were measured by trained examiners, following standard procedures. The BMI Z-scores were then calculated and categorized according to the WHO criteria. In the majority of previous research, children were categorized as overweight or obese according to other criteria, such as the International Obesity Task Force, hampering comparisons between different studies⁽²¹⁾.

There is also variability in the dietary assessment method and in the definition of eating frequency, meal and snack frequency. Food diaries, 24 h dietary recalls and self-reported questionnaires (by children or by their caregivers) have been used to assess eating frequency. In the present study we defined eating occasions using the 'participant-identified' approach, as it was the child's caregiver who decided the food consumption occasions in the food diaries, but it was the researchers who categorized the food consumption occasions, that provided at least 209 kJ (50 kcal), into main meals (breakfast, lunch and dinner) and snacks (the remaining food consumption occasions). We were also able to identify the time of day when children usually had each of the food consumption occasions.

Furthermore, different studies used different cut-offs to determine high or low eating frequency or using the exposure as a continuous variable. This last approach might not be the best one as we believe the association between



eating frequency and obesity is not linear, and the extremes are the ones most harmful; correspondingly, we used the middle category as the reference in the eating occasion variable.

The strengths of the present study include a national representative sample of children and results weighted for the Portuguese population distribution regarding sex, age and region, following harmonized European methodology⁽²⁸⁾. We also used two non-consecutive one-day food diaries followed by a face-to-face interview to estimate children's eating frequency, which is an advantage in comparison to simply asking parents to specify the usual eating frequency of their child. Trained interviewers, following standard procedures, performed objective measures of children's body size, which is also a strength compared with parents reporting the weight and height of the children.

There are also some limitations that deserve discussion. First, under-reporting of dietary intake among overweight or obese people is widely described in the literature^(46,47) and might have an impact on the association between eating frequency and obesity. When we included only plausible reporters for energy intake, the association between daily eating occasions and body weight ceased to be statistically significant. This effect might be due to a loss of power, as we excluded some children, and not due to a bias regarding the dietary report. Literature has reported a mismatch in mothers' perception regarding their children's weight. For example, in a cohort study among children aged 6–8 years, the majority of mothers misclassified the child's weight and had the tendency to normalize the child's weight status⁽⁴⁸⁾. Mothers were also poor at recognizing overweight and obesity in their children at 4 or 6 years⁽⁴⁹⁾ or in pre-school children⁽⁵⁰⁾. As parents might fail to recognize their child as overweight or obese, it is not expected that they change their report based on the child's weight status. A residual confounding because of the use of a broad categorization of physical activity level cannot be ruled out. However, in a sensitivity analysis, we included information for physical activity level evaluated by diaries for children aged 6 years or above, and for younger children, we have assumed the values defined according to Torun⁽³⁸⁾. This variable was tested as a confounder in the association between eating frequency and children's overweight/obesity, but it did not change the estimate by 10% nor was it significant.

A social desirability bias might have occurred, resulting in an over-reporting of healthy foods and/or under-reporting of unhealthy foods. However, this bias might have been minimized as our main exposure variable, eating frequency, was calculated independently of the quality or types of foods.

Another limitation is the cross-sectional nature of the study, which does not allow assessing the temporal relationship between eating frequency and obesity. Moreover, we could not exclude a potential reverse causality effect, since obese people could change their frequency of meals in an

attempt to lose weight. However, this approach is more likely to occur in adults than in children⁽⁵¹⁾.

Conclusion

Portuguese children aged 3–9 years had on average six daily eating occasions, divided into three main meals and three snacks. In these children, and after adjustment for potential confounders, lower daily snack frequency (<3/d) was positively associated with being overweight or obese. The present study supports that increasing the number of daily meals, maintaining the same energy intake, contributes to a healthy body weight in children.

Acknowledgements

This study had institutional support from the General Directorate of Health (DGS), the Regional Health Administration Departments, the Central Administration of the Health System (ACSS) and the European Food Safety Authority (CFT/EFSA/DCM/2012/01-C03). The researchers acknowledge all these institutions and persons involved in all phases of the IAN-AF 2015–2016, as well as the participants. *Financial support:* The IAN-AF 2015–2016 received funding from the EEA Grants Program, Public Health Initiatives (grant number PT06-000088SI3). The EEA Grant Program had no role in the design, analysis or writing of this article. *Conflict of interest:* None. *Authorship:* S.V. contributed to the design of the study, interpretation of the data and wrote the first draft of the paper. D.C. and M.S. contributed the statistical analysis and to the interpretation of data. A.O., C.L. and D.T. contributed to the design of the study, coordination of data collection and contributed to the discussion of results. All authors critically reviewed the manuscript and approved the final version as submitted. *Ethics of human subject participation:* Ethical approval was obtained from the National Commission for Data Protection, the Ethical Committee of the Institute of Public Health of the University of Porto and from the Ethical Commissions of each one of the Regional Administrations of Health. Written agreements from the parents or legal representatives were required for children. All documents with identification data were treated separately and stored in a different data set.

References

1. World Health Organization (2017) Obesity and overweight. <http://www.who.int/mediacentre/factsheets/fs311/en/> (accessed October 2017).
2. de Onis M, Blossner M & Borghi E (2010) Global prevalence and trends of overweight and obesity among preschool children. *Am J Clin Nutr* **92**, 1257–1264.
3. Adair L (2008) Child and adolescent obesity: epidemiology and developmental perspectives. *Physiol Behav* **94**, 8–16.



4. Reilly JJ (2005) Descriptive epidemiology and health consequences of childhood obesity. *Best Pract Res Clin Endocrinol Metab* **19**, 327–341.
5. Cornette R (2008) The emotional impact of obesity on children. *Worldviews Evid Based Nurs* **5**, 136–141.
6. Field AE, Camargo CA Jr, Taylor CB *et al.* (1999) Overweight, weight concerns, and bulimic behaviors among girls and boys. *J Am Acad Child Adolesc Psychiatry* **38**, 754–760.
7. Reinehr T, Andler W, Denzer C *et al.* (2005) Cardiovascular risk factors in overweight German children and adolescents: relation to gender, age and degree of overweight. *Nutr Metab Cardiovasc Dis* **15**, 181–187.
8. Visser M, Bouter LM, McQuillan GM *et al.* (2001) Low-grade systemic inflammation in overweight children. *Pediatrics* **107**, E13.
9. Pena AS, Wiltshire E, MacKenzie K *et al.* (2006) Vascular endothelial and smooth muscle function relates to body mass index and glucose in obese and nonobese children. *J Clin Endocrinol Metab* **91**, 4467–4471.
10. Meyer AA, Kundt G, Steiner M *et al.* (2006) Impaired flow-mediated vasodilation, carotid artery intima-media thickening, and elevated endothelial plasma markers in obese children: the impact of cardiovascular risk factors. *Pediatrics* **117**, 1560–1567.
11. Chiarelli F & Marcovecchio ML (2008) Insulin resistance and obesity in childhood. *Eur J Endocrinol* **159**, Suppl. 1, S67–S74.
12. Gardner DS, Hosking J, Metcalf BS *et al.* (2009) Contribution of early weight gain to childhood overweight and metabolic health: a longitudinal study (EarlyBird 36). *Pediatrics* **123**, e67–e73.
13. Ramsay SA, Bloch TD, Marriage B *et al.* (2018) Skipping breakfast is associated with lower diet quality in young US children. *Eur J Clin Nutr* **72**, 548–556.
14. Deshmukh-Taskar PR, Nicklas TA, O'Neil CE *et al.* (2010) The relationship of breakfast skipping and type of breakfast consumption with nutrient intake and weight status in children and adolescents: the National Health and Nutrition Examination Survey 1999–2006. *J Am Diet Assoc* **110**, 869–878.
15. Deshmukh-Taskar P, Nicklas TA, Radcliffe JD *et al.* (2013) The relationship of breakfast skipping and type of breakfast consumed with overweight/obesity, abdominal obesity, other cardiometabolic risk factors and the metabolic syndrome in young adults. The National Health and Nutrition Examination Survey (NHANES): 1999–2006. *Public Health Nutr* **16**, 2073–2082.
16. Pot GK, Hardy R & Stephen AM (2014) Irregular consumption of energy intake in meals is associated with a higher cardiometabolic risk in adults of a British birth cohort. *Int J Obes (Lond)* **38**, 1518–1524.
17. Mills JP, Perry CD & Reicks M (2011) Eating frequency is associated with energy intake but not obesity in midlife women. *Obesity (Silver Spring)* **19**, 552–559.
18. Howarth NC, Huang TT, Roberts SB *et al.* (2007) Eating patterns and dietary composition in relation to BMI in younger and older adults. *Int J Obes (Lond)* **31**, 675–684.
19. van der Heijden AA, Hu FB, Rimm EB *et al.* (2007) A prospective study of breakfast consumption and weight gain among US men. *Obesity (Silver Spring)* **15**, 2463–2469.
20. Canuto R, da Silva Garcez A, Kac G *et al.* (2017) Eating frequency and weight and body composition: a systematic review of observational studies. *Public Health Nutr* **20**, 2079–2095.
21. Kaisari P, Yannakoulia M & Panagiotakos DB (2013) Eating frequency and overweight and obesity in children and adolescents: a meta-analysis. *Pediatrics* **131**, 958–967.
22. Mesas AE, Munoz-Pareja M, Lopez-Garcia E *et al.* (2012) Selected eating behaviours and excess body weight: a systematic review. *Obes Rev* **13**, 106–135.
23. Leech RM, Worsley A, Timperio A *et al.* (2015) Understanding meal patterns: definitions, methodology and impact on nutrient intake and diet quality. *Nutr Res Rev* **28**, 1–21.
24. Johnson GH & Anderson GH (2010) Snacking definitions: impact on interpretation of the literature and dietary recommendations. *Crit Rev Food Sci Nutr* **50**, 848–871.
25. Jennings A, Cassidy A, van Sluijs EM *et al.* (2012) Associations between eating frequency, adiposity, diet, and activity in 9–10 year old healthy-weight and centrally obese children. *Obesity (Silver Spring)* **20**, 1462–1468.
26. Berg C & Forslund HB (2015) The influence of portion size and timing of meals on weight balance and obesity. *Curr Obes Rep* **4**, 11–18.
27. Lopes C, Torres D, Oliveira A *et al.* (2018) National Food, Nutrition, and Physical Activity Survey of the Portuguese general population (2015–2016): protocol for design and development. *JMIR Res Protoc* **7**, e42.
28. European Food Safety Authority (2014) Guidance on the EU Menu methodology. *EFSA J* **12**, 3944–3977.
29. Torres D, Faria N, Sousa N *et al.* (2017) *Inquérito Alimentar Nacional e de Atividade Física, IAN-AF 2015–2016: Manual Fotográfico de Quantificação de Alimentos*. Porto: Universidade do Porto.
30. Vilela S, Lopes C, Guiomar S *et al.* (2018) Validation of a picture book to be used in a pan-European dietary survey. *Public Health Nutr* **21**, 1654–1663.
31. National Institute of Health Dr Ricardo Jorge (2006) *Food Composition Table* (in Portuguese). Lisbon: Center of Food Safety and Nutrition.
32. Stewart A & Marfell-Jones M (2001) International Standards for Anthropometric Assessment. International Society for the Advancement of Kinanthropometry. <http://www.ceap.br/material/MAT17032011184632.pdf> (accessed February 2018).
33. WHO Multicentre Growth Reference Study Group (2006) *WHO Child Growth Standards: Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index-for-Age: Methods and Development*. Geneva: WHO.
34. Tremblay MS, LeBlanc AG, Kho ME *et al.* (2011) Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *Int J Behav Nutr Phys Act* **8**, 98.
35. Bouchard C, Tremblay A, Leblanc C *et al.* (1983) A method to assess energy expenditure in children and adults. *Am J Clin Nutr* **37**, 461–467.
36. Goldberg GR, Black AE, Jebb SA *et al.* (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* **45**, 569–581.
37. Schofield WN (1985) Predicting basal metabolic rate, new standards and review of previous work. *Hum Nutr Clin Nutr* **39**, Suppl. 1, 5–41.
38. Torun B (2005) Energy requirements of children and adolescents. *Public Health Nutr* **8**, 968–993.
39. Lumley T (2004) Analysis of complex survey samples. *J Stat Softw* **9**, 1–19.
40. Leidy HJ & Campbell WW (2011) The effect of eating frequency on appetite control and food intake: brief synopsis of controlled feeding studies. *J Nutr* **141**, 154–157.
41. Farshchi HR, Taylor MA & Macdonald IA (2005) Beneficial metabolic effects of regular meal frequency on dietary thermogenesis, insulin sensitivity, and fasting lipid profiles in healthy obese women. *Am J Clin Nutr* **81**, 16–24.
42. Alhussain MH, Macdonald IA & Taylor MA (2016) Irregular meal-pattern effects on energy expenditure, metabolism, and appetite regulation: a randomized controlled trial in healthy normal-weight women. *Am J Clin Nutr* **104**, 21–32.



43. Murakami K & Livingstone MB (2015) Variability in eating frequency in relation to adiposity measures and blood lipid profiles in British children and adolescents: findings from the National Diet and Nutrition Survey. *Int J Obes (Lond)* **39**, 608–613.
44. Murakami K & Livingstone MB (2014) Associations of eating frequency with adiposity measures, blood lipid profiles and blood pressure in British children and adolescents. *Br J Nutr* **111**, 2176–2183.
45. Taylor RW, Iosua E, Heath AM *et al.* (2017) Eating frequency in relation to BMI in very young children: a longitudinal analysis. *Public Health Nutr* **20**, 1372–1379.
46. Wehling H & Lusher J (2017) People with a body mass index 30 under-report their dietary intake: a systematic review. *J Health Psychol*. Published online: 1 July 2017. doi: 10.1177/1359105317714318.
47. Murakami K & Livingstone MB (2015) Prevalence and characteristics of misreporting of energy intake in US adults: NHANES 2003–2012. *Br J Nutr* **114**, 1294–1303.
48. McDonald SW, Ginez HK, Vinturache AE *et al.* (2016) Maternal perceptions of underweight and overweight for 6–8 years olds from a Canadian cohort: reporting weights, concerns and conversations with healthcare providers. *BMJ Open* **6**, e012094.
49. Mitchell R, Wake M, Canterford L *et al.* (2008) Does maternal concern about children's weight affect children's body size perception at the age of 6.5? – A community-based study. *Int J Obes (Lond)* **32**, 1001–1007.
50. Campbell MW, Williams J, Hampton A *et al.* (2006) Maternal concern and perceptions of overweight in Australian preschool-aged children. *Med J Aust* **184**, 274–277.
51. Rietmeijer-Mentink M, Paulis WD, van Middelkoop M *et al.* (2013) Difference between parental perception and actual weight status of children: a systematic review. *Matern Child Nutr* **9**, 3–22.

Appendix: Members of the IAN-AF Consortium

Carla Lopes, Andreia Oliveira, Milton Severo (Faculty of Medicine, University of Porto); Duarte Torres, Sara Rodrigues (Faculty of Nutrition and Food Sciences, University of Porto); Elisabete Ramos, Sofia Vilela (EPIUnit, Institute of Public Health, University of Porto); Sofia Guiomar, Luísa Oliveira (National Health Institute Doutor Ricardo Jorge); Violeta Alarcão, Paulo Nicola (Institute of Preventive Medicine and Public Health, Faculty of Medicine, University of Lisbon); Jorge Mota (CIAFEL, Faculty of Sports, University of Porto); Pedro Teixeira (Faculty of Human Kinetics, CIPER, University of Lisbon); Simão Soares (SilicoLife, Lda, Portugal); Lene Frost Andersen (Faculty of Medicine, University of Oslo).