

Letter to the Editor

Comment on Sergeant *et al.*: Impact of methods used to express levels of circulating fatty acids on the degree and direction of associations with blood lipids in humans

(First published online 15 April 2016)

In their paper in the January issue of this journal, Sergeant *et al.*⁽¹⁾, reported on the impact of different methods to express circulating fatty acids on the association with blood lipids. Using two human cohorts, the authors showed that the direction of the correlation between a number of individual circulating fatty acids and TAG reversed when expressing the circulating fatty acids as an absolute concentration *v.* a proportion of total fatty acids. On the basis of their findings, the authors suggest that expressing circulating fatty acids as concentrations is superior to expressing them as proportions, and they concluded that earlier relationships between (dietary) fatty acids and other biomarkers and indices of CHD should be re-visited with this in mind.

With their paper, Sergeant *et al.* point at an unresolved question on how to express and interpret levels of circulating fatty acids^(2,3). Both methods to express levels of circulating fatty acids have their merits and disadvantages. However, and contrary to the authors' suggestion, the reversal of the association with blood lipids when using different methods to express levels of circulating fatty acids does in itself not provide sufficient arguments for favouring one way of expression above the other. In fact, several key arguments in favour of expressing circulating fatty acids as a proportion of total fatty acids were not addressed.

First, circulating fatty acids expressed as proportions better reflect dietary intake than when expressed as concentrations, particularly for PUFA⁽²⁾. This is supported by data from our Hoom⁽⁴⁾ and Cohort Study on Diabetes and Atherosclerosis Maastricht (CODAM)⁽⁵⁾ cohorts. In 1171 subjects, intakes of dietary fatty acids were measured by a FFQ and circulating fatty acids by GLC. The correlation between dietary linoleic acid (LA) and proportions of circulating LA was r 0.306, whereas the correlation between dietary LA and concentrations of LA ($\mu\text{g/ml}$) was r 0.175. Higher correlations when expressed in proportions were observed across PUFA.

Second, the pool of total circulating fatty acids comprises all individual circulating fatty acids, mainly present as TAG, cholesterol esters and phospholipids. Therefore, the concentration of individual circulating fatty acids is positively related to the concentration of TAG and LDL-cholesterol, HDL-cholesterol and total cholesterol (TC), as shown by Sergeant *et al.* and others^(1,2). Consequently, using circulating fatty acids as concentrations to study the relationship between intakes of dietary fatty acids and blood lipids can be confounded by effects of the individual dietary fatty acid on concentrations

of the blood lipids itself, and may lead to spurious conclusions. In fact, strictly controlled dietary intervention studies on blood lipids provide strong evidence that increasing the intakes of dietary PUFA (mostly LA) at the expense of carbohydrates and SFA lowers TAG, LDL-cholesterol and TC concentrations⁽⁶⁾.

We observe similar discrepancy in the associations of the concentration of TAG with circulating LA expressed in proportions (r -0.448) or concentrations (r 0.395), as shown by Sergeant *et al.*⁽¹⁾. It is, however, important to note that this discrepancy may be specific to the association of circulating fatty acids with blood lipids, as circulating fatty acids are part of the pool of these blood lipids. The question whether such a discrepancy exists in the associations of circulating fatty acids with other CVD risk markers such as circulating cytokines or with the incidence of CVD or other outcomes is unresolved and deserves to be explored separately. For example, a strong association of DHA and EPA with brain atrophy rates has recently been reported, which was similarly inverse when circulating fatty acids were expressed as absolute concentrations or as proportions⁽⁷⁾.

It can be argued that, when circulating fatty acids are used as a marker of dietary intake, the most important criterion for evaluating different methods of expressing circulating fatty acids is the strength of the relationship with their levels of dietary intake. This favours the expression of circulating fatty acids as a proportion of total fatty acids rather than as absolute concentrations, particularly for PUFA. However, our observations, combined with the findings of Sergeant *et al.*⁽¹⁾ and others^(2,7), indicate that the optimal method of expressing fatty acid status depends on the biological assumptions and the biological question at hand.

Acknowledgements

A. J. W., M. A., M. J. V. G., A. E., P. L. Z., J. M. D. and I. A. B. contributed to the writing of the letter; A. J. W. and M. A. coordinated the process.

Funding: all authors contributed independently to this work.

Conflict of interest: A. J. W., M. A., P. L. Z. are employed by Unilever; Unilever markets food products made of vegetable oils, including margarines and dressings. No other authors declared a conflict of interest.

Anne J. Wanders¹, Marjan Alsema^{1,2,3},
Marleen J. van Greevenbroek⁴, Amany Elshorbagy⁵,
Peter L. Zock¹, Jacqueline M. Dekker^{2,3} and
Ingeborg A. Brouwer^{2,6}

¹*Unilever Research and Development, Olivier van
Noortlaan 120, 3133 AT Vlaardingen,
The Netherlands*

²*EMGO Institute for Health and Care Research,
VU University Medical Center, Van der Boechorststraat 7,
1081 BT Amsterdam, The Netherlands*

³*Department of Epidemiology and Biostatistics,
VU University Medical Center, De Boelelaan 1117,
1007 MB Amsterdam, The Netherlands*

⁴*Department of Internal Medicine, Maastricht University,
Universiteitssingel 50, 6229 ER Maastricht,
The Netherlands*

⁵*Department of Physiology, Faculty of Medicine,
Alexandria University, Chamblion Street, El Azareeta,
Alexandria, Egypt*

⁶*Department of Health Sciences, VU University
Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam,
The Netherlands*

email anne.wanders@unilever.com

doi:10.1017/S0007114516000933

References

1. Sergeant S, Ruczinski I, Ivester P, *et al.* (2016) Impact of methods used to express levels of circulating fatty acids on the degree and direction of associations with blood lipids in humans. *Br J Nutr* **115**, 251–261.
2. Bradbury KE, Skeaff CM, Green TJ, *et al.* (2010) The serum fatty acids myristic acid and linoleic acid are better predictors of serum cholesterol concentrations when measured as molecular percentages rather than as absolute concentrations. *Am J Clin Nutr* **91**, 398–405.
3. Hodson L, Skeaff CM & Fielding BA (2008) Fatty acid composition of adipose tissue and blood in humans and its use as a biomarker of dietary intake. *Prog Lipid Res* **47**, 348–380.
4. Reinders I, van Ballegooijen AJ, Visser M, *et al.* (2013) Associations of serum *n*-3 and *n*-6 polyunsaturated fatty acids with echocardiographic measures among older adults: the Hoorn study. *Eur J Clin Nutr* **67**, 1277–1283.
5. Van Woudenbergh GJ, Kuijsten A, Van der Kallen CJ, *et al.* (2012) Comparison of fatty acid proportions in serum cholesteryl esters among people with different glucose tolerance status: the CoDAM study. *Nutr Metab Cardiovasc Dis* **22**, 133–140.
6. Mensink RP, Zock PL, Kester AD, *et al.* (2003) Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. *Am J Clin Nutr* **77**, 1146–1155.
7. Jerneren F, Elshorbagy AK, Oulhaj A, *et al.* (2015) Brain atrophy in cognitively impaired elderly: the importance of long-chain omega-3 fatty acids and B vitamin status in a randomized controlled trial. *Am J Clin Nutr* **102**, 215–221.